


# Pythagorean theorem

In [mathematics](#), the **Pythagorean theorem** or **Pythagoras' theorem** is a fundamental relation in [Euclidean geometry](#) between the three sides of a [right triangle](#). It states that the area of the [square](#) whose side is the [hypotenuse](#) (the side opposite the [right angle](#)) is equal to the sum of the areas of the squares on the other two sides.

The [theorem](#) can be written as an [equation](#) relating the lengths of the sides  $a$ ,  $b$  and the hypotenuse  $c$ , sometimes called the **Pythagorean equation**:<sup>[1]</sup>

<div><div>Pythagorean theorem</div></div>	
Type	Theorem
Field	<a href="#">Euclidean geometry</a>
Statement	The sum of the areas of the two squares on the legs ( $a$ and $b$ ) equals the area of the square on the hypotenuse ( $c$ ).
Symbolic statement	$a^2 + b^2 = c^2$
Generalizations	<a href="#">Law of cosines</a> <a href="#">Solid geometry</a> <a href="#">Non-Euclidean geometry</a> <a href="#">Differential geometry</a>
Consequences	<a href="#">Pythagorean triple</a> <a href="#">Reciprocal Pythagorean theorem</a> <a href="#">Complex number</a> <a href="#">Euclidean distance</a> <a href="#">Pythagorean trigonometric identity</a>


$$a^2 + b^2 = c^2.$$

The theorem is named for the [Greek](#) philosopher [Pythagoras](#), born around 570 BC. The theorem has been [proved](#) numerous times by many different methods – possibly the most for any mathematical theorem. The proofs are diverse, including both [geometric](#) proofs and [algebraic](#) proofs, with some dating back thousands of years.

When [Euclidean space](#) is represented by a [Cartesian coordinate system](#) in [analytic geometry](#), [Euclidean distance](#) satisfies the Pythagorean relation: the squared distance between two points equals the sum of squares of the difference in each coordinate between the points.

The theorem can be [generalized](#) in various ways: to [higher-dimensional spaces](#), to [spaces that are not Euclidean](#), to objects that are not right triangles, and to objects that are not triangles at all but [n-dimensional](#) solids.

## Proofs using constructed squares



Rearrangement proof of the Pythagorean theorem.

(The area of the white space remains constant throughout the translation rearrangement of the triangles. At all moments in time, the area is always  $c^2$ . And likewise, at all moments in time, the area is always  $a^2 + b^2$ .)

### Rearrangement proofs

In one rearrangement proof, two squares are used whose sides have a measure of  $a + b$  and which contain four right triangles whose sides are  $a$ ,  $b$  and  $c$ , with the hypotenuse being  $c$ . In the square on the right side, the triangles are placed such that the corners of the square correspond to the corners of the right angle in the triangles, forming a square in the center whose sides are length  $c$ . Each outer square has an area of  $(a + b)^2$  as well as  $2ab + c^2$ , with  $2ab$  representing the total area of the four triangles. Within the big square on the left side, the four triangles are moved to form two similar rectangles with sides of length  $a$  and  $b$ . These rectangles in their new position have now delineated two new squares, one having side length  $a$  is formed in the bottom-left corner, and another square of side length  $b$  formed in the top-right corner. In this new position, this left side now has a square of area  $(a + b)^2$  as well as  $2ab + a^2 + b^2$ . Since both squares have the area of  $(a + b)^2$  it follows that the other measure of the square area also equal each other such that  $2ab + c^2 = 2ab + a^2 + b^2$ . With the area of the four triangles removed from both side of the equation what remains is  $a^2 + b^2 = c^2$ .<sup>[2]</sup>

In another proof rectangles in the second box can also be placed such that both have one corner that correspond to consecutive corners of the square. In this way they also form two boxes, this time in consecutive corners, with areas  $a^2$  and  $b^2$  which will again lead to a second square of with the area  $2ab + a^2 + b^2$ .

English mathematician [Sir Thomas Heath](#) gives this proof in his commentary on Proposition I.47 in [Euclid's Elements](#), and mentions the proposals of German mathematicians [Carl Anton Bretschneider](#) and [Hermann Hankel](#) that Pythagoras may have known this proof. Heath himself favors a different proposal for a Pythagorean proof, but acknowledges from the outset of his discussion "that the Greek literature which we possess belonging to the first five centuries after Pythagoras contains no statement specifying this or any other particular great geometric discovery to him."<sup>[3]</sup> Recent scholarship has cast increasing doubt on any sort of role for Pythagoras as a creator of mathematics, although debate about this continues.<sup>[4]</sup>

## Algebraic proofs




Diagram of the two algebraic proofs

The theorem can be proved algebraically using four copies of the same triangle arranged symmetrically around a square with side  $c$ , as shown in the lower part of the diagram.<sup>[5]</sup> This results in a larger square, with side  $a + b$  and area  $(a + b)^2$ . The four triangles and the square side  $c$  must have the same area as the larger square,