

REVIEW

Chronic Intermittent Hypoxia and Hypertension: A Review of Systemic Inflammation and Chinese Medicine*

WU Chun-xiao (邬春晓), LIU Yue (刘 玥), and ZHANG Jing-chun (张京春)

ABSTRACT Obstructive sleep apnea syndrome (OSAS) and hypertension commonly coexist. Clinical studies indicate that OSAS plays a key role in increasing the risk of prevalent hypertension. Chronic intermittent hypoxia (CIH) is the core pathological mechanism of OSAS, and has a close relationship with systemic inflammation. Growing evidence shows that CIH and hypertension are strongly related, involving markers or pathways indicative of systemic inflammation, such as high-sensitivity C-reactive protein (hs-CRP), interleukin-6, nuclear factor-kappa B, tumor necrosis factor- α , interleukin-8 and p38 mitogen-activated protein kinase (MAPK)-dependent pathways. Oxidative stress also plays an important role in this process, including in the activation of polymorphonuclear neutrophils. However, the pathophysiological and clinical significance of systemic inflammation in CIH and hypertension is not proven. This review article highlights the relationship between CIH and hypertension through systemic inflammation and the current interventions available in Chinese medicine, to offer a background for the future treatment of OSAS-related hypertension with integrative medicine.

KEYWORDS chronic intermittent hypoxia, obstructive sleep apnea syndrome, hypertension, systemic inflammation, Chinese medicine

Obstructive sleep apnea syndrome (OSAS) is a common disease, affecting about 2%–4% of the general population and characterized by repetitive upper airway collapse, frequent arousals and sleep disruption.⁽¹⁾ Data convincingly show a high correlation between OSAS and cardiovascular disorders such as hypertension, coronary artery disease, arrhythmia and stroke.⁽²⁾ Great strides have been made in establishing OSAS as an independent risk factor for hypertension.⁽³⁾ The high correlation between OSAS and hypertension was first discovered in the 1970s; scholars reported an improvement in blood pressure (BP) after surgical intervention for a variety of sleep disorder-related breathing problems.^(4–6) In the past several decades, a number of studies have found a relationship between OSAS and hypertension. Additionally, hypertension has been observed in 40% of OSAS patients. Meanwhile, OSAS is found in 30%–40% of patients with hypertension. These results indicate that OSAS and hypertension are intimately related.⁽⁷⁾ However, the mechanism of OSAS-related hypertension remains unclear. Because of upper airway obstruction, disordered breathing events lead to the intermittent occurrence of hypoxia throughout the sleep period. This phenomenon is called chronic intermittent hypoxia (CIH).⁽⁸⁾ CIH is different from continuous hypoxia. The former is distinguished by periods of low oxygen between periods of normal

oxygen. However, both of them independently lead to inflammation. While the underlying mechanisms of OSAS-related hypertension are poorly understood, systemic inflammation may play a key role in the process.

Indeed, the circulating levels of C-reactive protein (CRP), tumor necrosis factor- α (TNF- α), interleukin (IL)-6 and IL-8 are elevated in patients with OSAS,^(9–12) suggesting that CIH may be pivotal in systemic inflammation. Inflammation has been receiving a fair amount of attention as the etiology for the initiation and progression of hypertension. In brief, endothelial dysfunction develops at an early stage in response to the afore-mentioned inflammatory cytokines. This process leads to the up-regulation

©The Chinese Journal of Integrated Traditional and Western Medicine Press and Springer-Verlag Berlin Heidelberg 2013

*Supported by the Special Research Foundation of National Clinical Research Facility of Traditional Chinese Medicine (No. JDZX2012108); China Postdoctoral Foundation (No. 200902187); Famous Doctor Inheritance Research Foundation of China Academy of Chinese Medical Sciences (CM20121018); "3+3" Famous Doctor Inheritance Program of Beijing Administration of Traditional Chinese Medicine

Department of Cardiovascular Disease, Xiyuan Hospital, China Academy of Chinese Medical Sciences, Beijing (100091), China
Correspondence to: Prof. ZHANG Jing-chun, Tel: 86-10-62835341, E-mail: zhangjingchun276@vip.sohu.com

DOI: 10.1007/s11655-013-1459-x

of intercellular adhesion molecules, which promote leukocyte adherence to and extravasation into the endothelium.⁽¹³⁾ This is a potential mechanism leading to hypertension.

Few studies have evaluated the potential interactions among OSAS, hypertension and systemic inflammation in the development of these disorders. The objectives of this paper are, first, to review the correlation between systemic inflammation caused by CIH and the pathogenesis of hypertension, including biochemical/molecular mechanisms; and second, to review the progress of interventions with Chinese medicine in treating OSAS-hypertension. CM interventions could provide a useful alternative treatment to prevent cardiovascular complications after hypertension.

CIH and Inflammatory Activation Pathways

CIH and Nuclear Factor-Kappa B

Recent studies exploring the relationship between hypertension and OSAS have focused on the role of the transcription factor nuclear factor-kappa B (NF- κ B). Many studies have revealed the complex regulation by NF- κ B of genes involved in cell growth, survival, and division, as well as in apoptosis, stress, hypoxia, and the immune system.⁽¹⁴⁻¹⁶⁾ CIH increases the expression of NF- κ B, which then up-regulates the expression of inflammatory factors, including chemokines, adhesion molecules, cytokines, nitric oxide synthase, tissue factor and blood coagulation factor VIII.⁽¹⁷⁾ Ryan, et al⁽¹⁸⁾ found higher NF- κ B expression in intermittent hypoxia (IH) mice, in accordance with recent studies showing NF- κ B activation in cardiovascular tissue from hypoxic mice⁽¹⁷⁾ and in cells cultured under IH.⁽¹⁹⁾ IH-induced NF- κ B activation could underlie the inflammatory alterations in OSAS patients.

IH-induced activation of NF- κ B could also improve the survival rate of polymorphonuclear neutrophils (PMNs).⁽²⁰⁾ The long-term presence of a growing number of PMNs may lead to hypertension by causing endothelial dysfunction. After apoptosis, PMNs are unresponsive to inflammatory stimuli. Simultaneously, apoptosis improves the clearance of PMNs from the blood.^(21,22) Apoptosis is thus considered a fundamental mechanism for halting inflammation.

NF- κ B can provide a strong anti-apoptotic

signal in PMNs.⁽²³⁾ The up-regulation of NF- κ B under CIH, which suppresses PMN apoptosis, was confirmed by using the NF- κ B inhibitors-gliotoxin⁽²⁴⁾ and parthenolid.^(25,26) Interestingly, parthenolid has been shown to increase apoptosis only in PMNs exposed to CIH and it did not affect the apoptosis of PMNs under normoxia.⁽²⁷⁾ NF- κ B is clearly implicated in the survival of PMNs under CIH conditions.

Expression of the downstream survival gene-IL-8, induced by NF- κ B, may activate the inflammatory pathways.⁽²⁰⁾ CIH induces the activation of NF- κ B p65, after which levels of Bcl-2 and IL-8 are elevated, and PMN apoptosis is suppressed. IL-8, one of the most important survival cytokines,^(28,29) has been shown to increase PMN survival.^(30,31) IL-8 transduces its signals through interactions with chemokine receptors (CXCR) 1 and 2.⁽³²⁾ The expression of CXCR2 is down-regulated during PMN apoptosis, and up-regulation of CXCR2 during activation protects PMNs from apoptosis.^(30,33) CIH-induced expression of IL-8 thus parallels the up-regulation of CXCR2 expression and increased PMN survival.

CIH and p38 Mitogen-activated Protein Kinase

P38 mitogen-activated protein kinase (MAPK) belongs to a serine/threonine kinase family that is involved in a variety of cellular stress responses.⁽³⁴⁾ P38 MAPK was reported to increase the number of PMNs⁽³⁵⁾ and to improve PMN survival under continuous hypoxia.⁽³⁶⁾ Furthermore, a study by Dyugovskaya⁽³⁷⁾ found that the anti-PMN-apoptosis role of p38 MAPK signaling pathways is more significant under IH than under normoxia. The question arises whether p38 MAPK and NF- κ B result in the same ongoing inflammatory response. Therefore, the possibility of p38 MAPK elevating the number of PMNs through interactions with NF- κ B and TNF- α is discussed follow.

Further connecting p38 MAPK and NF- κ B, p38 MAPK plays an important role in the IH-induced activation of NF- κ B. The p38 MAPK inhibitor SB203580 increased apoptosis and decreased NF- κ B expression, and using siRNA to degrade p38 MAPK can significantly attenuate the IH-induced activation of NF- κ B.⁽³⁸⁾

Furthermore, important interactions occur between p38 MAPK and TNF- α in the development

of inflammation. TNF- α -stimulated PMN survival is entirely IL-8-dependent,⁽²⁹⁾ but the expression of IL-8 requires p38 MAPK.⁽²¹⁾ Exposure of PMNs to TNF- α results in phosphorylation of I κ B α , which leads to the activation of NF- κ B. It is widely accepted that two main TNF- α receptors are involved in PMN apoptosis: TNF receptor 1 (TNFR1) and TNFR2. Both independently mediate the activation of NF- κ B, thereby promoting PMN survival.^(39,40) Together, these results suggest that CIH, and therefore TNF- α , suppress PMN apoptosis.⁽⁴¹⁾

CIH and Oxidative Stress

Oxidative stress and inflammation have a close relationship; each appears capable of triggering the other. Under normal conditions, oxidizing agents and endogenous anti-oxidants maintain a balanced state. Once this homeostasis is disrupted, cell damage and inflammatory responses follow because of an increasingly oxidative environment.^(42,43)

Is there an increased incidence of oxidative stress in CIH? Though scholars have demonstrated that inflammation can be activated by oxidative stress, the molecular events upstream of the oxidative stress biomarkers remain unknown. The only mechanism discovered so far is that reactive oxygen species (ROS) are produced by a protein enzyme called xanthine oxidase.⁽⁴⁴⁾ Studies have shown that CIH contributes to an increase in oxidative stress markers, providing strong reasons to explore the activation of oxidative stress combined with CIH in the pathophysiology of OSAS. Discussions of clinical trials, animal experiments and treatments affecting oxidative stress follow.

Currently, small-scale clinical studies have proved that systemic oxidative stress is enhanced in OSAS patients.⁽⁴⁵⁻⁴⁷⁾ Oxidative stress biomarkers, such as advanced oxidation protein products, ferric reducing antioxidant power, superoxide radicals, serum nitrate and nitrite levels, are significantly different between OSAS patients and matched controls.⁽⁴⁸⁾ Serious OSAS patients even appear to have decreased resistance to oxidation.⁽⁴⁶⁾

Sleep-disordered breathing is related to many other factors, such as obesity, awakening stimuli and sympathetic nervous system problems, all of which promote oxidative stress.⁽⁴⁹⁾ If CIH really plays a key

role in the oxidative stress response, increases in oxidative stress markers would also be expected among animal models of hypoxia. Xu, et al⁽⁵⁰⁾ found increased levels of protein oxidation, lipid peroxidation and nucleic acid oxidation in hypoxic mouse brain cortex. Liu, et al and Nair, et al exposed mice to an IH environment, and observed a significant elevation in superoxide dehydrogenase activity and nicotinamide adenine dinucleotide phosphate oxidase.^(51,52)

Many studies have shown that continuous positive airway pressure (CPAP) treatment, for either a long or a short time, can relieve oxidative stress;⁽⁵³⁻⁵⁵⁾ the production of oxidative stress markers, including asymmetrical dimethylarginine, lipid peroxide, superoxide dismutase and thiobarbituric acid reactive substances,⁽⁵⁶⁾ is inhibited by this OSAS treatment. Del, et al⁽⁵⁷⁾ observed 138 OSAS patients and found that oxidative stress markers were reversed after CPAP treatment for 6 months. Together, these findings suggest an important inhibitory role of CPAP in oxidative stress caused by CIH.

Chinese Medicine Interventions: Clinical and Experimental Evidence

The Chinese medicines (CM) ligustrazine, rauwolfia and kudzu vine have been shown to reverse hypertension and improve related symptoms.⁽⁵⁸⁾ There is evidence of CM reversing endothelial injury. Repairing endothelial damage also has antihypertensive effects.⁽⁵⁹⁾ This has led to investigations designed to clarify whether CM interventions can protect endothelial function and reduce BP by blocking the activation of proinflammatory pathways. Is it possible for CM to make an impact on inflammatory processes? Could this propensity of inflammatory mediators to induce endothelial injury and hypertension be decreased by CM treatment? Some clinical trials found that the production and migration of inflammatory mediators, including NF- κ B, TNF- α , high-sensitive (hs)-CRP and IL-6, are favorably modified by treatment with herbal preparations.^(60,61)

NF- κ B is a central activating factor of the inflammatory cascade. Thus, research has been undertaken to investigate whether CM antihypertensive agents reduce NF- κ B levels. Zhang, et al⁽⁶²⁾ used different concentrations of astragalus polysaccharide (APS) to affect the lipopolysaccharide (LPS)-induced activation of NF- κ B in human umbilical

venous endothelial cells. LPS-induced I- κ B mRNA degradation was significantly inhibited by APS, leading to inhibition of NF- κ B in a dose-dependent manner. Sun, et al⁽⁶⁰⁾ explored the effects of puerarin on NF- κ B expression in kidneys of spontaneously hypertensive rats (SHR) by dividing 24 SHR into three groups: control group, puerarin group and benazepril group. Results showed that NF- κ B was expressed by SHR, and NF- κ B expression decreased remarkably in the benazepril group and puerarin group compared with the control group. Puerarin significantly inhibited NF- κ B.

A common inflammatory marker, high-sensitivity hs-CRP, has been considered one of the risk factors for cardiovascular disease in recent years by the European hypertension guidelines.⁽⁶³⁾ A variety of CM preparations can reduce hs-CRP, including Xuezhikang Capsule,⁽⁶¹⁾ Qingxuan Tiaoya Recipe (清眩调压方),⁽⁶⁴⁾ and calming Liver and restraining Yang formula.⁽⁶⁵⁾ These treatments showed favorable effects in depressing blood pressure and decreasing the hs-CRP levels in patients.

The level of hs-CRP has been reported to be increased significantly in cases of excess syndrome combined with deficiency syndrome compared with cases of deficiency syndrome.⁽⁶⁶⁾ The hs-CRP level was statistically higher in yin deficiency and overwhelming yang patients than in yin and yang deficiency or overwhelming liver fire patients.^(67,68) Furthermore, a small-scale randomized clinical trial showed that Jiangya Capsule (降压胶囊, a CM preparations having the efficacy of calming Liver yang) could improve the blood pressure and control the increase in serum hs-CRP in hypertensive patients.⁽⁶⁹⁾

As noted above, CM interventions reversed hypertension and inflammation, but the mechanism by which they act is unclear. However, an excess accumulation of blood stasis or a hyperactivity of yang and phlegm may indicate a higher risk of hypertension and other cardiovascular events.⁽⁷⁰⁾ Phlegm and stasis was positively correlated with IL-6 and hs-CRP levels.⁽⁷¹⁾ These findings imply that CM solved the pathological factors and the corresponding inflammatory factors were also suppressed. Wang, et al⁽⁷²⁾ divided 60 hypertension patients into excessive accumulation of phlegm dampness type cases and non-excessive accumulation of phlegm dampness type cases. The former patients received Tanreqing

Injection (痰热清注射液) plus telmisartan and the latter cases received telmisartan only. Six weeks later, compared with control patients, the levels of hs-CRP, TNF- α and IL-6 were significantly increased in hypertensive patients. After treatment, levels of the three inflammatory factors aforementioned were significantly reduced in the excessive accumulation of phlegm dampness group.

Overall, the data indicate that CM reduced BP and inhibited NF- κ B, TNF- α and IL-6. These findings imply that CM interventions are more effective than Western medicine in improving the inflammatory response in patients with hypertension. CM treatment against hypertension was also shown to improve the apnea hypopnea index (AHI) of OSAS patients.⁽⁷³⁾ Together, these results strengthen the need for early CM intervention in apneic individuals at risk for hypertension, and suggest new therapeutic options for these patients.

Conclusions and Perspectives

Despite the high correlation between OSAS and hypertension, little evidence is available regarding their pathophysiological and clinical outcomes. Most information involves cross-sectional analyses of selected clinical cohorts and there are no long-term follow-up studies. Future research into the relationship between CIH and hypertension needs to include matched control subjects, more consistent definitions and stricter criteria for outcomes. The finding that CIH-induced hypertension mainly involves systemic inflammation needs to be confirmed.

Although there is evidence in CIH and hypertension of overlapping mechanisms involving inflammation, oxidative stress and leukocyte dysfunction, there may be differences in the magnitude and consequences of these responses. Furthermore, the role of inflammatory markers in predicting hypertension in OSAS is unclear and lacks long-term prospective studies. If systemic inflammation is important in CIH and hypertension, studies in overlapping patients should provide insights regarding the nature and significance of these responses. These findings are clinically relevant because systemic inflammation may contribute to the pathogenesis of hypertension, and the molecular pathways involved are similar in OSAS. Studies of patients with OSAS and hypertension should provide insights into the mechanisms of systemic inflammation and help to

develop new therapies. Great strides have been made in establishing obstructive sleep apnea as a target for therapy in hypertensive patients. Additionally, CM interventions have shown some advantages in controlling OSAS-related hypertension. In the future, based on the relationship between CIH, inflammation and hypertension, further clinical and animal experimental studies of integrative medicine will be needed to offer high quality evidence for the prevention and control of OSAS-related hypertension.

REFERENCES

- Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 1993;328:1230-1235.
- Parish JM, Somers VK. Obstructive sleep apnea and cardiovascular disease. *Mayo Clin Proc* 2004;79:1036-1046.
- Mancia G, De Backer G, Dominiczak A, Cifkova R, Fagard R, Germano G. 2007 Guidelines for the management of arterial hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J* 2007;28:1462-1536.
- Lugaresi E, Coccagna G, Mantovani M, Brignani F. Effects of tracheostomy in two cases of hypersomnia with periodic breathing. *J Neurol Neurosurg Psychiatry* 1973;36:15-26.
- Coccagna G, Mantovani M, Brignani F, Parchi C, Lugaresi E. Tracheostomy in hypersomnia with periodic breathing. *Bull Physiopathol Respir (Nancy)* 1972;8:1217-1227.
- Motta J, Guilleminault C, Schroeder JS, Dement WC. Tracheostomy and hemodynamic changes in sleep-inducing apnea. *Ann Intern Med* 1978;89:454-458.
- Das AM, Khayat R. Hypertension in obstructive sleep apnea: risk and therapy. *Expert Rev Cardiovasc Ther* 2009;7:619-626.
- Adedayo AM, Olafiranye O, Smith D, Hill A, Zizi F, Brown C, et al. Obstructive sleep apnea and dyslipidemia: evidence and underlying mechanism. *Sleep Breath* 2012;Aug 18. [Epub ahead of print]
- Karamanli H, Ozol D, Ugur KS, Yildirim Z, Armutcu F, Bozkurt B, et al. Influence of CPAP treatment on airway and systemic inflammation in OSAS patients. *Sleep Breath* 2012; Sep 4. [Epub ahead of print]
- Svensson M, Venge P, Janson C, Lindberg E. Relationship between sleep-disordered breathing and markers of systemic inflammation in women from the general population. *J Sleep Res* 2012;21:147-154.
- Kaviraj B, Bai SC, Su L, Zheng XO, Huang R, Li TP, et al. Effect of obstructive sleep apnea syndrome on serum C-reactive protein level, left atrial size and premature atrial contraction. *J Southern Med Univ (Chin)* 2011;31:197-200.
- Chen J, Zhang JC, Chen YY. Relationship between hs-CRP and obstructive sleep apnea in hypertension patients. *Chin J Integr Med Cardio-/Cerebrovascular Dis (Chin)* 2012;3:280-282.
- McNicholas WT. Chronic obstructive pulmonary disease and obstructive sleep apnea: overlaps in pathophysiology, systemic inflammation, and cardiovascular disease. *Am J Respir Crit Care Med* 2009;180:692-700.
- Rauch DA, L Ratner. Targeting HTLV-1 activation of NF-kappa B in mouse models and ATLL patients. *Viruses* 2011;3:886-900.
- Barnes PJ. Pathophysiology of allergic inflammation. *Immunol Rev* 2011; 242:31-50.
- Dejean E, Foisseau M, Lagarrigue F, Lamant L, Prade N, Marfak A, et al. ALK+ALCLs induce cutaneous, HMGB-1-dependent IL-8/CXCL8 production by keratinocytes through NF-kappa B activation. *Blood* 2012;119:4698-4707.
- Greenberg H, Ye X, Wilson D, Htoo AK, Hendersen T, Liu SF. Chronic intermittent hypoxia activates nuclear factor-kappa B in cardiovascular tissues *in vivo*. *Biochem Biophys Res Commun* 2006; 343:591-596.
- Ryan S, Taylor CT, McNicholas WT. Selective activation of inflammatory pathways by intermittent hypoxia in obstructive sleep apnea syndrome. *Circulation* 2005;112:2660-2667.
- Li S, Qian XH, Zhou W, Zhang Y, Feng J, Wan NS, et al. Time-dependent inflammatory factor production and NF-kappa B activation in a rodent model of intermittent hypoxia. *Swiss Med Wkly* 2011;14:w13309.
- Walmsley SR, Print C, Farahi N, Peyssonnaud C, Johnson RS, Cramer T, et al. Hypoxia-induced neutrophil survival is mediated by HIF-1alpha-dependent NF-kappaB activity. *J Exp Med* 2005;201:105-115.
- Duncan AL, Leuenroth SJ, Grutkoski P, Ayala A, Simms HH. TNF alpha-induced suppression of PMN apoptosis is mediated through interleukin-8 production. *Shock* 2000;14:284-289.
- Maiani AN, Kuijpers TW, Roos D. Apoptosis of neutrophils. *Acta Haematol* 2004;111:56-66.
- Ward C, Walker A, Dransfield I, Haslett C, Rossi AG. Regulation of granulocyte apoptosis by NF-kappa B. *Biochem Soc Trans* 2004;32:465-467.
- Pahl HL, Krauss B, Schulze-Osthoff K, Decker T, Traenckner EB, Vogt M. The immunosuppressive fungal metabolite gliotoxin specifically inhibits transcription factor NF-kappaB. *J Exp Med* 1996;183:1829-1840.
- Magni P, Ruscica M, Dozio E, Rizzi E, Beretta G, Facino RM. Parthenolide inhibits the LPS-induced secretion of IL-6 and TNF-alpha and NF-kappa B nuclear translocation in BV-2 microglia. *Phytother Res* 2012;1405-1409.

26. Dyugovskaya L, Polyakov A, Ginsberg D, Lavie P, Lavie L. Molecular pathways of spontaneous and TNF (alpha)-mediated neutrophil apoptosis under intermittent hypoxia. *Am J Respir Cell Mol Biol* 2011;45:154-162.
27. Hehner SP, Hofmann TG, Droge W, Schmitz ML. The antiinflammatory sesquiterpene lactone parthenolide inhibits NF-kappa B by targeting the I kappa B kinase complex. *J Immunol* 1999;163:5617-5623.
28. Luo HR, Loison F. Constitutive neutrophil apoptosis: mechanisms and regulation. *Am J Hematol*, 2008;83:288-295.
29. Cowburn AS, Deighton J, Walmsley SR, Chilvers ER. The survival effect of TNF-alpha in human neutrophils is mediated via NF-kappa B-dependent IL-8 release. *Eur J Immunol* 2004; 34:1733-1743.
30. Glynn PC, Henney E, Hall IP. The selective CXCR2 antagonist SB272844 blocks interleukin-8 and growth-related oncogene-alpha-mediated inhibition of spontaneous neutrophil apoptosis. *Pulm Pharmacol Ther* 2002;15:103-110.
31. Acorci MJ, Dias-Melicio LA, Golim MA, Bordon-Graciani AP, Peracoli MT, Soares AM. Inhibition of human neutrophil apoptosis by *Paracoccidioides brasiliensis*: role of interleukin-8. *Scand J Immunol* 2009;69:73-79.
32. Javor J, Bucova M, Cervenova O, Kralinsky K, Sadova E, Suchankova M, et al. Genetic variations of interleukin-8, CXCR1 and CXCR2 genes and risk of acute pyelonephritis in children. *Int J Immunogenet* 2012;39:338-345.
33. Allegretti M, Bertini R, Cesta MC, Bizzarri C, Di Bitondo R, Di Cioccio. V 2-Arylpropionic CXC chemokine receptor 1 (CXCR1) ligands as novel noncompetitive CXCL8 inhibitors. *J Med Chem* 2005;48:4312-4331.
34. Alvarado-Kristensson M, Melander F, Leandersson K, Ronnstrand L, Wernstedt C, Andersson T. p38-MAPK signals survival by phosphorylation of caspase-8 and caspase-3 in human neutrophils. *J Exp Med* 2004;199:449-458.
35. Villunger A, O'Reilly LA, Holler N, Adams J, Strasser A. Fas ligand, Bcl-2, granulocyte colony-stimulating factor, and p38 mitogen-activated protein kinase: regulators of distinct cell death and survival pathways in granulocytes. *J Exp Med* 2000;192:647-658.
36. Leuenroth SJ, Grutkoski PS, Ayala A, Simms HH. Suppression of PMN apoptosis by hypoxia is dependent on Mcl-1 and MAPK activity. *Surgery* 2000;128:171-177.
37. Dyugovskaya L, Polyakov A, Lavie P, Lavie L. Delayed neutrophil apoptosis in patients with sleep apnea. *Am J Respir Crit Care Med* 2008;177:544-554.
38. Ryan S, McNicholas WT, Taylor CT. A critical role for p38 map kinase in NF-kappa B signaling during intermittent hypoxia/reoxygenation. *Biochem Biophys Res Commun* 2007;355:728-733.
39. Murray J, Barbara JA, Dunkley SA, Lopez AF, Van Ostade X, Condliffe AM, et al. Regulation of neutrophil apoptosis by tumor necrosis factor-alpha: requirement for TNFR55 and TNFR75 for induction of apoptosis *in vitro*. *Blood* 1997;90:2772-2783.
40. Cabrini M, K Nahmod, J Geffner. New insights into the mechanisms controlling neutrophil survival. *Curr Opin Hematol* 2010;17:31-35.
41. Kilpatrick LE, Sun S, Mackie D, Baik F, Li H, Korchak, HM. Regulation of TNF mediated antiapoptotic signaling in human neutrophils: role of delta-PKC and ERK1/2. *J Leukoc Biol* 2006;80:1512-21.
42. Chen SJ, Yen CH, Huang YC, Lee BJ, Hsia S, Lin PT. Relationships between inflammation, adiponectin, and oxidative stress in metabolic syndrome. *PLoS One* 2012;7:e45693.
43. Sesti F, Tsitsilonis OE, Kotsinas A, Trougakos IP. Oxidative stress-mediated biomolecular damage and inflammation in tumorigenesis. *In Vivo* 2012;26:395-402.
44. Murata M, Fukushima K, Takao T, Seki H, Takeda S, Wake N. Oxidative stress produced by xanthine oxidase induces apoptosis in human extravillous trophoblast cells. *J Reprod Dev* 2012; Sep 14. [Epub ahead of print].
45. Baysal E, Taysi S, Aksoy N, Uyar M, Celenk E, Karatas ZA, et al. Serum paraoxonase, arylesterase activity and oxidative status in patients with obstructive sleep apnea syndrome (OSAS). *Eur Rev Med Pharmacol Sci* 2012;16:770-774.
46. Simiakakis M, Kapsimalis F, Chaligiannis E, Loukides S, Sitaras N, Alchanatis M. Lack of effect of sleep apnea on oxidative stress in obstructive sleep apnea syndrome (OSAS) patients. *PLoS One* 2012;7:e39172.
47. Ntalapascha M, Makris D, Kyparos A, Tsilioni I, Kostikas K, Gourgoulialis K, et al. Oxidative stress in patients with obstructive sleep apnea syndrome. *Sleep Breath* 2012; May 18. [Epub ahead of print].
48. Mancuso M, Bonanni E, LoGerfo A, Orsucci D, Maestri M, Chico L, et al. Oxidative stress biomarkers in patients with untreated obstructive sleep apnea syndrome. *Sleep Med* 2012;13:632-636.
49. Williams A, Scharf SM. Obstructive sleep apnea, cardiovascular disease, and inflammation—is NF-kappa B the key? *Sleep Breath* 2007;11:69-76.
50. Xu W, Chi L, Row BW, Xu R, Ke Y, Xu B, et al. Increased oxidative stress is associated with chronic intermittent hypoxia-mediated brain cortical neuronal cell apoptosis in a mouse model of sleep apnea. *Neuroscience* 2004;126:313-323.
51. Liu JN, Zhang JX, Lu G, Qiu Y, Yang D, Yin GY, et al. The effect of oxidative stress in myocardial cell injury in mice exposed to chronic intermittent hypoxia. *Chin Med J*

- 2010;123:74-78.
52. Nair D, Dayyat EA, Zhang SX, Wang Y, Gozal D. Intermittent hypoxia-induced cognitive deficits are mediated by NADPH oxidase activity in a murine model of sleep apnea. *PLoS One* 2011;6:e19847.
 53. Barcelo A, Miralles C, Barbe F, Vila M, Pons S, Agusti AG. Antioxidant status in patients with sleep apnoea and impact of continuous positive airway pressure treatment. *Eur Respir J* 2006;27:756-760.
 54. Minoguchi K, Yokoe T, Tanaka A, Ohta S, Hirano T, Yoshino G, et al. Association between lipid peroxidation and inflammation in obstructive sleep apnoea. *Eur Respir J* 2006;28:378-385.
 55. Alzoughaibi MA, Bahammam AS. The effect of one night of continuous positive airway pressure therapy on oxidative stress and antioxidant defense in hypertensive patients with severe obstructive sleep apnea. *Sleep Breath* 2012;16:499-504.
 56. Oyama J, Yamamoto H, Maeda T, Ito A, Node K, Makino N. Continuous positive airway pressure therapy improves vascular dysfunction and decreases oxidative stress in patients with the metabolic syndrome and obstructive sleep apnea syndrome. *Clin Cardiol* 2012;35:231-236.
 57. Del Ben M, Fabiani M, Loffredo L, Polimeni L, Carnevale R, Baratta F, et al. Oxidative stress mediated arterial dysfunction in patients with obstructive sleep apnoea and the effect of continuous positive airway pressure treatment. *BMC Pulm Med* 2012;12:36.
 58. Liu LS, Chen MQ, Zeng GY, Zhou BF. A forty-year study on hypertension. *Acta Acad Med Sini (Chin)* 2002;24:401-408.
 59. Chen YY, Zhang JC. Research status of endothelial mechanism in high blood pressure of traditional Chinese and Western medicine. *Chin J Gerontol (Chin)* 2010;30:3010-3013.
 60. Sun WC, Zhou HP. The effects of puerarin on renal arterial NF- κ B expression in spontaneous hypertensive rats. *J Emerg Tradit Chin Med (Chin)* 2008;17:965-966.
 61. Xiong IH, Wang M, Zhang RL, Huang WJ. Observation on the intervention efficacy of Xuezhikang Capsule on levels of CRP and IL-6 in hypertension patients with different complications. *Liaoning J of Tradit Chin Med (Chin)* 2012;39:1337-1340.
 62. Zhang JZ, Chen LG, Hu XQ, Jia HX. Influence of astragalus polysaccharide on the expression of Toll-like receptor 4 and nuclear transcription factor- κ B in essential hypertension patients with blood stasis syndrome. *J Traditi Chin Med (Chin)* 2011;52:1286-1304.
 63. Whitworth JA. 2003 World Health Organization (WHO)/ International Society of Hypertension (ISH) statement on management of hypertension. *J Hypertens* 2003;21:1983-1992.
 64. Tao LL, Ma XC, Chen KJ. Clinical study on effect of Qingxuan Tiaoya Recipe in treating menopausal women with hypertension. *Chin J Integr Tradit West Med (Chin)* 2009;8:680-684.
 65. Zhong GW, Luo YH, Xiang LL, Xie Y, Xie QQ, Li YH, et al. Clinical efficacy study on calming Liver and restraining Yang formula in treating patients with mild or moderate degree of essential hypertension. *Chin J Chin Mate Med (Chin)* 2010;35:776-781.
 66. Suo HL, Wang SR, Wu AM, Zhang DM, Shi FQ. Risk stratification of EH and correlation between TCM syndromes of EH and high-sensitivity C-reactive protein in clinic. *J Beijing Univ Tradit Chin Med (Chin)* 2009;32:265-269.
 67. Qing Y, Zhang ZZ, Yang LY. Connection between pattern of traditional Chinese medicine syndrome in primary hypertension and C-reactive protein. *J Guangxi Tradit Chin Med Univ (Chin)* 2008;11:18-19.
 68. Zhang ZB, Zhou CG, Lu S. Distribution of TCM syndrome types of essential hypertension and their relationship to biochemical indicators. *Liaoning J Tradit Chin Med (Chin)* 2010;37:969-971.
 69. Li H, Zhao WM, Han YX. Effect of a integrative medical regimen on levels of vascular endothelial function and hypersensitive C-reactive protein in elderly patients with isolated systolic hypertension. *Chin J Integr Tradit West Med (Chin)* 2009;29:115-119.
 70. Fu DY, Zu LH, Xin XY, He SY, Zhang XM, Wang SH, et al. Analysis of cardiovascular vulnerable factors among 409 cases of EH patients with different patterns of TCM. *Univ Tradit Med Sin Pharmacol Shanghai (Chin)* 2009;23:15-18.
 71. Liu LL, Guo RJ, Zhang YL. Relationship between syndrome factor of hypertension and interleukin-6, highly sensitive C-reactive protein and matrix metalloproteinase-9. *Chin J Tradit Chin Med Pharm (Chin)* 2011; 26:1583-1586.
 72. Wang X, Li WQ. The expression and the intervention of Tanreqing Injection of the pulse pressure and the levels of serum inflammatory factors in hypertensive patients of excessive accumulation of phlegm dampness type. *J Emerg Tradit Chin Med (Chin)* 2009;18:1389-1414.
 73. Zhang JC, Chen YY, Chen J, Liu JG. The impact of Qingxuan Granule on endothelial function in hypertensive patients with and without obstructive sleep apnea. *Chin J Gerontol (Chin)* 2012;23:672-675.

(Received December 27, 2012)
 Edited by CHEN Yi-yu