

HYPERTENSION COMPENDIUM

The Gut Microbiome in Hypertension

Recent Advances and Future Perspectives

Ellen G. Avery¹, Hendrik Bartolomaeus², Andras Maifeld³, Lajos Marko⁴, Helge Wiig⁵, Nicola Wilck⁶,
Stephan P. Rosshart⁷, Sofia K. Forslund⁸, Dominik N. Müller⁹

ABSTRACT: The pathogenesis of hypertension is known to involve a diverse range of contributing factors including genetic, environmental, hormonal, hemodynamic and inflammatory forces, to name a few. There is mounting evidence to suggest that the gut microbiome plays an important role in the development and pathogenesis of hypertension. The gastrointestinal tract, which houses the largest compartment of immune cells in the body, represents the intersection of the environment and the host. Accordingly, lifestyle factors shape and are modulated by the microbiome, modifying the risk for hypertensive disease. One well-studied example is the consumption of dietary fibers, which leads to the production of short-chain fatty acids and can contribute to the expansion of anti-inflammatory immune cells, consequently protecting against the progression of hypertension. Dietary interventions such as fasting have also been shown to impact hypertension via the microbiome. Studying the microbiome in hypertensive disease presents a variety of unique challenges to the use of traditional model systems. Integrating microbiome considerations into preclinical research is crucial, and novel strategies to account for reciprocal host-microbiome interactions, such as the wildling mouse model, may provide new opportunities for translation. The intricacies of the role of the microbiome in hypertensive disease is a matter of ongoing research, and there are several technical considerations which should be accounted for moving forward. In this review we provide insights into the host-microbiome interaction and summarize the evidence of its importance in the regulation of blood pressure. Additionally, we provide recommendations for ongoing and future research, such that important insights from the microbiome field at large can be readily integrated in the context of hypertension.

Key Words: blood pressure ■ fasting ■ fatty acids, volatile ■ hypertension ■ immune system ■ microbiota ■ translational medical research

Microbes are everywhere. They self-organize, creating complex ecosystems within otherwise uninhabitable niches, rapidly adapting to their environment. The human holobiont is the assembly of many different species to form a singular functional unit. Adult human beings contain slightly >50% microbial cells, outnumbering the cells of the human host (3.0×10^{13}). During pregnancy, the developing fetus is virtually sterile, though upon birth, newborns immediately collect up to 3.8×10^{13} bacteria from 500 to 1000 different microbial species¹ at important epithelial barrier sites, reaching a total mass of about 1.5 kg during adulthood, about the same weight as the liver. The microbiome, defined as a catalog of all microbes in a site and their genes, encodes over 40 million distinct gene variants.² Half of these gene variants are unique to a single individual, which demonstrates

the striking heterogeneity of human microbiome data and perhaps may explain the elusiveness of a universally healthy microbiome configuration.^{2,3} The host depends on the microbiome for several fundamental symbiotic functions, such as the priming of the immune system and the production of essential vitamins, as well as for the energy harvest from foods. The gut microbiota, defined as the microbial taxa within a human, is now regarded an endocrine organ that generates metabolites that can act as effectors in the host, triggering responses in the local microenvironment or distant target organ systems such as the heart, kidney, vasculature, and brain (Figure 1).⁴

The first description of a living organism in the human gastrointestinal tract dates back to 1681 when Antonie van Leeuwenhoek reported a number of little animals in his stool samples; followed by the isolation of *Escherichia*

Correspondence to: Dominik N. Müller, PhD, Experimental and Clinical Research Center and Max Delbrück Center for Molecular Medicine in the Helmholtz Association, Lindenberger Weg 80, 13125 Berlin, Germany. Email dominik.mueller@mdc-berlin.de

For Sources of Funding and Disclosures, see page 945.

© 2021 American Heart Association, Inc.

Circulation Research is available at www.ahajournals.org/journal/res

Nonstandard Abbreviations and Acronyms

AT1	angiotensin II type 1
BP	blood pressure
CVD	cardiovascular disease
DASH	Dietary Approaches to Stop Hypertension
F/B	Firmicutes-to-Bacteroidetes
FFAR	free fatty acid receptor
FMT	fecal microbiota transplantation
GF	germ free
Gpr41	G-protein–coupled receptor 41
Gpr43	G-protein–coupled receptor 43
Gpr109a	G-protein–coupled receptor 109 A
HDAC	histone deacetylase
IFN	interferon
IL	interleukin
LI	large intestine
mTOR	mechanistic target of rapamycin
Olfr78	olfactory receptor 78
Olfr558	olfactory receptor 558
SCFA	short-chain fatty acid
SHR	spontaneously hypertensive rat
SI	small intestine
STAT3	signal transducer and activator of transcription 3
Treg	regulatory T cell
WKY	Wistar-Kyoto

coli as the first species from the gastrointestinal tract in 1885. Dramatic technological developments ensued over the proceeding centuries with the establishment of high-throughput sequencing technologies and metabolomics, as well as developments in high-performance computing and artificial intelligence. These advancements have opened a new avenue to decipher the interrelationship between lifestyle, diet, pharmacotherapy, and the gut microbiome. Each individual's respective healthy gut microbiome is relatively stable over time and coexists in equilibrium with the surrounding environment.⁵ Perturbations such as antibiotics, intestinal infections, and profound dietary or lifestyle changes such as moving on or off a Western diet can induce transient or persistent changes to this ecosystem.^{5,6} Microbiomes from diseased and nondiseased individuals differ (exhibiting a dysbiotic as opposed to eubiotic state), and this has been demonstrated to hold for several inflammatory (eg, colitis), cardiovascular, and metabolic disorders.^{7–11} In recent years, the microbiome field has increasingly shifted focus from taxonomic (Who's there?) to functional (What are they doing?) analyses with an accompanying emphasis on host-microbiome interactions at the molecular

mechanistic level across time and space.¹² Furthermore, the interplay between the host and its microbiomes is a critically important consideration. As this reciprocal interaction becomes more evident, novel strategies to account for it in preclinical disease modeling are emerging. A major future challenge is to move beyond biomarker identification to identify bioactive metabolites, which could be candidates for therapeutic use. Nevertheless, improved understanding of the host-microbiome interface provides an exciting new avenue for the development of pharmacological and nonpharmacological treatments for hypertensive disease.

GUT MICROBIOME-HOST IMMUNE INTERACTION

The internal environment of the intestines acts as an interface between the external environment and the host and is constantly challenged by the consumption habits of the host. On the luminal side, microbes are able to attach and colonize this space, while on the host side, the gastrointestinal tract acts as the largest compartment for immune cells in the body. The physiology of the gastrointestinal tract and accompanying immune cells has been extensively reviewed (by Mowat and Agace¹³). Additionally, the structural and functional comparability of human and mouse gastrointestinal physiology, and the resultant implications for translational microbiome research, has been outlined elegantly elsewhere (Nguyen et al¹⁴). Therefore, here we will briefly describe the aspects necessary to understand gastrointestinal involvement in hypertensive disease, though our review of this topic is by no means comprehensive.

Anatomically, the intestines consist of different segments. The duodenum, jejunum, and ileum represent the small intestine (SI), which occupies much more physical space than the large intestine (LI), made up of the colon and rectum. Rodents have an enlarged cecum compared to humans, which is a blind-ended sac connecting the SI and LI.¹⁴ In mice, the cecum acts as a large reservoir for the commensal microbes that are involved in the fermentation of fibers that cannot be otherwise cleaved by SI enzymes.¹³ The role of the cecum in mice is nontrivial, as this acts as the major sight of production for short-chain fatty acids (SCFAs), and removal of this compartment led to increased inflammation at distal sites within the gastrointestinal tract.¹⁵ In humans, the volume of microbes is far smaller than in mice, but this compartment still plays an important role in facultative anaerobic fermentation.¹⁶

Of note, the composition and abundance of commensal microbes is distinct in the different gastrointestinal regions, for example, abundance of microbes is low in the adult SI ($<10^5$ microorganisms/mL) and increases to 10^{12} in the colon.¹³ The SI and LI have distinct physiological functions. While the duodenum and jejunum are




Figure 1. The relationship between blood pressure and the gut microbiome.

Ingested food is transformed by the gut microbiome into small metabolites. Food antigens, microbially produced metabolites, and the microbes themselves all contribute to immune homeostasis. Perturbations to the symbiotic relationship between host and the microbiome can lead directly or indirectly, via the immune system, to changes in blood pressure and associated heart, vascular, or kidney damage. Illustration credit: Ben Smith.

involved in the process of digestion, absorption of nutrients, and motility, the LI has 3 primary functions: absorption of water and electrolytes, production and absorption of vitamins, and formation and transport of feces for elimination.¹⁷ Mucus lining the gut lumen represents a physiological barrier against bacterial infections and can bind toxins.¹⁸ In addition, the mucus serves as a nutrient source for bacteria and thus affects colonization by microbes that have the ability to survive and expand in the mucus layer.¹⁹ *Akkermansia muciniphila* and *Citrobacter rodentium* are capable of degrading mucin, and the latter blooms during fiber deprivation.²⁰ Loss of integrity of the colonic mucus layer increases host susceptibility to pathogens. Under healthy conditions, the tight epithelial layer prevents the invasion of pathogenic microbes while certain stimuli like inflammatory disease or a Western diet can lead intestinal permeability and the development of so-called leaky gut syndrome.²¹ Altogether, the evidence pinpoints to the pivotal role of the gut microbiota in gut epithelial health.²²

The intestine is continually exposed food particles and food antigens, physiological or opportunistic microbiota-derived metabolites, and other immunomodulatory stimuli. Immune cells within the gastrointestinal tract not only respond to antigenic provocations within the gut but have been shown to egress to distal organs throughout the body, indicating their importance in system-wide inflammatory homeostasis.^{23,24} We are now just beginning to understand the full breadth of spatial interactions of these stimuli with the respective immune compartment (Figure 2). Gnotobiotic mice are those with a defined community. Germ-free (GF) mice, which are devoid of all bacteria, are an extreme but useful gnotobiotic model system used to elucidate the impact of bacteria on the host immune system and physiology. GF mice have increased gut permeability, which can be reversed upon colonization with microbes.²⁵ GF mice are also severely compromised in immune cell function and lymphoid organ development.²⁶ Colonization experiments have demonstrated the importance of certain




Figure 2. Intestinal spatial variability can be found on both the host and microbiome sides. Relative levels of luminal and tissue-associated content are illustrated here, suggestive of the regional specialization of both features. The luminal contents of the intestines are known to vary significantly, in terms of microbial load, the microbial inhabitants, and the resultant microbially produced metabolites.^{13,187} Shown here is a proposed scheme, although the inhabitants and regional specifications throughout the gastrointestinal tract are subject to individual differences in both mice and men. In accordance with the variations in luminal content, the host immune system is likewise regionally specific.^{13,178} Shown here are immune cells where spatial dynamics have been demonstrated during immune homeostasis. SCFA indicates short-chain fatty acid; Th1, type 1 helper; Th17, T helper 17; CD8+ T, cytotoxic T; and Treg, regulatory T cell. Modified from Mowat and Agace.¹³

microbes such as segmented filamentous bacterium for the regional specialization of immune cells along the length of the intestine.²⁷ They have also provided detailed insight as to how human-resident bacterial species may modulate the host immune system.²⁸ The importance of the microbiota-immune dynamics in human disease is clear as well. For example, fecal microbiota transplantation (FMT) affects the severely perturbed immune system in patients after radiation and chemotherapy during hematopoietic cell transplantation.²⁹ Furthermore, microbial colonization in early life is critical for the maturation of the human immune system,³⁰ and perturbations during this phase have been shown to influence susceptibility to allergies³¹ and infectious diseases.³²

Over the last few decades, experimental and clinical studies have demonstrated that cells of the innate

and adaptive immune system play pivotal roles in the pathogenesis of hypertension, target organ damage, and cardiovascular disease (CVD).^{33–35} Proinflammatory effector memory T cells and T helper cell subtypes T helper 17 cells (Th17; producing IL [interleukin]-17) and type 1 helper cells (producing IFN [interferon]- γ) promote hypertension and cardiovascular target organ damage, whereas regulatory T cells (Tregs) typically produce high amounts of anti-inflammatory IL-10 and can ameliorate vascular, cardiac, and renal damage.^{36–41} In addition, gamma delta T cells⁴² and myeloid-derived suppressor cells⁴³ also play an important role in the pathogenesis of hypertension. Dendritic cells, which can alter the activation state of several T-cell subtypes, have been shown to increase in salt-responsive hypertension and are suggested to play a role in the interplay between

microbial dysbiosis and blood pressure (BP).⁴⁴ Bacteria can communicate with different immune cells involved in CVD, either directly or through the metabolites that they produce. For example, segmented filamentous bacteria or *Bifidobacterium adolescentis* induce Th17 cells,^{27,45–47} whereas *Lactobacillus murinus* and their tryptophan metabolite indole-3 lactic acid inhibits Th17 cells.⁴⁸ Further, *Clostridium* spp. and the SCFA butyrate are outstanding inducers of Tregs in the colon.^{49,50}

GUT MICROBIOTA AND HYPERTENSION

The development of high BP is a complex, multifactorial process that involves both genetic and environmental risk factors. While 901 loci have been identified in the latest genome-wide association study, altogether this can only explain 5.7% of BP variability.⁵¹ Additionally, population data from the UK Biobank suggest that lifestyle factors can account for up to 4 to 5 mmHg.⁵² Of note, even a 2-mmHg BP reduction would decrease overall CVD mortality by 7%.⁵³ Much of the evidence for the role of the gut microbiome and hypertension has accumulated in the last decade due to the reduction in costs for next-generation sequencing. Several cross-sectional studies in humans show an association between gut microbiome and BP or hypertension.^{54–63} Alpha diversity describes the microbial variance within a given ecosystem, captured as a biological sample. The majority of published microbiome studies in humans identified a reduced alpha diversity in hypertensive patients or in patients with higher BP,^{55–60,63} which has also been observed in obesity, hyperinsulinemia, and dyslipidemia. Numerous human gut microbiome studies have reported an association between a higher abundance of Gram-negative microbiota including *Klebsiella*, *Parabacteroides*, *Desulfovibrio*, and *Prevotella* and higher BP, although not all studies were able to identify this pattern.^{54,56,59,61} Data from the cross-sectional HELIUS cohort study⁵⁸ (HEalthy Life In an Urban Setting study) demonstrated that *Klebsiella* spp. and *Streptococcaceae* spp. were positively correlated with BP, as found previously.^{59,61} Moreover, GF mice that received FMT from a hypertensive human donor developed a similar gut microbiota to their donor, as well as elevated systolic and diastolic BP after 8 weeks when compared with GF mice, which received an FMT from 2 normotensive donors.⁵⁶ In addition, there are several valuable rodent hypertension models that have analyzed the role of the gut microbiome and BP.^{48,60,64–67} Adnan et al⁶⁴ demonstrated that the gut microbiota of stroke-prone SHR (spontaneously hypertensive rats) is dysbiotic and significantly different than the microbiota of normotensive WKY (Wistar-Kyoto) control rats. Furthermore, FMT from stroke-prone SHR into WKY controls increased the systolic BP of these otherwise normotensive rats.⁶⁴ Dysbiosis was also described for hypertension models such as angiotensin II-infused

mice,⁶⁶ Dahl salt-sensitive rats,⁶⁵ as well as high salt-treated mice,⁴⁸ and deoxycorticosterone acetate-salt hypertensive mice.⁶⁷ Additionally, a recent study demonstrated in SHR rats that microbial dysbiosis was linked to pathophysiological changes in the gastrointestinal tract and diminished intestinal integrity.⁶⁸ Furthermore, intestinal permeability and dysbiosis in SHR could be remedied by treating rats with the antihypertensive drug losartan.⁶⁹

The link between the gut microbiome and hypertension is not species specific. For example, high salt treatment in both mice and humans reduced *Lactobacillus* spp. abundance and led to an increase in BP.⁴⁸ Of note, a modest reduction in salt in therapy-naïve hypertensive patients was able to reduce BP and improve arterial compliance.⁷⁰ Improved clinical outcomes were accompanied by an increase in 8 circulating SCFAs (including 2-methylbutyrate, butyrate, hexanoate, isobutyrate, and valerate).⁷⁰ Furthermore, it was shown that probiotic *Lactobacillus* treatment inhibited Th17 cells and ameliorated salt-sensitive hypertension by restoring indole-3 lactic acid levels, which is product of microbial tryptophan metabolism.⁴⁸ In addition, *Lactobacillus coryniformis* has been shown to improve vascular function and insulin sensitivity.⁷¹ *Lactobacillus* treatment not only ameliorates CVDs but also improves experimental autoimmune disease outcomes (detailed in depth in a previous review⁷²). A systematic review of randomized controlled trials investigating the role of probiotics on high BP showed that *Lactobacillus*-containing probiotics are effective, if used in sufficiently high doses and for at least 8 weeks.⁷³ Despite preliminary advances to understand the role of the microbiome in the regulation of BP, it is also essential to elucidate how environmental factors affect this nexus.

LIFESTYLE AFFECTS THE GUT MICROBIOME

In humans, core microbial communities in the gut are stable and change only in response to major perturbations such as enteric infection, global travel, or medication, leading to transient or persistent changes in the gut microbiome.^{5,74} In addition, members of the gut microbiota are not only reactive to proportions of certain dietary stimuli but may also respond in a spatiotemporal context. Currently, our understanding of the exact mechanisms by which specific dietary changes affect susceptibility to inflammatory, autoimmune, and cardiovascular diseases is rather vague. Using machine learning algorithms trained on microbiome composition and function offers an exciting opportunity to facilitate better predictions of responsiveness to nutritional stimuli. A landmark study by Segal et al⁷⁵ investigated a large cohort of healthy individuals and found that they could predict glycemic variability in response to a meal challenge using individual microbiome and nutritional information.

Emerging research suggests that dietary factors (high salt or high fiber) and lifestyle interventions (salt restriction or caloric restriction) influence microbial community structure and function, and this has major implications for immune cell activation and BP. A Western lifestyle typically involves consuming several main meals per day and leads to decreased bacterial diversity, the overgrowth of some commensals, and concomitant suppression of others.^{76,77} Accordingly, production of metabolites by the bacterial community is shifted,⁷⁸ which promotes inflammation⁷⁹ and ultimately can lead to the development of diseases such as obesity⁸⁰ and atherosclerosis.⁸¹ While historically, meals were often cooked freshly, nowadays humans more frequently consume processed foods, which generally have a higher salt content.⁸² This lifestyle often leads to a higher salt intake than the recommendations of medical guidelines or experts.^{83–85} To reduce the risk of cardiometabolic disease, a healthy diet and exercise are often prescribed. Most recommendations center around changing the Western diet, which is rich in saturated fat, sugar, salt, and calories but low in fiber, to a healthier Mediterranean-like Dietary Approaches to Stop Hypertension (DASH) diet⁸⁶ to achieve optimal nutrition, including negative energy balance and lower salt intake, although compliance is a major challenge. Fasting, as an extreme form of caloric restriction, plays an important role in different cultural and religious practices. Dramatic caloric restriction not only affects host health and physiology but also lowers BP.⁸⁷ Lifestyle and diet-induced perturbations of the microbiota and its metabolites can directly affect epithelial cell and immune cell homeostasis.^{9,46,48,88–90} Our understanding of the connection between nutrition, the microbiota and microbial products, the immune system, and host health or disease is still in its infancy.^{4,91}

MICROBIOTA-DERIVED SCFAs

One of the most well-characterized microbiota-derived metabolite classes is SCFAs, which are produced during the process of fermentation of undigestible fibers. Acetate, propionate, and butyrate are the 3 SCFAs of high abundance. Dietary fiber is a hypernym for dietary carbohydrates made up of ≥ 3 monomers, such as non-starch polysaccharides, resistant starches, inulin, pectins, β -glucan, and oligosaccharides. Most of these fibrous compounds are digested by microbes from the Bacteroidetes, Firmicutes and Actinobacteria phyla. The species *Bifidobacterium adolescentis*, *Eubacterium rectale*, *Eubacterium hallii*, *Faecalibacterium prausnitzii*, and *Ruminococcus bromii* typically colonize in the LI and possess enzymes to digest fibers for SCFA production. The LI has about 4-fold higher propionate and butyrate levels compared with SI.⁹² SCFAs are rapidly absorbed in the colon,⁹³ and butyrate is utilized to a large extent as fuel to supply energy to colonic epithelial cells.⁹⁴ Of note,

intestinal SCFAs are much higher compared with portal blood, whereas SCFAs are higher in the portal, then hepatic blood, and least in the peripheral blood, suggesting SCFAs are substantially taken up by the liver.⁹² Propionate uptake by the liver serves as precursor for gluconeogenesis, liponeogenesis, and protein synthesis,⁹⁵ while acetate enters the circulation and is metabolized by several tissues and is a substrate for cholesterol synthesis.⁹⁶ SCFAs can bind to the G-protein-coupled receptors Gpr41 (G-protein-coupled receptor 41), Gpr43 (G-protein-coupled receptor 43), Gpr109a (G-protein-coupled receptor 109 A), Olfr558 (olfactory receptor 558), and Olfr78 (olfactory receptor 78) in mice, which are also known as FFARs (free fatty acid receptors).⁹⁷ FFARs can be found in a variety of tissues, including the vasculature and kidney, and are involved in the regulation of vasoreactivity in response to propionate, acetate, and butyrate.⁹⁸ SCFA-induced G-protein-coupled receptor signal transduction has been elegantly reviewed by Poll et al.⁹⁷ Of note, Gpr41 and Olfr78 are both involved in the regulation of BP, although they seem to promote opposing actions.⁹⁹ Renin secretion is induced upon Olfr78 activation.⁹⁹ In line with this, Gpr41 knockout mice are hypertensive, while Olfr78 null animals are hypotensive.¹⁰⁰ Interestingly, acetate was formerly used in hemodialysis buffers but was largely abandoned because of its hypotensive action,¹⁰¹ consistent with the concept that for the most part, SCFAs lower BP. Even though epidemiological and interventional studies have suggested that dietary fiber intake lowers BP, the daily fiber intake in Western societies is often below the recommendations found in nutritional guidelines.^{102–104} Fiber itself has been suggested to shape microbiome composition to some extent. With regard to BP, the stimulatory action of fibers increases the abundance of the SCFA producers *F. prausnitzii* and *E. rectale*, as well as *Lactobacillus* spp., and possesses bifidogenic properties.^{105,106} A landmark study showed that European children consuming a Western diet have significantly lower SCFA levels and a high Firmicutes-to-Bacteroidetes (F/B) ratio compared with African children, who traditionally had an unprocessed diet high in fiber.¹⁰⁷ Since that study, a high F/B ratio is often used a surrogate marker of gut dysbiosis, although some Firmicutes bacteria are also known produce microbial metabolites that contribute to a healthy microbiome. Likewise, experimental work often relies on the F/B ratio as a disease marker. SHR and the stroke-prone SHR strain show an increase in F/B ratio, supporting the concept that this can act as a marker for gut dysbiosis.⁶⁴

BP and SCFA in Health and Disease

Various experimental or clinical studies have demonstrated the effects of prebiotic higher fiber or postbiotic SCFA treatment on BP. Pluznick et al⁹⁹ reported that propionate induces an acute dose-dependent hypotensive

response in anesthetized mice, which was mediated by Gpr41. Kaye et al⁶⁶ elegantly showed that prebiotic fiber not only prevents CVD but deficiencies in these nutrients may be a risk factor for the development of hypertension and CVD. The addition of the postbiotics acetate, propionate, or butyrate to a low-fiber diet was also found to improve BP and reduce target organ damage.⁶⁶ Further, FMT in GF mice demonstrated that the gut microbiome resultant of a diet lacking resistant starch compared with the high-fiber situation not only resulted in higher BP upon angiotensin II challenge but also contributed to the pathogenesis of cardiac and renal damage.⁶⁶ Similarly, our group recently tested the properties of oral propionate treatment in hypertensive mice, with and without atherosclerosis.¹⁰⁸ Propionate treatment decreased systemic and local inflammatory responses, BP, and cardiac damage in both models.¹⁰⁸ The therapeutic effects of propionate were mediated by Treg cells.¹⁰⁸ In our study, the BP-lowering potential of propionate was not acute but occurred over time, suggesting that anti-inflammatory properties of SCFAs indirectly contributed to the improvement of the vascular phenotype.¹⁰⁸ Furthermore, others have shown that Th17 cells and the balance of Th17 to Tregs mediate the role of SCFAs in BP regulation.^{109,110}

Human studies on the role of SCFAs on BP are rather rare. Some studies of microbiota composition and hypertension have indicated SCFA producers *Ruminococcaceae* spp., *Rothia*, or *Roseburia* spp. were associated with lower BP.^{57–59,61,62} In a small intervention trial, the postbiotic butyrate (600 mg/d), the prebiotic inulin (10 g/d), and the combination of these two all reduced diastolic BP in patients with metabolic syndrome.¹¹¹ In the HELIUS cohort,⁵⁸ machine learning algorithms applied to microbiome data identified *Roseburia* spp. to account for the largest absolute effect on BP. The highest tertile of *Roseburia* spp. abundance was associated with a 4.1-mmHg lower systolic BP even after adjusting for confounders, including the use of medication.⁵⁸ Conversely, fecal SCFA levels were higher in patients with higher BP.⁵⁸ This positive correlation is in line with previous studies^{55,59} but seems to contradict to the negative correlation between BP and microbial SCFA producers within the gastrointestinal tract. However, fecal SCFA levels do not necessarily reflect the SCFA levels within the intestine but rather reflect the SCFA generated in the gut, which was not absorbed by the host. This notion is supported by experimental work in SHR rats demonstrating that colonic butyrate absorption into the host is reduced in experimental hypertension.¹¹² In addition, the AT1 (angiotensin II type 1) receptor blocker Candesartan—a drug frequently used for the treatment of hypertension—was found to increase *Lactobacillus* abundance and fecal SCFA levels, improve intestinal integrity, and reduce BP in SHR rats.¹¹³ In the human MetaCardis cohort,^{114,115} depletion of butyrate production genes in the gut of severely obese subjects was ameliorated by

Candesartan treatment (Forslund et al, in revision). Altogether in the HELIUS cohort, machine learning models based on gut microbiota composition explained 4.4% and 4.3% of systolic and diastolic BP variability, respectively.

Fiber-derived SCFAs do not solely affect BP but also play a pivotal role in other CVDs and autoimmunity.^{46,60,99,100,108,116–122} For instance, postbiotic treatment with acetate, propionate, or butyrate ameliorates acute kidney injury.¹²¹ Renoprotection was associated with a reduced local and systemic inflammatory response, oxidative cellular stress, and apoptosis.¹²¹ In an animal model of multiple sclerosis—a T cell-mediated inflammatory disease of the central nervous system—propionate increased the frequency of anti-inflammatory Tregs in the gut and spleen, and this was accompanied by an amelioration of clinical symptoms.⁴⁶ High fiber intake and increased SCFA concentrations have also been shown to protect the central nervous system.¹¹⁷ Of note, patients with multiple sclerosis were shown to benefit from propionate treatment.¹²³ Short-term propionate treatment resulted in a significant and sustained enrichment of functionally competent Tregs, while type 1 helper cells and Th17 cells were depleted simultaneously.¹²³ In addition, SCFA supplementation or high fiber intake positively influences outcomes in rheumatoid arthritis—a chronic inflammatory disorder of the joints.^{116,124,125} Propionate can increase bone mass, and SCFAs were found to stimulate bone formation by increasing the number of Treg cells.¹¹⁶

Interplay Between SCFA and the Immune System

Mechanistically, SCFAs can affect distinct immune cell populations. For instance, neutrophils were found to produce less inflammatory cytokines upon propionate and butyrate treatment.¹²⁶ Butyrate can also reduce oxidative stress and phagocytic capacity.¹²⁷ SCFAs modulate the inflammatory process by decreasing dendritic cell maturation and inhibiting CD4 (cluster of differentiation 4) and CD8 (cluster of differentiation 8) T-cell proliferation.¹²¹ In contrast to acetate, butyrate or propionate affects dendritic cell maturation from bone marrow precursor cells by HDAC (histone deacetylase) inhibition.¹²⁸ Moreover, butyrate was shown to provoke M1 macrophages to secrete fewer inflammatory cytokines and increases secretion of the anti-inflammatory cytokine IL-10.¹²⁹

SCFAs also elicit the expression of anti-inflammatory signatures in human monocytes and T cells. For example, butyrate inhibits IL-12 production in *Staphylococcus aureus*—stimulated human monocytes and enhances the secretion of IL-10.¹³⁰ Recently, we have demonstrated that propionate decreases the rate of Th17 cell differentiation.^{46,123} Butyrate was also found to increase the secretion of IL-10 in type 1 helper cell-differentiated cells via Gpr43.¹³¹ SCFA-driven induction of IL-10 activates STAT3 (signal transducer and activator of transcription

3) and mTOR (mechanistic target of rapamycin) and consequently upregulates transcription factor B lymphocyte-induced maturation protein 1.¹³¹ Further, one of the most well-studied properties of SCFAs is their role in the induction of anti-inflammatory Tregs. We and others showed that butyrate and propionate increase the differentiation of murine and human Tregs and enhance their suppressive capacity.^{46,50,123,132} In addition to butyrate, de novo Treg cell formation in the periphery was also induced by propionate, but not acetate, via HDAC inhibition.¹³² Of note, *Clostridia*—a dominant class of commensal microbe—mediated the induction of colonic Tregs,⁵⁰ which is in line with findings that *Clostridium butyricum* induces Tregs and reduces Th17 cells, thereby ameliorating symptoms in experimental autoimmunity.¹³³

Fasting as a Novel Treatment Strategy for BP Control

Accumulating evidence suggests that fasting is an effective tool to manage metabolic and inflammatory diseases.⁸⁷ The rationale that caloric restriction impacts the microbiome is intriguing; nevertheless, robust data in humans are still scarce.^{90,134–137} Mesnage et al⁹⁰ investigated the effect of a 10-day periodic fast on the fecal microbiota of 15 healthy men. Fasting caused a decrease in *Lachnospiraceae* and *Ruminococcaceae*.⁹⁰ A small human pilot study showed that Ramadan fasting affected the microbiome of healthy subjects enriching several SCFA producers.^{135,138} We evaluated the impact of fasting in metabolic syndrome patients. In a clinical study, 35 metabolic syndrome patients underwent a 5-day fast followed by 3 weeks of DASH diet. A control group received DASH diet only. Fasting followed by the DASH diet reduced BP, need for antihypertensive medication, and body weight 3 months post-intervention and altered the gut microbiome impacting SCFA producers. Stratification of the cohort to BP responsiveness showed that immune cell changes present in the fasting arm are more pronounced in BP responders than in nonresponders. Furthermore, the immune shift in the fasting arm was fundamentally different from observed changes seen in the DASH arm. A BP responder-specific microbiome change in the fasting arm post-intervention (enrichment of *F. prausnitzii*, *Bacteroides*, and Firmicutes; depletion of *Actinomyces*) was observed. Of note, the enrichment of the butyrate producer *F. prausnitzii* remained even 3 months post-fasting. Computational analyses revealed that BP responders and nonresponders not only reacted differentially to fasting but differed at baseline with regard to their propionate synthesis capacity. Applying machine learning algorithms to either baseline immunome or 16S microbiome data, we were able to make effective predictions regarding which patients would respond to fasting with a reduction in BP. The prediction model was confirmed by reanalyzing the data from the only other existing

cohort (the Mesnage dataset) investigating the effect of fasting and BP.⁹⁰ A significant long-term BP decrease in the Mesnage cohort was predicted with about 70% accuracy, further supporting the idea that these findings are likely generalizable. It is important to emphasize that the aforementioned studies established associations between the microbiome and BP in response to fasting. Fasting is a daunting challenge for many patients. Being able to manipulate the mechanisms responsible for the change in BP in response to fasting would be of high clinical utility. However, few studies have succeeded in establishing mechanisms in microbiome research, which hold up in a clinical setting.¹³⁹ The high-profile failure of SER-109—a stool-derived bacterial spore mixture aimed at recurrent *Clostridium difficile* infections—during phase 2 trials is a prime example.^{139,140} The pharmacological was designed for a different disease state; however, the translation challenges facing the microbiome field at large also apply in the context of hypertension.

FROM MICE TO MEN: TRANSLATIONAL CHALLENGES

As detailed throughout this review, the host-microbiome interaction is clearly influential in human health and disease. The interactions between the host and its various microbial inhabitants are reciprocal in nature, meaning the host genetics, microbial genetics, and the interplay of these two spaces all could be explanations for the resultant phenotype.¹⁴¹ Model systems are often used in basic and preclinical hypertension research to study the pathogenesis and progression of disease. Mouse and rat models are extremely informative and can provide information that would not be available from a human cohort study. However, there are many barriers to the study of the human host-microbiome interaction in model systems.

There are many aspects of gastrointestinal physiology and morphology, which are distinct between human and rodent species.¹⁴ The cecum is perhaps the most obvious example of divergent speciation in humans and rodents, as well as the thinness of the mucus layer in the colon of mice compared with humans.¹⁴ Indeed, the genetics of model systems themselves is a double-edged sword; inbreeding and genetic homogeneity provides an easy platform for genetic manipulation, but one may question the utility of this oversimplification for translational research. The diversity of human genomes indelibly impacts individual susceptibility to disease, and the interactions between host and microbial genes is a growing area of research for the treatment of complex ailments such as type 1 diabetes and Crohn disease.¹⁴² Additionally, the genomic responses to inflammatory stressors in mice and humans are starkly contrasting, which could be related to either host-specific or microbiome-specific

features or combination of the two.¹⁴³ The microbes inhabiting the gastrointestinal tract are often also distinct between mice and humans. Although they may be comparable on the genus or phylum level, species-specific changes are often shown to have clinical importance in hypertension.¹⁴⁴ Additionally, it is known that within each microbial clade, the degree to which functional properties are shared between member species differs.¹⁴⁵ For example, *Firmicutes* species are particularly metabolically inconsistent as a clade,¹⁴⁵ which again presents an issue when considering the ubiquitous use of the F/B ratio as a marker for dysbiosis. Furthermore, because of the isolation of laboratory-raised mouse communities for extended periods of time, which is frequently unavoidable, the microbiome and associated immune competency of substrains can also be divergent from one another.^{146–148} Interestingly, a recent study found that within 2 substrains of mice with subtle differences in the host genome, colonization with an inflammatory bowel disease–associated *Bacteroides* species induced disease in only one mouse strain, despite robust colonization of all mice.¹⁴⁹ Littermate controls have, therefore, become an important standardization technique for microbiome studies.^{141,150} Use of littermate controls can provide important context, for example, 2 recent studies suggested a role for Nlrp6- and ASC (adaptor apoptosis-associated speck-like protein)-mediated inflammasomes in shaping commensal gut microbiota composition,^{151,152} although none of these results were reproducible when the appropriate littermate controls were used.¹⁵³ Additionally, the coprophagic nature of rodent models has been suggested to have a unique impact on the microbiome, which can be avoided using single housing strategies, although this induces a stress response in mice, adding an additional confounding factor.

Many researchers attempt to circumvent the issue of noncomparable species in mice compared with humans by using human microbes to colonize mice. This presents 2 important challenges, which should be considered. First, there is the issue of the reciprocal nature of interactions between the host and its microbes, of which the host is unavoidably different when using rodent models. Indeed, the importance of this interaction was exemplified in a recent study that showed that the colonization of GF mice with human or rat microbes did not induce immune maturation and only mouse-specific microbes were able to induce full immune competence.¹⁵⁴ Several differences have been noted between mouse and human immune composition, which may be related to the immune-microbiome axis, for example, the proportion of neutrophils in the peripheral blood is about twice as high (50%–70%) in humans compared with mice (10%–25%).^{155,156} Furthermore, the distribution of CD8+ T cells in nonlymphoid tissue in adult humans is much higher than what is found in specific pathogen-free mice, which could have implications for the progression of intracellular

infections or cancer.^{157,158} This is not surprising considering the coevolution of hosts and their respective microbiomes.^{141,156} Human and mice only share about 15% of bacterial lineages.¹⁵⁹ Due to the niche-specific specialization of microbes,¹⁴⁵ even implantation of defined species may not induce the desired effect and can vary in effect depending on the rodent model used.¹⁴ Second, given that the human microbiome is so highly diverse, how should one determine which human microbiome to use for experimentation? Protocols in this sense vary between studies.¹⁶⁰ Standardized procedures have been suggested throughout the literature,^{160,161} such as household member or internal baseline controls, but have not been adopted universally. Despite these caveats, mice implanted with a microbiome grafted from diseased patients have been found to recapitulate clinical findings in several disease states, including in hypertension.⁵⁶

FROM MICE TO MEN: FUTURE PERSPECTIVES

Despite the ubiquitousness of using specific pathogen-free inbred rodent models for disease research, alternative approaches are gaining traction. The intention to avoid infections that may alter experimental results is logically sound. However, there is mounting evidence that suggests humans are relatively dirty and that exposure to a full range of fungi, viruses, microbes, and so on is needed for the robust development of a healthy immune system.^{162,163} Indeed, a recent interventional study exposed children whose environment was otherwise highly hygienic to increased microbial biodiversity, which led to an increase in the amount of healthy commensal bacteria in the gut and skin, and a shift toward anti-inflammatory cytokine production in the skin.¹⁶⁴ It has been suggested that preclinical modeling of immune response in rodents likely contributes to the high failure rate in clinical trials.^{157,165} Humans are acutely and chronically infected with pathogens throughout their lifetime, which shapes their immune systems.^{166,167} Furthermore, infections can also alter reactivity to vaccinations and subsequent unrelated infections, and it was recently shown that sequential infection in mice can recapitulate these effects.¹⁶⁸

Several studies have shown that using a mouse with a more natural environment or mice captured from the wild are different and may more accurately recapitulate human physiology than laboratory-raised specific pathogen-free mice.^{165,169,170} It has been proposed that the scientific community may be able to improve preclinical pipelines with the use of mice with a wild mouse microbiome, rich in both commensals and opportunistic pathogens that are not normally present in laboratory-raised mice. Wild mice that were selected for their close similarity in genomic background were shown to

have a distinct immune profile compared with laboratory mice.¹⁷¹ The applicability of a strategy involving the capture of mice from the wild before experimentation is unlikely to be adopted widely. However, a recent study demonstrated that implantation of the full breadth of microbes from wild mice onto a C57BL/6NTac background (referred to as wildling mice) generates equally promising results (Figure 3).¹⁶⁵ Wildling mice, compared with standard laboratory-raised mice, had much higher concordance of mouse data with the results of clinical trials¹⁶⁵ and were more resistant to disease.¹⁶⁹ The wildling mouse microbiome was also more resilient to a high-fat dietary challenge over the course of 10 weeks than their laboratory mouse counterparts, whose microbiome drastically shifted in response to the challenge and was not able to fully recover.¹⁶⁵ Laboratory mice in this regard are divergent from humans because of the relative stability of the human microbiome over time.⁵ Additionally, it has been shown that the introduction of a wild microbiome leads to the convergence of laboratory-raised mouse microbial communities.¹⁶⁵ Cohousing experiments of laboratory mice with pet store mice also shifted the immune cell profile toward that of the wild mouse, which was more like the adult human phenotype.¹⁵⁷ Interestingly, wild mice also have a thicker mucus layer in the colon than laboratory mice.¹⁷² These findings suggest the potential of this technique, particularly the wildling strategy, to be used to standardize the research into the host-microbiome interaction, thereby improving reproducibility.

IMPORTANT CONSIDERATIONS FOR ONGOING RESEARCH

Throughout this review, we have addressed many issues and offered solutions to ongoing problems in the study of the microbiome. Nevertheless, microbiome science would also benefit from healthy doses of skepticism.¹⁷³ Indeed, both animal and human studies present many challenges for ongoing research. In mouse studies, standardization of procedures throughout the field, such as the use of littermate controls and robust documentation of conditions that could impact the microbiome, is essential. Factors like caging, bedding, and diet are likely to substantially impact results.^{174,175} Furthermore, it has been recently shown that the microbiome has a diurnal circadian rhythm, in which case, time of sample collection should become a consideration.¹⁷⁶ The site of sampling the microbiome is also important, as the microbial cell density, composition, and the production of microbial metabolites varies throughout the gastrointestinal tract.¹³ A recent study demonstrated that in addition to the spatial dynamics of different gastrointestinal regions, the microbiome sampled from the mucosa and the luminal space were unique in mice and in men.¹⁷⁷

Because the gastrointestinal tract is the site of action for polarization of immune cells and absorption of microbially produced metabolites,¹⁷⁸ many have questioned whether fecal sampling is the right avenue for studying the host-microbiome interface. Feces represents the excretory products from this system. However, fecal sampling is the most common and practically applicable way to examine the microbiome, particularly for longitudinal studies where noninvasive methods are necessitated. The collection of fecal matter no doubt substantially contributes to our understanding of the host-microbiome interaction. Although the relevance of locally produced microbial biproducts is suspected to be of importance, particularly impacting the uptake of metabolites to the circulation and affecting the activity of gastrointestinal immune cells, measurements of this compartment are underdeveloped. The ability to identify microbially produced compounds at the site of action in the interstitial fluid would likely provide a different perspective on host-microbiome dynamics.

Microbiome analysis adds another layer of complexity to experimental design. Particularly in human studies, procedural standardization is necessary to increase reproducibility and accelerate progress. Recent studies have suggested that sample collection,¹⁷⁹ storage,¹⁸⁰ and extraction methods^{181,182} can all contribute to resultant microbiome quantification. Particularly, the extraction method appears to play a major role in the overall microbiome signature, as it was found that the extraction method used resulted in taxonomic shifts mimicking important biologically relevant features like the enterotype or diet.¹⁸² Two sequencing methods, 16s rRNA sequencing and shotgun metagenomic sequencing, can be used to understand the composition of the microbiome. Recent findings suggest results from 16s and shotgun are largely concordant, regardless of which technology is selected,¹⁸³ although importantly the resolution of shotgun is better than 16s. The main differences are that 16s sequencing only allows genus-level taxonomic resolution, whereas shotgun offers species-level information and that shotgun data additionally can be used to assess the functional potential of the microbiome.¹⁸⁴ Similar inferences can be made from 16s using projected annotations (eg, using PICRUSt¹⁸⁵); however, this is not as reliable or robust and does not necessarily match up with comparable annotation from shotgun data.¹⁸³ However, it should also be noted that 16s is more cost-effective than shotgun, and in the case that high-resolution data are not crucial, this remains a viable option for sequencing.¹⁸⁴ Choice of software pipeline, settings, and particularly reference databases also may have impact on results, though substantially less than biosample handling. However, when comparing multiple datasets, the analysis pipeline used for either 16s or shotgun data must be carefully considered so as not to introduce a source of bias. Beyond this, investigators




Figure 3. Modeling the human immune phenotype for basic and preclinical research.

The mammalian phenotype is driven by the combination of the host genome and the microbial genome (microbiome), together referred to as the metagenome. However, the repertoire of microbes encountered in the wild is not replicated in a laboratory setting. This can substantially distort how the immune system develops and functions, leading to false assumptions of how our own wild immune system works. Thus, laboratory mice are too far removed from natural environmental conditions to faithfully mirror the physiology of free-living mammals such as humans. To address this shortcoming, embryos of laboratory mice can be transferred into wild mice to generate wildlings that more closely resemble the natural mammalian metaorganism with coevolved microbes and pathogens, while preserving the research benefits of tractable genetics of laboratory mice (interventionalist approaches, mechanistic studies, etc). Natural microbiota has been shown to be multigenerationally stable and resilient against environmental challenges, thereby providing a model system for long-term work and reproducible experimentation. Moreover, in 2 preclinical trials,^{188,189} where conventional laboratory mice as well as rat and nonhuman primate models had failed to predict the human response to harmful drug treatments, wildlings accurately mirrored the human phenotype.¹⁶⁵ Such models may enhance the validity and reproducibility of biomedical studies among research institutes, facilitate the discovery of disease mechanisms and treatments that cannot be studied in conventional laboratory mice, and increase the safety and the success of translating results from animal models to humans.

must also ask themselves whether the quantification of absolute or relative abundance of microbes is best able to answer their research question. Because the variation in DNA yield from sequenced samples can influence relative abundance quantification, it is important to interpret these data with caution to avoid false positives (including through methods such as rarefaction or explicit modeling of proportions), and it may be useful to verify findings using absolute abundances, considering again possible sources of error.^{161,186}

Ultimately, results must always be interpreted in the context with which they were collected. Most microbiome research to date (particularly human studies) suggests associations between phenotypic data and microbes or metrics of microbial diversity. To move the field forward and begin to address whether the microbiome can be targeted or manipulated to influence the prevalence or progression of hypertension, studies that can establish causation and uncover mechanisms are urgently needed. The two areas that are promising in terms of the development of novel treatment strategies are the targeting of microbially produced molecules (such as SCFA) and modulation of the microbiome-immune axis.¹³⁹ Considering the challenges in preclinical research using rodent models, we suggest the use of novel strategies to address these ongoing inquiries.

In conclusion, although prudence is needed in the interpretation of microbiome data,¹⁷³ the study of the microbiome-host interface in hypertension is a promising and rapidly accelerating field of research. With a variety of opportunities for further advancement, we anticipate that both pharmacological and lifestyle-centered treatment options addressing the microbiome space are likely to emerge in the not-so-distant future.

ARTICLE INFORMATION

Affiliations

Experimental and Clinical Research Center, a cooperation of Charité-Universitätsmedizin Berlin and Max Delbrück Center for Molecular Medicine, Berlin, Germany (E.G.A., H.B., A.M., L.M., N.W., S.K.F., D.N.M.). Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany (H.B., A.M., L.M., N.W., S.K.F., D.N.M.). For Max Delbrück Center for Molecular Medicine in the Helmholtz Association, Berlin, Germany (E.G.A., H.B., N.W., S.K.F., D.N.M.). DZHK (German Centre for Cardiovascular Research), partner site Berlin, Germany (E.G.A., H.B., A.M., L.M., N.W., S.K.F., D.N.M.). For Berlin Institute of Health at Charité-Universitätsmedizin Berlin, Berlin, Germany (N.W.). Freie Universität Berlin, Department of Biology, Chemistry, Pharmacy, Berlin, Germany (E.G.A.). Department of Biomedicine, University of Bergen, Norway (H.W.). Medical Center-University of Freiburg, Department of Medicine II, Gastroenterology, Hepatology, Endocrinology, and Infectious Diseases, Freiburg, Germany (S.P.R.).

Sources of Funding

D.N. Müller, H. Bartolomaeus, N. Wilck, and S.K. Forslund were supported by the Deutsche Forschungsgemeinschaft (German Research Foundation; Projektnummer 394046635-SFB 1365); D.N. Müller was supported by the German Center for Cardiovascular Research (81Z0100106); N. Wilck was supported by the European Research Council under the European Union Horizon 2020 Research and Innovation Program (NW: 852796); N. Wilck is supported by a grant from the Corona-Stiftung; N. Wilck is a participant in the Clinician Scientist Program funded by the Berlin Institute of Health. Financial support from the Research Council of Norway (project number 262079), the Norwegian Health Association, and from

the Western Norway Regional Health Authority (project number 912168) to H. Wiig is gratefully acknowledged. S.P. Rosshart was supported by the Deutsche Forschungsgemeinschaft DFG (German Research Foundation; Emmy Noether-Programm RO 6247/1-1 and SFB 1160 IMPATH).

Disclosures

S.P. Rosshart declares no competing interests and discloses that Taconic Biosciences licensed WildR mice with natural gut microbiota from NIDDK.

REFERENCES

- Sender R, Fuchs S, Milo R. Revised estimates for the number of human and bacteria cells in the body. *PLoS Biol*. 2016;14:e1002533. doi: 10.1371/journal.pbio.1002533
- Tierney BT, Yang Z, Luber JM, Beaudin M, Wibowo MC, Baek C, Mehlenbacher E, Patel CJ, Kostic AD. The landscape of genetic content in the gut and oral human microbiome. *Cell Host Microbe*. 2019;26:283–295. e8. doi: 10.1016/j.chom.2019.07.008
- Human Microbiome Project C. Structure, function and diversity of the healthy human microbiome. *Nature*. 2012;486:207–214. doi: 10.1038/nature11234
- Zhang LS, Davies SS. Microbial metabolism of dietary components to bioactive metabolites: opportunities for new therapeutic interventions. *Genome Med*. 2016;8:46. doi: 10.1186/s13073-016-0296-x
- David LA, Materna AC, Friedman J, Campos-Baptista MI, Blackburn MC, Perrotta A, Erdman SE, Alm EJ. Host lifestyle affects human microbiota on daily timescales. *Genome Biol*. 2014;15:R89. doi: 10.1186/gb-2014-15-7-r89
- Thompson AL, Monteagudo-Mera A, Cadenas MB, Lampl ML, Azcarate-Peril MA. Milk- and solid-feeding practices and daycare attendance are associated with differences in bacterial diversity, predominant communities, and metabolic and immune function of the infant gut microbiome. *Front Cell Infect Microbiol*. 2015;5:3. doi: 10.3389/fcimb.2015.00003
- Forslund K, Hildebrand F, Nielsen T, Falony G, Le Chatelier E, Sunagawa S, Prifti E, Vieira-Silva S, Gudmundsdottir V, Pedersen HK, et al; MetaHIT Consortium. Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. *Nature*. 2015;528:262–266. doi: 10.1038/nature15766
- Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, Liang S, Zhang W, Guan Y, Shen D, et al. A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature*. 2012;490:55–60. doi: 10.1038/nature11450
- Koeth RA, Wang Z, Levison BS, Buffa JA, Org E, Sheehy BT, Britt EB, Fu X, Wu Y, Li L, et al. Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat Med*. 2013;19:576–585. doi: 10.1038/nm.3145
- Tang WH, Wang Z, Fan Y, Levison B, Hazen JE, Donahue LM, Wu Y, Hazen SL. Prognostic value of elevated levels of intestinal microbe-generated metabolite trimethylamine-N-oxide in patients with heart failure: refining the gut hypothesis. *J Am Coll Cardiol*. 2014;64:1908–1914. doi: 10.1016/j.jacc.2014.02.617
- Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, Wu Y, Hazen SL. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med*. 2013;368:1575–1584. doi: 10.1056/NEJMoa1109400
- Donia MS, Fischbach MA. HUMAN MICROBIOTA. Small molecules from the human microbiota. *Science*. 2015;349:1254766. doi: 10.1126/science.1254766
- Mowat AM, Agace WW. Regional specialization within the intestinal immune system. *Nat Rev Immunol*. 2014;14:667–685. doi: 10.1038/nri3738
- Nguyen TL, Vieira-Silva S, Liston A, Raes J. How informative is the mouse for human gut microbiota research? *Dis Model Mech*. 2015;8:1–16. doi: 10.1242/dmm.017400
- Brown K, Abbott DW, Uwiera RRE, Inglis GD. Removal of the cecum affects intestinal fermentation, enteric bacterial community structure, and acute colitis in mice. *Gut Microbes*. 2018;9:218–235. doi: 10.1080/19490976.2017.1408763
- Marteau P, Pochart P, Doré J, Béra-Maillet C, Bernalier A, Corthier G. Comparative study of bacterial groups within the human cecal and fecal microbiota. *Appl Environ Microbiol*. 2001;67:4939–4942. doi: 10.1128/aem.67.10.4939-4942.2001
- Stevens CE, Hume ID. Contributions of microbes in vertebrate gastrointestinal tract to production and conservation of nutrients. *Physiol Rev*. 1998;78:393–427. doi: 10.1152/physrev.1998.78.2.393

18. Lievin-Le Moal V, Servin AL. The front line of enteric host defense against unwelcome intrusion of harmful microorganisms: mucins, antimicrobial peptides, and microbiota. *Clin Microbiol Rev*. 2006;19:315–337. doi: 10.1128/CMR.19.2.315–337.2006
19. Derrien M, Vaughan EE, Plugge CM, de Vos WM. *Akkermansia muciniphila* gen. nov., sp. nov., a human intestinal mucin-degrading bacterium. *Int J Syst Evol Microbiol*. 2004;54:1469–1476. doi: 10.1099/ijs.0.02873-0
20. Collado MC, Derrien M, Isolauri E, de Vos WM, Salminen S. Intestinal integrity and *Akkermansia muciniphila*, a mucin-degrading member of the intestinal microbiota present in infants, adults, and the elderly. *Appl Environ Microbiol*. 2007;73:7767–7770. doi: 10.1128/AEM.01477-07
21. Camilleri M. Leaky gut: mechanisms, measurement and clinical implications in humans. *Gut*. 2019;68:1516–1526. doi: 10.1136/gutjnl-2019-318427
22. Desai MS, Seekatz AM, Koropatkin NM, Kamada N, Hickey CA, Wolter M, Pudlo NA, Kitamoto S, Terrapon N, Muller A, et al. A dietary fiber-deprived gut microbiota degrades the colonic mucus barrier and enhances pathogen susceptibility. *Cell*. 2016;167:1339–1353.e21. doi: 10.1016/j.cell.2016.10.043
23. Krebs CF, Paust HJ, Krohn S, Koyro T, Brix SR, Riedel JH, Bartsch P, Wiech T, Meyer-Schwesinger C, Huang J, et al. Autoimmune renal disease is exacerbated by S1P-receptor-1-dependent intestinal Th17 cell migration to the kidney. *Immunity*. 2016;45:1078–1092. doi: 10.1016/j.immuni.2016.10.020
24. Hart AL, Ng SC, Mann E, Al-Hassi HO, Bernardo D, Knight SC. Homing of immune cells: role in homeostasis and intestinal inflammation. *Inflamm Bowel Dis*. 2010;16:1969–1977. doi: 10.1002/ibd.21304
25. Hayes CL, Dong J, Galipeau HJ, Jury J, McCarville J, Huang X, Wang XY, Naidoo A, Anbazhagan AN, Libertucci J, et al. Commensal microbiota induces colonic barrier structure and functions that contribute to homeostasis. *Sci Rep*. 2018;8:14184. doi: 10.1038/s41598-018-32366-6
26. MacDonald TT, Carter PB. Requirement for a bacterial flora before mice generate cells capable of mediating the delayed hypersensitivity reaction to sheep red blood cells. *J Immunol*. 1979;122:2624–2629.
27. Ivanov II, Atarashi K, Manel N, Brodie EL, Shima T, Karaoz U, Wei D, Goldfarb KC, Santee CA, Lynch SV, et al. Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell*. 2009;139:485–498. doi: 10.1016/j.cell.2009.09.033
28. Geva-Zatorsky N, Sefik E, Kua L, Pasman L, Tan TG, Ortiz-Lopez A, Yanortsang TB, Yang L, Jupp R, Mathis D, et al. Mining the human gut microbiota for immunomodulatory organisms. *Cell*. 2017;168:928–943.e11. doi: 10.1016/j.cell.2017.01.022
29. Schluter J, Peled JU, Taylor BP, Markey KA, Smith M, Taur Y, Niehus R, Staffas A, Dai A, Fontana E, et al. The gut microbiota is associated with immune cell dynamics in humans. *Nature*. 2020;588:303–307. doi: 10.1038/s41586-020-2971-8
30. Gensollen T, Iyer SS, Kasper DL, Blumberg RS. How colonization by microbiota in early life shapes the immune system. *Science*. 2016;352:539–544. doi: 10.1126/science.aad9378
31. Russell SL, Gold MJ, Hartmann M, Willing BP, Thorson L, Wlodarska M, Gill N, Blanchet MR, Mohn WW, McNagny KM, et al. Early life antibiotic-driven changes in microbiota enhance susceptibility to allergic asthma. *EMBO Rep*. 2012;13:440–447. doi: 10.1038/embor.2012.32
32. Zhang X, Zhivaki D, Lo-Man R. Unique aspects of the perinatal immune system. *Nat Rev Immunol*. 2017;17:495–507. doi: 10.1038/nri.2017.54
33. Wilck N, Balogh A, Markó L, Bartolomeus H, Müller DN. The role of sodium in modulating immune cell function. *Nat Rev Nephrol*. 2019;15:546–558. doi: 10.1038/s41581-019-0167-y
34. Libby P, Pasterkamp G, Crea F, Jang IK. Reassessing the mechanisms of acute coronary syndromes. *Circ Res*. 2019;124:150–160. doi: 10.1161/CIRCRESAHA.118.311098
35. Xiao L, Harrison DG. Inflammation in hypertension. *Can J Cardiol*. 2020;36:635–647. doi: 10.1016/j.cjca.2020.01.013
36. Itani HA, Harrison DG. Memories that last in hypertension. *Am J Physiol Renal Physiol*. 2015;308:F1197–F1199. doi: 10.1152/ajprenal.00633.2014
37. Itani HA, Xiao L, Saleh MA, Wu J, Pilkinton MA, Dale BL, Barbaro NR, Foss JD, Kirabo A, Montanié KR, et al. CD70 Exacerbates blood pressure elevation and renal damage in response to repeated hypertensive stimuli. *Circ Res*. 2016;118:1233–1243. doi: 10.1161/CIRCRESAHA.115.308111
38. Madhur MS, Lob HE, McCann LA, Iwakura Y, Blinder Y, Guzik TJ, Harrison DG. Interleukin 17 promotes angiotensin II-induced hypertension and vascular dysfunction. *Hypertension*. 2010;55:500–507. doi: 10.1161/HYPERTENSIONAHA.109.145094
39. Markó L, Kvakan H, Park JK, Qadri F, Spallek B, Binger KJ, Bowman EP, Kleinewietfeld M, Fokuhl V, Dechend R, et al. Interferon- γ signaling inhibition ameliorates angiotensin II-induced cardiac damage. *Hypertension*. 2012;60:1430–1436. doi: 10.1161/HYPERTENSIONAHA.112.199265
40. Didion SP, Kinzenbaw DA, Schrader LI, Chu Y, Faraci FM. Endogenous interleukin-10 inhibits angiotensin II-induced vascular dysfunction. *Hypertension*. 2009;54:619–624. doi: 10.1161/HYPERTENSIONAHA.109.137158
41. Kvakan H, Kleinewietfeld M, Qadri F, Park JK, Fischer R, Schwarz I, Rahn HP, Plehm R, Wellner M, Elitok S, et al. Regulatory T cells ameliorate angiotensin II-induced cardiac damage. *Circulation*. 2009;119:2904–2912. doi: 10.1161/CIRCULATIONAHA.108.832782
42. Caillon A, Mian MOR, Fraulob-Aquino JC, Huo KG, Barhoumi T, Ouerd S, Sinnaeve PR, Paradis P, Schiffrin EL. $\gamma\delta$ T cells mediate angiotensin II-induced hypertension and vascular injury. *Circulation*. 2017;135:2155–2162. doi: 10.1161/CIRCULATIONAHA.116.027058
43. Shah KH, Shi P, Giani JF, Janjulia T, Bernstein EA, Li Y, Zhao T, Harrison DG, Bernstein KE, Shen XZ. Myeloid suppressor cells accumulate and regulate blood pressure in hypertension. *Circ Res*. 2015;117:858–869. doi: 10.1161/CIRCRESAHA.115.306539
44. Ferguson JF, Aden LA, Barbaro NR, Van Beusecum JP, Xiao L, Simmons AJ, Warden C, Pasic L, Himmel LE, Washington MK, et al. High dietary salt-induced dendritic cell activation underlies microbial dysbiosis-associated hypertension. *JCI Insight*. 2019;5:e126241. doi: 10.1172/jci.insight.126241
45. Honda K, Littman DR. The microbiota in adaptive immune homeostasis and disease. *Nature*. 2016;535:75–84. doi: 10.1038/nature18848
46. Haghighi A, Jörg S, Duscha A, Berg J, Manzel A, Waschbisch A, Hammer A, Lee DH, May C, Wilck N, et al. Dietary fatty acids directly impact central nervous system autoimmunity via the small intestine. *Immunity*. 2015;43:817–829. doi: 10.1016/j.immuni.2015.09.007
47. Tan TG, Sefik E, Geva-Zatorsky N, Kua L, Naskar D, Teng F, Pasman L, Ortiz-Lopez A, Jupp R, Wu HJ, et al. Identifying species of symbiotic bacteria from the human gut that, alone, can induce intestinal Th17 cells in mice. *Proc Natl Acad Sci USA*. 2016;113:E8141–E8150. doi: 10.1073/pnas.1617460113
48. Wilck N, Matus MG, Kearney SM, Olesen SW, Forslund K, Bartolomeus H, Haase S, Mähler A, Balogh A, Markó L, et al. Salt-responsive gut commensal modulates TH17 axis and disease. *Nature*. 2017;551:585–589. doi: 10.1038/nature24628
49. Atarashi K, Tanoue T, Shima T, Imaoka A, Kuwahara T, Momose Y, Cheng G, Yamasaki S, Saito T, Ohba Y, et al. Induction of colonic regulatory T cells by indigenous *Clostridium* species. *Science*. 2011;331:337–341. doi: 10.1126/science.1198469
50. Furusawa Y, Obata Y, Fukuda S, Endo TA, Nakato G, Takahashi D, Nakanishi Y, Uetake C, Kato K, Kato T, et al. Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature*. 2013;504:446–450. doi: 10.1038/nature12721
51. Evangelou E, Warren HR, Mosen-Ansorena D, Mifsud B, Pazoki R, Gao H, Ntritsos G, Dimou N, Cabrera CP, Karaman I, et al. Million Veteran Program. Genetic analysis of over 1 million people identifies 535 new loci associated with blood pressure traits. *Nat Genet*. 2018;50:1412–1425. doi: 10.1038/s41588-018-0205-x
52. Pazoki R, Dehghan A, Evangelou E, Warren H, Gao H, Caulfield M, Elliott P, Tzoulaki I. Genetic predisposition to high blood pressure and lifestyle factors: associations with midlife blood pressure levels and cardiovascular events. *Circulation*. 2018;137:653–661. doi: 10.1161/CIRCULATIONAHA.117.030898
53. Lewington S, Clarke R, Qizilbash N, Peto R, Collins R; Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2002;360:1903–1913. doi: 10.1016/s0140-6736(02)11911-8
54. Dan X, Mushi Z, Baili W, Han L, Enqi W, Huanhu Z, Shuchun L. Differential analysis of hypertension-associated intestinal microbiota. *Int J Med Sci*. 2019;16:872–881. doi: 10.7150/ijms.29322
55. de la Cuesta-Zuluaga J, Mueller NT, Alvarez-Quintero R, Velasquez-Mejia EP, Sierra JA, Corrales-Agudelo V, Carmona JA, Abad JM, Escobar JS. Higher fecal short-chain fatty acid levels are associated with gut microbiome dysbiosis, obesity, hypertension and cardiometabolic disease risk factors. *Nutrients*. 2018;11:51. doi: 10.3390/nu11010051
56. Li J, Zhao F, Wang Y, Chen J, Tao J, Tian G, Wu S, Liu W, Cui Q, Geng B, et al. Gut microbiota dysbiosis contributes to the development of hypertension. *Microbiome*. 2017;5:14. doi: 10.1186/s40168-016-0222-x
57. Sun S, Lulla A, Sioda M, Winglee K, Wu MC, Jacobs DR Jr, Shikany JM, Lloyd-Jones DM, Launer LJ, Fodor AA, et al. Gut microbiota composition and blood pressure. *Hypertension*. 2019;73:998–1006. doi: 10.1161/HYPERTENSIONAHA.118.12109

58. Verhaar BJH, Collard D, Prodan A, Levels JHM, Zwinderman AH, Bäckhed F, Vogt L, Peters MJL, Muller M, Nieuwdorp M, et al. Associations between gut microbiota, faecal short-chain fatty acids, and blood pressure across ethnic groups: the HELIUS study. *Eur Heart J*. 2020;41:4259–4267. doi: 10.1093/eurheartj/ehaa704
59. Yan Q, Gu Y, Li X, Yang W, Jia L, Chen C, Han X, Huang Y, Zhao L, Li P, et al. Alterations of the gut microbiome in hypertension. *Front Cell Infect Microbiol*. 2017;7:381. doi: 10.3389/fcimb.2017.00381
60. Yang T, Santisteban MM, Rodriguez V, Li E, Ahmari N, Carvajal JM, Zadeh M, Gong M, Qi Y, Zubecevic J, et al. Gut dysbiosis is linked to hypertension. *Hypertension*. 2015;65:1331–1340. doi: 10.1161/HYPERTENSIONAHA.115.05315
61. Kim S, Goel R, Kumar A, Qi Y, Lobaton G, Hosaka K, Mohammed M, Handberg EM, Richards EM, Pepine CJ, et al. Imbalance of gut microbiome and intestinal epithelial barrier dysfunction in patients with high blood pressure. *Clin Sci (Lond)*. 2018;132:701–718. doi: 10.1042/CS20180087
62. Huart J, Leenders J, Taminiau B, Descy J, Saint-Remy A, Daube G, Krzesinski JM, Melin P, de Tullio P, Jouret F. Gut microbiota and fecal levels of short-chain fatty acids differ upon 24-hour blood pressure levels in men. *Hypertension*. 2019;74:1005–1013. doi: 10.1161/HYPERTENSIONAHA.118.12588
63. Jackson MA, Verdi S, Maxan ME, Shin CM, Zierer J, Bowyer RCE, Martin T, Williams FMK, Menni C, Bell JT, et al. Gut microbiota associations with common diseases and prescription medications in a population-based cohort. *Nat Commun*. 2018;9:2655. doi: 10.1038/s41467-018-05184-7
64. Adnan S, Nelson JW, Ajami NJ, Venna VR, Petrosino JF, Bryan RM Jr, Durgan DJ. Alterations in the gut microbiota can elicit hypertension in rats. *Physiol Genomics*. 2017;49:96–104. doi: 10.1152/physiolgenomics.00081.2016
65. Mell B, Jala VR, Mathew AV, Byun J, Wagholde H, Zhang Y, Haribabu B, Vijay-Kumar M, Pennathur S, Joe B. Evidence for a link between gut microbiota and hypertension in the Dahl rat. *Physiol Genomics*. 2015;47:187–197. doi: 10.1152/physiolgenomics.00136.2014
66. Kaye DM, Shihata WA, Jama HA, Tsyganov K, Ziemann M, Kiriazis H, Horlock D, Vijay A, Giam B, Vinh A, et al. Deficiency of prebiotic fiber and insufficient signaling through gut metabolite-sensing receptors leads to cardiovascular disease. *Circulation*. 2020;141:1393–1403. doi: 10.1161/CIRCULATIONAHA.119.043081
67. Marques FZ, Nelson E, Chu PY, Horlock D, Fiedler A, Ziemann M, Tan JK, Kuruppu S, Rajapakse NW, El-Osta A, et al. High-fiber diet and acetate supplementation change the gut microbiota and prevent the development of hypertension and heart failure in hypertensive mice. *Circulation*. 2017;135:964–977. doi: 10.1161/CIRCULATIONAHA.116.024545
68. Santisteban MM, Qi Y, Zubecevic J, Kim S, Yang T, Shenoy V, Cole-Jeffrey CT, Lobaton GO, Stewart DC, Rubiano A, et al. Hypertension-linked pathophysiological alterations in the gut. *Circ Res*. 2017;120:312–323. doi: 10.1161/CIRCRESAHA.116.309006
69. Robles-Vera I, Toral M, de la Visitación N, Sánchez M, Gómez-Guzmán M, Muñoz R, Algieri F, Vezza T, Jiménez R, Gálvez J, et al. Changes to the gut microbiota induced by losartan contributes to its antihypertensive effects. *Br J Pharmacol*. 2020;177:2006–2023. doi: 10.1111/bph.14965
70. Chen L, He FJ, Dong Y, Huang Y, Wang C, Harshfield GA, Zhu H. Modest sodium reduction increases circulating short-chain fatty acids in untreated hypertensives: a randomized, double-blind, placebo-controlled trial. *Hypertension*. 2020;76:73–79. doi: 10.1161/HYPERTENSIONAHA.120.14800
71. Toral M, Gómez-Guzmán M, Jiménez R, Romero M, Sánchez M, Utrilla MP, Garrido-Mesa N, Rodríguez-Cabezas ME, Olivares M, Gálvez J, et al. The probiotic *Lactobacillus coryniformis* CECT5711 reduces the vascular pro-oxidant and pro-inflammatory status in obese mice. *Clin Sci (Lond)*. 2014;127:33–45. doi: 10.1042/CS20130339
72. Müller DN, Wilck N, Haase S, Kleinewietfeld M, Linker RA. Sodium in the microenvironment regulates immune responses and tissue homeostasis. *Nat Rev Immunol*. 2019;19:243–254. doi: 10.1038/s41577-018-0113-4
73. Kalesi S, Sun J, Buys N, Jayasinghe R. Effect of probiotics on blood pressure: a systematic review and meta-analysis of randomized, controlled trials. *Hypertension*. 2014;64:897–903. doi: 10.1161/HYPERTENSIONAHA.114.03469
74. Maier L, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, Brochado AR, Fernandez KC, Dose H, Mori H, et al. Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature*. 2018;555:623–628. doi: 10.1038/nature25979
75. Zeevi D, Korem T, Zmora N, Israeli D, Rothschild D, Weinberger A, Ben-Yacov O, Lador D, Avnit-Sagi T, Lotan-Pompan M, et al. Personalized Nutrition by prediction of glycemic responses. *Cell*. 2015;163:1079–1094. doi: 10.1016/j.cell.2015.11.001
76. Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, Sogin ML, Jones WJ, Roe BA, Affourtit JP, et al. A core gut microbiome in obese and lean twins. *Nature*. 2009;457:480–484. doi: 10.1038/nature07540
77. Turnbaugh PJ, Ridaura VK, Faith JJ, Rey FE, Knight R, Gordon JL. The effect of diet on the human gut microbiome: a metagenomic analysis in humanized gnotobiotic mice. *Sci Transl Med*. 2009;1:6ra14. doi: 10.1126/scitranslmed.3000322
78. Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, Griffin NW, Lombard V, Henrissat B, Bain JR, et al. Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science*. 2013;341:1241214. doi: 10.1126/science.1241214
79. Christ A, Günther P, Lauterbach MAR, Duesell P, Biswas D, Pelka K, Scholz CJ, Oosting M, Haendler K, Baßler K, et al. Western diet triggers NLRP3-dependent innate immune reprogramming. *Cell*. 2018;172:162–175.e14. doi: 10.1016/j.cell.2017.12.013
80. Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, Burcelin R. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes*. 2008;57:1470–1481. doi: 10.2337/db07-1403
81. Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, Feldstein AE, Britt EB, Fu X, Chung YM, et al. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature*. 2011;472:57–63. doi: 10.1038/nature09922
82. He FJ, MacGregor GA. Reducing population salt intake worldwide: from evidence to implementation. *Prog Cardiovasc Dis*. 2010;52:363–382. doi: 10.1016/j.pcad.2009.12.006
83. Taylor J. 2013 ESH/ESC guidelines for the management of arterial hypertension. *Eur Heart J*. 2013;34:2108–2109.
84. World Health Organization. Guideline: sodium intake for adults and children. 2012;ISBN 978-92-4-150483-6.
85. European Union. Implementation of the EU Salt Reduction Framework: results of member states survey. 2012.
86. Whelton PK, Carey RM, Aronow WS, Casey DE Jr, Collins KJ, Dennison Himmelfarb C, DePalma SM, Gidding S, Jamerson KA, Jones DW, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APHA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2018;138:e484–e594. doi: 10.1161/CIR.0000000000000596
87. de Cabo R, Mattson MP. Effects of intermittent fasting on health, aging, and disease. *N Engl J Med*. 2019;381:2541–2551. doi: 10.1056/NEJMr1905136
88. Bäckhed F, Manchester JK, Semenkovich CF, Gordon JL. Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. *Proc Natl Acad Sci USA*. 2007;104:979–984. doi: 10.1073/pnas.0605374104
89. Tremaroli V, Bäckhed F. Functional interactions between the gut microbiota and host metabolism. *Nature*. 2012;489:242–249. doi: 10.1038/nature11552
90. Mesnage R, Grundler F, Schwiertz A, Le Maho Y, Wilhelmi de Toledo F. Changes in human gut microbiota composition are linked to the energy metabolic switch during 10 d of Buchinger fasting. *J Nutr Sci*. 2019;8:e36. doi: 10.1017/jns.2019.33
91. Zmora N, Suez J, Elinav E. You are what you eat: diet, health and the gut microbiota. *Nat Rev Gastroenterol Hepatol*. 2019;16:35–56. doi: 10.1038/s41575-018-0061-2
92. Cummings JH, Pomare EW, Branch WJ, Naylor CP, Macfarlane GT. Short chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut*. 1987;28:1221–1227. doi: 10.1136/gut.28.10.1221
93. Moore WE, Cato EP, Holdeman LV. Some current concepts in intestinal bacteriology. *Am J Clin Nutr*. 1978;31:S33–S42. doi: 10.1093/ajcn/31.10.S33
94. Roediger WE. Utilization of nutrients by isolated epithelial cells of the rat colon. *Gastroenterology*. 1982;83:424–429.
95. Wolever TM, Spadafora P, Eshuis H. Interaction between colonic acetate and propionate in humans. *Am J Clin Nutr*. 1991;53:681–687. doi: 10.1093/ajcn/53.3.681
96. Wolever TM, Brighenti F, Royall D, Jenkins AL, Jenkins DJ. Effect of rectal infusion of short chain fatty acids in human subjects. *Am J Gastroenterol*. 1989;84:1027–1033.
97. Poll BG, Cheema MU, Pluznick JL. Gut microbial metabolites and blood pressure regulation: focus on SCFAs and TMAO. *Physiology (Bethesda)*. 2020;35:275–284. doi: 10.1152/physiol.00004.2020
98. Muralitharan RR, Jama HA, Xie L, Peh A, Snelson M, Marques FZ. Microbial peer pressure: the role of the gut microbiota in hypertension and its complications. *Hypertension*. 2020;76:1674–1687. doi: 10.1161/HYPERTENSIONAHA.120.14473.

99. Pluznick JL, Protzko RJ, Gevorgyan H, Peterlin Z, Sipos A, Han J, Brunet I, Wan LX, Rey F, Wang T, et al. Olfactory receptor responding to gut microbiota-derived signals plays a role in renin secretion and blood pressure regulation. *Proc Natl Acad Sci USA*. 2013;110:4410–4415. doi: 10.1073/pnas.1215927110
100. Natarajan N, Hori D, Flavahan S, Stepan J, Flavahan NA, Berkowitz DE, Pluznick JL. Microbial short chain fatty acid metabolites lower blood pressure via endothelial G protein-coupled receptor 41. *Physiol Genomics*. 2016;48:826–834. doi: 10.1152/physiolgenomics.00089.2016
101. Pagel MD, Ahmad S, Vizzo JE, Scribner BH. Acetate and bicarbonate fluctuations and acetate intolerance during dialysis. *Kidney Int*. 1982;21:513–518. doi: 10.1038/ki.1982.54
102. Stephen AM, Champ MM, Cloran SJ, Fleith M, van Lieshout L, Mejbourn H, Burley VJ. Dietary fibre in Europe: current state of knowledge on definitions, sources, recommendations, intakes and relationships to health. *Nutr Res Rev*. 2017;30:149–190. doi: 10.1017/S095442241700004X
103. Reynolds A, Mann J, Cummings J, Winter N, Mete E, Te Morenga L. Carbohydrate quality and human health: a series of systematic reviews and meta-analyses. *Lancet*. 2019;393:434–445. doi: 10.1016/S0140-6736(18)31809-9
104. Hartley L, May MD, Loveman E, Colquitt JL, Rees K. Dietary fibre for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev*. 2016;2016:CD011472. doi: 10.1002/14651858.CD011472.pub2
105. Dewulf EM, Cani PD, Claus SP, Fuentes S, Puylaert PG, Neyrinck AM, Bindels LB, de Vos WM, Gibson GR, Thissen JP, et al. Insight into the prebiotic concept: lessons from an exploratory, double blind intervention study with inulin-type fructans in obese women. *Gut*. 2013;62:1112–1121. doi: 10.1136/gutjnl-2012-303304
106. Maier TV, Lucio M, Lee LH, VerBerkmoes NC, Brislawn CJ, Bernhardt J, Lamendella R, McDermott JE, Bergeron N, Heinzmann SS, et al. Impact of dietary resistant starch on the human gut microbiome, metaproteome, and metabolome. *mBio*. 2017;8:e01343-17. doi: 10.1128/mBio.01343-17
107. De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, Collini S, Pieraccini G, Lionetti P. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci USA*. 2010;107:14691–14696. doi: 10.1073/pnas.1005963107
108. Bartolomeaus H, Balogh A, Yakoub M, Homann S, Markó L, Höges S, Tsvetkov D, Krannich A, Wundersitz S, Avery EG, et al. Short-chain fatty acid propionate protects from hypertensive cardiovascular damage. *Circulation*. 2019;139:1407–1421. doi: 10.1161/CIRCULATIONAHA.118.036652
109. Robles-Vera I, Toral M, de la Visitación N, Sánchez M, Gómez-Guzmán M, Romero M, Yang T, Izquierdo-García JL, Jiménez R, Ruiz-Cabello J, et al. Probiotics prevent dysbiosis and the rise in blood pressure in genetic hypertension: role of short-chain fatty acids. *Mol Nutr Food Res*. 2020;64:e1900616. doi: 10.1002/mnfr.201900616
110. Robles-Vera I, de la Visitación N, Toral M, Sánchez M, Romero M, Gómez-Guzmán M, Yang T, Izquierdo-García JL, Guerra-Hernández E, Ruiz-Cabello J, et al. Probiotic bifidobacterium breve prevents DOCA-salt hypertension. *FASEB J*. 2020;34:13626–13640. doi: 10.1096/fj.202001532R
111. Roshanravan N, Mahdavi R, Alizadeh E, Jafarabadi MA, Hedayati M, Ghavami A, Alipour S, Alamdari NM, Barati M, Ostadrahimi A. Effect of butyrate and inulin supplementation on glycemic status, lipid profile and glucagon-like peptide 1 level in patients with type 2 diabetes: a randomized double-blind, placebo-controlled trial. *Horm Metab Res*. 2017;49:886–891. doi: 10.1055/s-0043-119089
112. Yang T, Magee KL, Colon-Perez LM, Larkin R, Liao YS, Balazic E, Cowart JR, Arocha R, Redler T, Febo M, et al. Impaired butyrate absorption in the proximal colon, low serum butyrate and diminished central effects of butyrate on blood pressure in spontaneously hypertensive rats. *Acta Physiol (Oxf)*. 2019;226:e13256. doi: 10.1111/apha.13256
113. Wu D, Tang X, Ding L, Cui J, Wang P, Du X, Yin J, Wang W, Chen Y, Zhang T. Candesartan attenuates hypertension-associated pathophysiological alterations in the gut. *Biomed Pharmacother*. 2019;116:109040. doi: 10.1016/j.biopha.2019.109040
114. Verger EO, Armstrong P, Nielsen T, Chakaroun R, Aron-Wisniewsky J, Göbel RJ, Schütz T, Delaere F, Gausseres N, Clément K, et al; MetaCardis Consortium. Dietary assessment in the metacardis study: development and relative validity of an online food frequency questionnaire. *J Acad Nutr Diet*. 2017;117:878–888. doi: 10.1016/j.jand.2016.10.030
115. Vieira-Silva S, Falony G, Belda E, Nielsen T, Aron-Wisniewsky J, Chakaroun R, Forslund SK, Assmann K, Valles-Colomer M, Nguyen TTD, et al; MetaCardis Consortium. Statin therapy is associated with lower prevalence of gut microbiota dysbiosis. *Nature*. 2020;581:310–315. doi: 10.1038/s41586-020-2269-x
116. Lucas S, Omata Y, Hofmann J, Böttcher M, Iljazovic A, Sarter K, Albrecht O, Schulz O, Krishnacoumar B, Krönke G, et al. Short-chain fatty acids regulate systemic bone mass and protect from pathological bone loss. *Nat Commun*. 2018;9:55. doi: 10.1038/s41467-017-02490-4
117. Mizuno M, Noto D, Kaga N, Chiba A, Miyake S. The dual role of short fatty acid chains in the pathogenesis of autoimmune disease models. *PLoS One*. 2017;12:e0173032. doi: 10.1371/journal.pone.0173032
118. Pluznick JL. Microbial short-chain fatty acids and blood pressure regulation. *Curr Hypertens Rep*. 2017;19:25. doi: 10.1007/s11906-017-0722-5
119. Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru C, Blanchard C, Junt T, Nicod LP, Harris NL, et al. Gut microbiota metabolism of dietary fiber influences allergic airway disease and hematopoiesis. *Nat Med*. 2014;20:159–166. doi: 10.1038/nm.3444
120. Zaiss MM, Rapin A, Lebon L, Dubey LK, Mosconi I, Sarter K, Piersigilli A, Menin L, Walker AW, Rougemont J, et al. The intestinal microbiota contributes to the ability of helminths to modulate allergic inflammation. *Immunity*. 2015;43:998–1010. doi: 10.1016/j.immuni.2015.09.012
121. Andrade-Oliveira V, Amano MT, Correa-Costa M, Castoldi A, Felizardo RJ, de Almeida DC, Bassi EJ, Moraes-Vieira PM, Hiyane MI, Rodas AC, et al. Gut bacteria products prevent AKI induced by ischemia-reperfusion. *J Am Soc Nephrol*. 2015;26:1877–1888. doi: 10.1681/ASN.2014030288
122. Bartolomeaus H, Avery EG, Bartolomeaus TUP, Kozhakhmetov S, Zhumadilov Z, Müller DN, Wilck N, Kushugulova A, Forslund SK. Blood pressure changes correlate with short-chain fatty acid production potential shifts under a synbiotic intervention. *Cardiovasc Res*. 2020;116:1252–1253. doi: 10.1093/cvr/cvaa083
123. Duschka A, Gisevius B, Hirschberg S, Yissachar N, Stangl GI, Eilers E, Bader V, Haase S, Kaisler J, David C, et al. Propionic acid shapes the multiple sclerosis disease course by an immunomodulatory mechanism. *Cell*. 2020;180:1067–1080.e16. doi: 10.1016/j.cell.2020.02.035
124. Hager J, Bang H, Hagen M, Frech M, Trager P, Sokolova MV, Steffen U, Tascilar K, Sarter K, Schett G, et al. The role of dietary fiber in rheumatoid arthritis patients: a feasibility study. *Nutrients*. 2019;11:2392. doi: 10.3390/nu11102392
125. Hui W, Yu D, Cao Z, Zhao X. Butyrate inhibit collagen-induced arthritis via Treg/IL-10/Th17 axis. *Int Immunopharmacol*. 2019;68:226–233. doi: 10.1016/j.intimp.2019.01.018
126. Vinolo MA, Rodrigues HG, Hatanaka E, Sato FT, Sampaio SC, Curi R. Suppressive effect of short-chain fatty acids on production of proinflammatory mediators by neutrophils. *J Nutr Biochem*. 2011;22:849–855. doi: 10.1016/j.jnutbio.2010.07.009
127. Vinolo MA, Hatanaka E, Lambertucci RH, Newsholme P, Curi R. Effects of short chain fatty acids on effector mechanisms of neutrophils. *Cell Biochem Funct*. 2009;27:48–55. doi: 10.1002/cbf.1533
128. Singh N, Thangaraju M, Prasad PD, Martin PM, Lambert NA, Boettger T, Offermanns S, Ganapathy V. Blockade of dendritic cell development by bacterial fermentation products butyrate and propionate through a transporter (Slc5a8)-dependent inhibition of histone deacetylases. *J Biol Chem*. 2010;285:27601–27608. doi: 10.1074/jbc.M110.102947
129. Wang F, Liu J, Weng T, Shen K, Chen Z, Yu Y, Huang Q, Wang G, Liu Z, Jin S. The inflammation induced by lipopolysaccharide can be mitigated by short-chain fatty acid, butyrate, through upregulation of IL-10 in septic shock. *Scand J Immunol*. 2017;85:258–263. doi: 10.1111/sji.12515
130. Säemann MD, Böhmig GA, Osterreicher CH, Burtscher H, Parolini O, Diakos C, Stöckl J, Hörl WH, Zlabinger GJ. Anti-inflammatory effects of sodium butyrate on human monocytes: potent inhibition of IL-12 and up-regulation of IL-10 production. *FASEB J*. 2000;14:2380–2382. doi: 10.1096/fj.00-0359fje
131. Sun M, Wu W, Chen L, Yang W, Huang X, Ma C, Chen F, Xiao Y, Zhao Y, Ma C, et al. Microbiota-derived short-chain fatty acids promote Th1 cell IL-10 production to maintain intestinal homeostasis. *Nat Commun*. 2018;9:3555. doi: 10.1038/s41467-018-05901-2
132. Arpaia N, Campbell C, Fan X, Dikiy S, van der Veeken J, deRoos P, Liu H, Cross JR, Pfeffer K, Coffey RJ, et al. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature*. 2013;504:451–455. doi: 10.1038/nature12726
133. Chen H, Ma X, Liu Y, Ma L, Chen Z, Lin X, Si L, Ma X, Chen X. Gut microbiota interventions with *Clostridium butyricum* and norfloxacin modulate immune response in experimental autoimmune encephalomyelitis mice. *Front Immunol*. 2019;10:1662. doi: 10.3389/fimmu.2019.01662
134. Frost F, Storck LJ, Kacprowski T, Gärtner S, Rühlemann M, Bang C, Franke A, Völker U, Aghdassi AA, Steveling A, et al. A structured weight

- loss program increases gut microbiota phylogenetic diversity and reduces levels of Collinsella in obese type 2 diabetics: a pilot study. *PLoS One*. 2019;14:e0219489. doi: 10.1371/journal.pone.0219489
135. Ozkul C, Yalinay M, Karakan T. Structural changes in gut microbiome after Ramadan fasting: a pilot study. *Benef Microbes*. 2020;11:227–233. doi: 10.3920/BM2019.0039
 136. Louis S, Tappu RM, Damms-Machado A, Huson DH, Bischoff SC. Characterization of the gut microbial community of obese patients following a weight-loss intervention using whole metagenome shotgun sequencing. *PLoS One*. 2016;11:e0149564. doi: 10.1371/journal.pone.0149564
 137. Guevara-Cruz M, Flores-López AG, Aguilar-López M, Sánchez-Tapia M, Medina-Vera I, Díaz D, Tovar AR, Torres N. Improvement of lipoprotein profile and metabolic endotoxemia by a lifestyle intervention that modifies the gut microbiota in subjects with metabolic syndrome. *J Am Heart Assoc*. 2019;8:e012401. doi: 10.1161/JAHA.119.012401
 138. Maifield SV. *Nat Commun* [online ahead of print], 2021. doi: 10.1038/s41467-021-22097-0
 139. Fischbach MA. Microbiome: focus on causation and mechanism. *Cell*. 2018;174:785–790. doi: 10.1016/j.cell.2018.07.038
 140. Ratner M. Seres's pioneering microbiome drug fails mid-stage trial. *Nat Biotechnol*. 2016;34:1004–1005. doi: 10.1038/nbt1016-1004b
 141. Stappenbeck TS, Virgin HW. Accounting for reciprocal host-microbiome interactions in experimental science. *Nature*. 2016;534:191–199. doi: 10.1038/nature18285
 142. Virgin HW, Todd JA. Metagenomics and personalized medicine. *Cell*. 2011;147:44–56. doi: 10.1016/j.cell.2011.09.009
 143. Seok J, Warren HS, Cuenca AG, Mindrinos MN, Baker HV, Xu W, Richards DR, McDonald-Smith GP, Gao H, Hennessy L, et al. Inflammation and Host Response to Injury, Large Scale Collaborative Research Program. Genomic responses in mouse models poorly mimic human inflammatory diseases. *Proc Natl Acad Sci USA*. 2013;110:3507–3512. doi: 10.1073/pnas.1222878110
 144. Bartolomaeus H, Markó L, Wilck N, Luft FC, Forslund SK, Muller DN. Precarious symbiosis between host and microbiome in cardiovascular health. *Hypertension*. 2019;73:926–935. doi: 10.1161/HYPERTENSIONAHA.119.11786
 145. Vieira-Silva S, Falony G, Darzi Y, Lima-Mendez G, Garcia Yunta R, Okuda S, Vandeputte D, Valles-Colomer M, Hildebrand F, Chaffron S, et al. Species-function relationships shape ecological properties of the human gut microbiome. *Nat Microbiol*. 2016;1:16088. doi: 10.1038/nmicrobiol.2016.88
 146. Randall DW, Kieswich J, Swann J, McCafferty K, Thiemermann C, Curtis M, Hoyle L, Yaqoob MM. Batch effect exerts a bigger influence on the rat urinary metabolome and gut microbiota than uraemia: a cautionary tale. *Microbiome*. 2019;7:127. doi: 10.1186/s40168-019-0738-y
 147. Enriquez JA. Mind your mouse strain. *Nat Metab*. 2019;1:5–7. doi: 10.1038/s42255-018-0018-3
 148. Omary MB, Cohen DE, El-Omar EM, Jalan R, Low MJ, Nathanson MH, Peek RM Jr, Turner JR. Not all mice are the same: standardization of animal research data presentation. *Hepatology*. 2016;63:1752–1754. doi: 10.1002/hep.28608
 149. Bloom SM, Bijanki VN, Nava GM, Sun L, Malvin NP, Donermeyer DL, Dunne WM Jr, Allen PM, Stappenbeck TS. Commensal Bacteroides species induce colitis in host-genotype-specific fashion in a mouse model of inflammatory bowel disease. *Cell Host Microbe*. 2011;9:390–403. doi: 10.1016/j.chom.2011.04.009
 150. Rogers GB, Kozłowska J, Keeble J, Metcalfe K, Fao M, Dowd SE, Mason AJ, McGuckin MA, Bruce KD. Functional divergence in gastrointestinal microbiota in physically-separated genetically identical mice. *Sci Rep*. 2014;4:5437. doi: 10.1038/srep05437
 151. Elinav E, Strowig T, Kau AL, Henao-Mejia J, Thaiss CA, Booth CJ, Peaper DR, Bertin J, Eisenbarth SC, Gordon JI, et al. NLRP6 inflammasome regulates colonic microbial ecology and risk for colitis. *Cell*. 2011;145:745–757. doi: 10.1016/j.cell.2011.04.022
 152. Levy M, Thaiss CA, Zeevi D, Dohnalová L, Zilberman-Schapira G, Mahdi JA, David E, Savidor A, Korem T, Herzig Y, et al. Microbiota-modulated metabolites shape the intestinal microenvironment by regulating NLRP6 inflammasome signaling. *Cell*. 2015;163:1428–1443. doi: 10.1016/j.cell.2015.10.048
 153. Mamantopoulos M, Ronchi F, Van Hauwermeiren F, Vieira-Silva S, Yilmaz B, Martens L, Saeys Y, Drexler SK, Yazdi AS, Raes J, et al. Nlrp6- and Asc-dependent inflammasomes do not shape the commensal gut microbiota composition. *Immunity*. 2017;47:339–348.e4. doi: 10.1016/j.immuni.2017.07.011
 154. Chung H, Pamp SJ, Hill JA, Surana NK, Edelman SM, Troy EB, Reading NC, Villablanca EJ, Wang S, Mora JR, et al. Gut immune maturation depends on colonization with a host-specific microbiota. *Cell*. 2012;149:1578–1593. doi: 10.1016/j.cell.2012.04.037
 155. Doeing DC, Borowicz JL, Crockett ET. Gender dimorphism in differential peripheral blood leukocyte counts in mice using cardiac, tail, foot, and saphenous vein puncture methods. *BMC Clin Pathol*. 2003;3:3. doi: 10.1186/1472-6890-3-3
 156. Mestas J, Hughes CC. Of mice and not men: differences between mouse and human immunology. *J Immunol*. 2004;172:2731–2738. doi: 10.4049/jimmunol.172.5.2731
 157. Beura LK, Hamilton SE, Bi K, Schenkel JM, Odumade OA, Casey KA, Thompson EA, Fraser KA, Rosato PC, Filali-Mouhim A, et al. Normalizing the environment recapitulates adult human immune traits in laboratory mice. *Nature*. 2016;532:512–516. doi: 10.1038/nature17655
 158. Selin LK, Brehm MA, Naumov YN, Cornberg M, Kim SK, Clute SC, Welsh RM. Memory of mice and men: CD8+ T-cell cross-reactivity and heterologous immunity. *Immunol Rev*. 2006;211:164–181. doi: 10.1111/j.0105-2896.2006.00394.x
 159. Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD, Gordon JI. Obesity alters gut microbial ecology. *Proc Natl Acad Sci USA*. 2005;102:11070–11075. doi: 10.1073/pnas.0504978102
 160. Arrieta MC, Walter J, Finlay BB. Human Microbiota-associated mice: a model with challenges. *Cell Host Microbe*. 2016;19:575–578. doi: 10.1016/j.chom.2016.04.014
 161. Marques FZ, Jama HA, Tsyganov K, Gill PA, Rhys-Jones D, Muralitharan RR, Muir J, Holmes A, Mackay CR. Guidelines for transparency on gut microbiome studies in essential and experimental hypertension. *Hypertension*. 2019;74:1279–1293. doi: 10.1161/HYPERTENSIONAHA.119.13079
 162. Bach JF. The hygiene hypothesis in autoimmunity: the role of pathogens and commensals. *Nat Rev Immunol*. 2018;18:105–120. doi: 10.1038/nri.2017.111
 163. Okada H, Kuhn C, Feillet H, Bach JF. The 'hygiene hypothesis' for autoimmune and allergic diseases: an update. *Clin Exp Immunol*. 2010;160:1–9. doi: 10.1111/j.1365-2249.2010.04139.x
 164. Roslund MI, Puhakka R, Gronroos M, Nurminen N, Oikarinen S, Gazali AM, Cineke O, Kramna L, Sitter N, Vari HK, et al. Biodiversity intervention enhances immune regulation and health-associated commensal microbiota among daycare children. *Sci Adv*. 2020;6:eaba2578. doi: 10.1126/sciadv.aba2578
 165. Rosshart SP, Herz J, Vassallo BG, Hunter A, Wall MK, Badger JH, McCulloch JA, Anastasakis DG, Sarshad AA, Leonardi I, et al. Laboratory mice born to wild mice have natural microbiota and model human immune responses. *Science*. 2019;365:eaaw4361. doi: 10.1126/science.aaw4361
 166. Oldstone MB. Molecular and cellular mechanisms, pathogenesis, and treatment of insulin-dependent diabetes obtained through study of a transgenic model of molecular mimicry. *Curr Top Microbiol Immunol*. 2005;296:65–87. doi: 10.1007/3-540-30791-5_5
 167. Foxman EF, Iwasaki A. Genome-virome interactions: examining the role of common viral infections in complex disease. *Nat Rev Microbiol*. 2011;9:254–264. doi: 10.1038/nrmicro2541
 168. Reese TA, Bi K, Kambal A, Filali-Mouhim A, Beura LK, Bürger MC, Pulendran B, Sekaly RP, Jameson SC, Masopust D, et al. Sequential infection with common pathogens promotes human-like immune gene expression and altered vaccine response. *Cell Host Microbe*. 2016;19:713–719. doi: 10.1016/j.chom.2016.04.003
 169. Rosshart SP, Vassallo BG, Angeletti D, Hutchinson DS, Morgan AP, Takeda K, Hickman HD, McCulloch JA, Badger JH, Ajami NJ, et al. Wild mouse gut microbiota promotes host fitness and improves disease resistance. *Cell*. 2017;171:1015–1028.e13. doi: 10.1016/j.cell.2017.09.016
 170. Sbiński-Kind J, Kath J, Brachs S, Streitz M, von Herrath MG, Kühl AA, Schmidt-Bleek K, Mai K, Spranger J, Volk HD. Distinct housing conditions reveal a major impact of adaptive immunity on the course of obesity-induced type 2 diabetes. *Front Immunol*. 2018;9:1069. doi: 10.3389/fimmu.2018.01069
 171. Abolins S, King EC, Lazarou L, Weldon L, Hughes L, Drescher P, Raynes JG, Hafalla JCR, Viney ME, Riley EM. The comparative immunology of wild and laboratory mice, *Mus musculus domesticus*. *Nat Commun*. 2017;8:14811. doi: 10.1038/ncomms14811
 172. Jakobsson HE, Rodríguez-Piñero AM, Schütte A, Ermund A, Boysen P, Bemark M, Sommer F, Bäckhed F, Hansson GC, Johansson ME. The composition of the gut microbiota shapes the colon mucus barrier. *EMBO Rep*. 2015;16:164–177. doi: 10.15252/embr.201439263
 173. Hanage WP. Microbiology: microbiome science needs a healthy dose of scepticism. *Nature*. 2014;512:247–248. doi: 10.1038/512247a

174. Zhang C, Zhang M, Pang X, Zhao Y, Wang L, Zhao L. Structural resilience of the gut microbiota in adult mice under high-fat dietary perturbations. *ISME J*. 2012;6:1848–1857. doi: 10.1038/ismej.2012.27
175. Ericsson AC, Gagliardi J, Bouhan D, Spollen WG, Givan SA, Franklin CL. The influence of caging, bedding, and diet on the composition of the microbiota in different regions of the mouse gut. *Sci Rep*. 2018;8:4065. doi: 10.1038/s41598-018-21986-7
176. Thaïss CA, Levy M, Korem T, Dohnalová L, Shapiro H, Jaitin DA, David E, Winter DR, Gury-BenAri M, Tatirovsky E, et al. Microbiota diurnal rhythmicity programs host transcriptome oscillations. *Cell*. 2016;167:1495–1510. e12. doi: 10.1016/j.cell.2016.11.003
177. Zmora N, Zilberman-Schapira G, Suez J, Mor U, Dori-Bachash M, Bashardes S, Kotler E, Zur M, Regev-Lehavi D, Brik RB, et al. Personalized gut mucosal colonization resistance to empiric probiotics is associated with unique host and microbiome features. *Cell*. 2018;174:1388–1405.e21. doi: 10.1016/j.cell.2018.08.041
178. Belkaid Y, Hand TW. Role of the microbiota in immunity and inflammation. *Cell*. 2014;157:121–141. doi: 10.1016/j.cell.2014.03.011
179. Wang Z, Zolnik CP, Qiu Y, Usyk M, Wang T, Strickler HD, Isasi CR, Kaplan RC, Kurland IJ, Qi Q, et al. Comparison of fecal collection methods for microbiome and metabolomics studies. *Front Cell Infect Microbiol*. 2018;8:301. doi: 10.3389/fcimb.2018.00301
180. Choo JM, Leong LE, Rogers GB. Sample storage conditions significantly influence faecal microbiome profiles. *Sci Rep*. 2015;5:16350. doi: 10.1038/srep16350
181. Costea PI, Zeller G, Sunagawa S, Pelletier E, Alberti A, Levenez F, Tramontano M, Driessen M, Hercog R, Jung FE, et al. Towards standards for human fecal sample processing in metagenomic studies. *Nat Biotechnol*. 2017;35:1069–1076. doi: 10.1038/nbt.3960
182. Bartolomaeus TUP, Forslund SK. Hitchhiker's guide to microbiome studies. *Cardiovasc Res*. 2020;116:e44–e47. doi: 10.1093/cvr/cvz316
183. Rausch P, Rühlemann M, Hermes BM, Doms S, Dagan T, Dierking K, Domin H, Fraune S, von Frieling J, Hentschel U, et al. Comparative analysis of amplicon and metagenomic sequencing methods reveals key features in the evolution of animal metaorganisms. *Microbiome*. 2019;7:133. doi: 10.1186/s40168-019-0743-1
184. Jovel J, Patterson J, Wang W, Hotte N, O'Keefe S, Mitchel T, Perry T, Kao D, Mason AL, Madsen KL, et al. Characterization of the gut microbiome using 16S or shotgun metagenomics. *Front Microbiol*. 2016;7:459. doi: 10.3389/fmicb.2016.00459
185. Langille MG, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, Clemente JC, Burkepile DE, Vega Thurber RL, Knight R, et al. Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nat Biotechnol*. 2013;31:814–821. doi: 10.1038/nbt.2676
186. Michalska-Smith MJ, Allesina S. Telling ecological networks apart by their structure: a computational challenge. *PLoS Comput Biol*. 2019;15: e1007076. doi: 10.1371/journal.pcbi.1007076
187. Donaldson GP, Lee SM, Mazmanian SK. Gut biogeography of the bacterial microbiota. *Nat Rev Microbiol*. 2016;14:20–32. doi: 10.1038/nrmicro3552
188. Suntharalingam G, Perry MR, Ward S, Brett SJ, Castello-Cortes A, Brunner MD, Panoskaltsis N. Cytokine storm in a phase 1 trial of the anti-CD28 monoclonal antibody TGN1412. *N Engl J Med*. 2006;355:1018–1028. doi: 10.1056/NEJMoa063842
189. Fisher CJ Jr, Agosti JM, Opal SM, Lowry SF, Balk RA, Sadoff JC, Abraham E, Schein RM, Benjamin E. Treatment of septic shock with the tumor necrosis factor receptor:Fc fusion protein. The Soluble TNF Receptor Sepsis Study Group. *N Engl J Med*. 1996;334:1697–1702. doi: 10.1056/NEJM199606273342603