1CD-U-3 (8380/3) (844:13) Carcinoma, mixed endometrional + Serous case to highest 8441/3

Surgical Pathology

Site Indometrium C54.1

Ju 4/3/11

TISSUE DESCRIPTION:

A1 A2 A3 A4 A5 A6 A7 A8 A9 J1 K1 L1 M1 N1 O1 P1 O1 R1 S1

B1 T1 U1 C1 C2 C3 C4 C5 D1 D2 D3 D4 E1 F1 G1 H1 I1 Uterus, right ovary (2.1 x 1.8 x 1.2 cm) with 11.0 cm segment of

right fallopian tube, and left ovary (2.0 \times 1.6 \times 1.0 cm) with 9.0

cm segment of left fallopian tube together weighing 50.0 grams;

portion of omentum (30.0 x 18.0 x 3.0 cm); right and left pelvic

lymph nodes; right and left para-aortic lymph nodes above and below

the inferior mesenteric artery; right gonadal vessels (11.0 cm in

length); left gonadal vessels (12.0 cm in length); and staging

biopsies [large bowel serosa, large bowel mesentery, right diaphragm, cul de sac, bladder peritoneum, left and right colic

gutters, right and left pelvic sidewall, small bowel serosa, and

small bowel mesentery ranging in size from 0.3 cm to 1.6 cm in

greatest dimension].

DIAGNOSIS:

Uterus, endometrium, hysterectomy: FIGO grade III (of III) mixed

serous and endometrioid carcinoma is identified forming a 3.6×3.1

 \times 2.0 cm polypoid mass. The tumor invades 0.2 cm into the myometrium (total myometrial thickness, 1.2 cm). Angiolymphatic

invasion is absent. The lower uterine segment and cervix are

uninvolved.

Uterus, myometrium, hysterectomy: No diagnostic abnormalities.

Uterus, cervix, hysterectomy: No diagnostic abnormalities.



Ovary and fallopian tube, right and left, salpingooophorectomy: No diagnostic abnormalities.

Omentum, omentectomy: Negative for tumor..

Lymph nodes, right pelvic, excision: Multiple right pelvic lymph nodes (4 external iliac, 7 internal iliac, 3 common iliac, 2 obturator) are negative for tumor.

Lymph nodes, left pelvic, excision: Multiple left pelvic lymph nodes (3 external iliac, 4 internal iliac, 7 common iliac, 7 obturator) are negative for tumor.

Lymph nodes, left and right para-aortic above and below the inferior

mesenteric artery, excision: Multiple para-aortic lymph nodes above

the inferior mesenteric artery (3 left, 2 right) are negative for

tumor. Multiple para-aortic lymph nodes below the inferior

mesenteric artery (5 left, 3 right) are negative for tumor.

Gonadal vessels, left and right, excision: The left and right gonadal vessels are negative for tumor.

Soft tissue; large bowel serosa, large bowel mesentery, right

diaphragm, cul de sac, bladder peritoneum, left and right colic

gutters, right and left pelvic sidewall, small bowel serosa, and

small bowel mesentery; staging biopsies: Negative for malignancy,

all sites.

Criteria		Yes	No
Diagnosis Discrepancy			
Primary Tumor Site D	screpancy		
HIPAA Discrepancy			
Prior Malignancy Hist	ory /		1-/-
Dual/Synchronous Rt	rhafy Noted		
Cuse is (circie):	QUALIFIED // DE	SQUARTED,	
Reviewer Initials	Date Reviewed:		