fled mnt	1"
MAY	LA
W 44	# y
-	
.,,	
	υ -4/-

Surgical Pathology Report

Final

		FI	NAL	100-0-3	_ (8380 3)
Physicien(s).	(Aga	Service: Gj .comition; Platient Lype:	a mecology Site	1CD-0-3 denocuccióna, mix Serous (8441/3) - 1 : Indometrium C54.1 Raporad:	ed endometriorid + code to highest 8441/3 LW 4/3/11
- ADENI - ADENI - SEE C JTERUS, MYON - ADENI THICKN - FOCAL	METRIUM, TOTAL / OCARCINOMA, MIXI OCARCINOMA INVO OMMENT AND SYN METRIUM, TOTAL AI OCARCINOMA BY D IESB IS 24 MM	BDOMINAL HYSTEREC	ID SEROUS TYPES ERINE SEGMENT TOMY VADING TO A DEPT	H OF 8 MM WHERE MYOMETRIAL ENTIFIED	
- SQUAI - CHROI - INCIDE ITERUS, SERO	MOUS METAPLASIA NIC INFLAMMATION ENTAL CAPILLARY I	I HEMANGIOMA INAL HYSTERECTOMY	,	UUID:75FBBF31-5 TCGA-FI- 	.c30-48C2-B4A5-2DA184DC039C A2D2-01A-PR Redact
	BES, RIGHT AND LE STOPATHOLOGIC A	FT, BILATERAL SALPI BNORMALITY	NGO-OOPHORECTO	PMY []] []] [] [] [] [] [] [
WADIES DICH	T AND LEFT, BILAT STOPATHOLOGIC A	ERAL SALPINGO-OOP BNORMALITY	HORECTOMY		
		By this signals examination	irs, I attest that the above d of the alidestandor other m	diagnosis is besed upon my personal network to the arrangement to the second to the arrangement).	

Page 1 of 3

bilateral fallopian tubes and ovaries' consists of a 7.1 x 6.2 x 5.2 cm uterus that measures 3.2 x 2.4 x 2.5 cm, cervix and attached adnexa. The specimen weighs 250 grams. Opened to show a 4.0 x 5.5 x 4.3 cm polypoid mass in the interie with no obvious invasion. Shown to the surgeon. Tissue for Rest for permanents," by

Microscopic Description and Comment:

Sections of the endometrial tumor show that it is composed of mixed histology. Approximately 70% of the tumor is endometrioid type adenocarcinoma, FIGO grade 1. The remaining 30% of the tumor is high grade serous adenocarcinoma.

and

History:

year old woman. Operative procedure: Examination under anesthesia with exploratory laparotomy with total abdominal hysterectomy, and bilateral salpingo-cophorectomy.

Specimen(s) Received:
A: UTERUS, CERVIX, BILATERAL TUBES AND OVARIES

Gross Description:

The specimen is received in a formalin-filled container labeled and "cervix, uterus, bilateral fallopian tubes and ovaries." It holds a 260-gram uterus with attached cervix, and bilateral adnexa. The uterus with attached cervix measures 7.5 cm from fundus to ectocervix, 5.2 cm from cornu to cornu, and 7.0 cm from anterior to posterior. The serosal surface is pink-tan, glistening, and shiny with a single subserosal fibroid that measures 5.0 x 5.0 cm. The $2.5 \times 3.0 \times 3.5$ cm cervix has a 3.0×3.5 cm beige, glistening, and shiny ectocervix, and a 1.0 cm diameter external os. The 2.2×0.5 cm endocervical canal is pink-tan. The 3.5×3.0 cm endometrial cavity is pink-tan, replaced by a polypoid mass that measures $4.5 \times 5.0 \times 2.3$ cm. The mass appears to be invading the lower uterine segment. The mass is 2.0 cm in maximum thickness, and it is 1.4 cm from the outer serosal surface. Sectioning shows well-circumscribed fibroid, comprised of whorled and rubbery tissue, ranging from 2.0 cm to 5.0 cm with calcification. The maximum myometrial thickness is up to 2.0 cm. The 2.5 x 2.0 x 1.5 cm overy is pink-tan and appears to be normal. Cut sections show normal ovarian parenchyma. The 7.0 x 1.5 x 1.0 cm pink-tan-brown tortuous fallopian tube has a normal outer surface. Cut section shows a large luman. The $2.5 \times 1.5 \times 1.0$ cm pink-tan ovary has a normal outer surface. Cut sections show normal ovarian parenchyma. The 5.0 x 1.2 x 1.0 cm tortuous falloplan tube with fimbrial end has a normal outer surface. Cut sections show a pinpoint lumen. The sections are submitted as follows: A1 and A2 - anterior and posterior cervix; A3 and A4 - anterior lower uterine segment; A5 - posterior jower uterine segment; A6 to A9 - anterior endomyometrium with a polypoid mass (A6 and A7 - one section); A10 to A14 - posterior endomyometrium with the mass (A10 and A11 - one section; A13 and A14 - one section); A15 and A16 - fibroid; A17 and A18 - left adnexa; A19 and A20 - right adnexa. Jar 2.

SYNOPTIC REPORTING FORM FOR MALIGNANT ENDOMETRIAL TUMORS

HISTOPATHOLOGIC TYPE

The histologic diagnosis is adenocarcinoma, mixed type (endometrioid FIGO grade 1 and serous types)

TUMOR INVASION

invasive tumor is present with superficial invasion into the luminal 1/3 of the myometrium

Page 2 of 3

Page: 2 of 3	Printed from

TUMOR SIZE The lumor invades to a depth of 8 mm The myometrial thickness is 24 mm

LOWER UTERINE SEGMENT INVOLVEMENT (does not change the stage)
The lower uterine segment is involved by turnor

ENDOCERVICAL INVOLVEMENT The endocervix is not involved by tumor

LYMPHVASCULAR SPACE INVASION Lymphvascular space invasion by tumor is present but limited in scope

REGIONAL LYMPH NODES (N)
Regional lymph nodes cannol be assessed (NX)

DISTANT METASTASIS (M) Distant metastasis cannol be assessed (MX)

PRIMARY TUMOR (TNM Category/FIGO Stage)
Tumor invades less than one-half of the myometrium (T1b/IB)

STAGE GROUPING Insufficient data to assign stage (Stage X)

The pathologic stage assigned here should be regarded as provisional, and may change after integration of clinical data not provided with this specimen.

Surgical Patholog	y report is available	on-line on		
	Surgical Patholog	Surgical Pathology report is available	Surgical Pathology report is available on-line on	Surgical Pathology report is available on-line on

Page 3 of 3

. ...