Sub Topics

Short Duration
Medium Duration
Long Duration
Ruling Varieties
New Varieties
Traditional Varieties

Paddy Varieties of Tamil Nadu

Short duration varieties

Duration (Days)	Suitable seasons	Suitable varieties
90 - 120 days	Navarai, Sornavari, Kar, Kuruvai, Late Thaladi	ADT 36, IR 50, IR 64, IR 36, MDU 3, CO 47, ASD 16, ASD 17, ADT 37, TPS 2, ADT 39, ASD 18, MDU 4, PMK 2, ADT 42, MDU 5, ASD 20, ADT 43, TKM 11, ADT (RH)1, ADT (R) 45, TKM (R) 12, TRY (R) 2, PMK (R) 3, ADT (R) 47, ADT (R) 48.

Medium duration varieties

Duration (Days)	Suitable seasons	Suitable varieties
120-140 days	Early Samba, Samba, Late Samba, Thaladi/Pishanam, Late Thaladi, Late Pishanam.	IR 20, Bhavani, IR 50, CO 43, White Ponni, ADT 38, TPS 2, TKM 10, TPS 3, ASD 19, TRY 1, CO 46, CORH 2& ADT(R) 46.

Top

Long duration varieties

Duration (Days)	Suitable seasons	Suitable varieties
140-180 days	Early Samba, Samba, Late Samba, Thaladi/Pishanam, Late Pishanam.	Ponmani, PY 4(JAWAHAR), ADT(R) 44

Top

Ruling varieties in Tamil Nadu

Short duration varieties	**IR-64, CO-47, ADT-36, ADT-37, ADT-43, ADT-45, ADT-47, ADT-48, ASD-16, ASD-17, ASD-20 and MDU-5**
Medium duration varieties	**IR 20, IR 36, CO 43, CO 46, ADT 38, ADT 39, ADT 46, Bhavani, MDU 3, MDU 4, TRY 1, ASD 19, TPS 2, TPS 3 and white ponni**
Long duration varieties	**Ponmani(CR 1009), BPT 5204, and ADT 44**
Hybrids	**CORH 1, CORH 2, CORH 3, ADTRH 1**

Ruling Varieties - Short Duration

IR 64

Parentage:IR 5657-33-2-1/ IR 2061-465-1-5-3
Duration(Days):115 – 120
Average yield(Kg/ha) :6146
1000 Grain weight(g) :23.1
Grain type : Long slender
Habit : Semi dwarf
Rice Color : White

Top

CO 47

Parentage – IR 50 / CO 43
Duration (Days) – 110-115
Average yield (Kg/ha) – 5832
1000 grain weight (g) – 20.6
Grain type – Medium slender
Habit – Erect
Rice Color – White
Special features – Resistant to Blast

Top

ADT 36

Parentage – Triveni x IR 20
Duration (Days) –110
Average yield (Kg/ha) –4000
1000 grain weight (g) – 20.6
Grain type – Medium
Habit – Erect
Rice Color - White
Special features – Resistant to Blast and Brown Plant Hopper

Top

ADT 37

Parentage – BG 280-12/ PTB 33
Duration (Days) –105
Average yield (Kg/ha) – 6200
1000 grain weight (g) – 23.4
Grain type – Short bold
Habit – Semi dwarf, Semi erect
Rice color - White
Special features – Resistant to many pest and diseases
Seed dormancy – 60 days

Top


ADT 43

Parentage – IR 50/White ponni
Duration (Days) –110
Average yield (Kg/ha) – 5900
1000 grain weight (g) – 15.5
Grain type – Medium slender
Habit – Semi dwarf
Rice color - White
Special features - Resistant to Green Leaf Hopper, high tillering and fine rice

Top


ADT 45

Parentage – IR 50/ADT 37
Duration (Days) – 110
Average yield (Kg/ha) – 6100
1000 grain weight (g) – 17.5
Grain type – Medium slender
Habit – Semi dwarf, Erect
Rice color – White
Special features - Resistant to Gall midge and moderately resistant to Brown Plant Hopper



Top

ADT 47




Parentage – ADT 43/Jeeragasamba
Duration (Days) –118
Average yield (Kg/ha) – 6200
1000 grain weight (g) – 13.5
Grain type – Medium slender
Habit – Semi dwarf and erect
Rice color – White
Special features – Resistant to stem borer, White Backed Plant Hopper and sheath blight.

Top


ADT 48

Parentage – IET 11412/ IR 64
Duration (Days) – 94-99
Average yield (Kg/ha) – 4800
1000 grain weight (g) – 22.0
Grain type – Long slender
Habit – Semi dwarf and erect
Rice color – White
Special features - Resistant to Green Leaf Hopper, Gall midge and Stem borer



Top

ASD 16




Parentage – ADT 31/CO 39
Duration (Days) – 110 - 115
Average yield (Kg/ha) – 5600
1000 grain weight (g) – 24.2
Grain type – Short Bold
Habit – Semi dwarf erect
Rice color – White
Special features – Medium tillering, Resistant to Blast

Top


ASD 17

Parentage – ADT 31/RATNA/ASD 8/IR 8
Duration (Days) – 95-101
Average yield (Kg/ha) – 5422
1000 grain weight (g) – 23.8
Grain type – Short bold
Habit – Semi dwarf, slightly open
Rice color – White
Special features – Moderately resistant to Brown Plant Hopper, Susceptible to gall midge



Top

ASD 20




Parentage – IR 18348/IR25863/IR 58
Duration (Days) – 110
Average yield (Kg/ha) – 6000
1000 grain weight (g) – 22.08
Grain type – Long Slender
Habit – Erect
Rice color – White
Special features – Resistant to many pest and diseases

Top

MDU 5


Parentage – O.glaberrima x Pokkali
Duration (Days) – 95 - 100
Average yield (Kg/ha) – 5000
1000 grain weight (g) – 21.1
Grain type – Medium slender
Habit – Erect
Rice color – White
Special features – Resistant to Drought, suitable for dry and transplanted condition



Top

Ruling Varieties - Medium Duration

IR 20




Parentage – IR 262 X TKM 6
Duration (Days) – 130 - 135
Average yield (Kg/ha) – 5000
1000 grain weight (g) – 19
Grain type – Medium slender
Habit – Semi-dwarf
Rice color – White
Special features – Resistant to stem borer

Top


IR 36

Parentage – IR 1561-228/1 IR 244/O.nivara/CR 94-13
Duration (Days) – 120
Average yield (Kg/ha) – 5000
1000 grain weight (g) – 21
Grain type – Medium
Habit – Dwarf Erect
Rice color – White
Special features – Resistant to Brown Plant Hopper, Moderately resistant to stem borer and Bacterial Leaf Blight



Top

CO 43




Parentage – Dasal x IR 20
Duration (Days) – 135 - 140
Average yield (Kg/ha) – 5200
1000 grain weight (g) – 20
Grain type – Medium slender
Habit – Erect
Rice color – White
Special features – Tolerant to saline and alkaline soil

Top


CO 46

Parentage – T7/IR 20
Duration (Days) – 125
Average yield (Kg/ha) – 6000
1000 grain weight (g) – 23.5
Grain type – Long Slender
Habit – Tall erect
Rice color – White
Special features – Resistant to Brown Plant Hopper



Top

ADT 38




Parentage – IR 1529-680-3-2/ IR 4432-52-6-4/ IR 7963-30-2
Duration (Days) – 130 - 135
Average yield (Kg/ha) – 6200
1000 grain weight (g) – 21
Grain type – Long Slender
Habit – Semi dwarf, erect
Rice color – White
Special features – Resistant to many pest, Susceptible to Bacterial Leaf Blight, Seed viability period is low

Top


ADT 39

Parentage – IR 8/IR 20
Duration (Days) – 120 - 125
Average yield (Kg/ha) – 5000
1000 grain weight (g) – 18
Grain type – Medium slender
Habit – Semi dwarf
Rice color – White
Special features – Resistant to Blast and sheath rot



Top

ADT 46




Parentage – ADT38 / CO 45
Duration (Days) – 135
Average yield (Kg/ha) – 6656
1000 grain weight (g) – 23.8
Grain type – Long Slender
Habit – Erect, semi-dwarf
Rice color – White
Special features- Resistant to Stem borer and Leaf folder

Top


Bhavani

Parentage – Peta x BPI 76
Duration (Days) – 130 - 135
Average yield (Kg/ha) – 5000
1000 grain weight (g) – 21.5
Grain type – Long Slender
Habit – Medium tall
Rice color – White



Top

MDU 3




Parentage – IR 8/W 1263
Duration (Days) – 120 - 125
Average yield (Kg/ha) - 5000
1000 grain weight (g) – 23.1
Grain type – Long Slender
Habit – Semi dwarf
Rice color – White
Special features - Highly resistant to Gall Midge, resistant to Brown Plant Hopper, leaf blast, moderately resistant to White Backed Plant Hopper, Brown Spot and neck blast

Top


MDU 4

Parentage – AC 2836/Jagannath
Duration (Days) – 120 - 125
Average yield (Kg/ha) – 5900
1000 grain weight (g) – 22.9
Grain type – Long slender
Habit – Erect, tall
Rice color – White
Special features - cold tolerant



Top

TRY 1




Parentage – IR578-172-2-2/BR-1-2-B-1
Duration (Days) – 135-140
Average yield (Kg/ha) – 5255
1000 grain weight (g) – 24
Grain type – Medium
Habit – Erect
Rice color – White
Special features - Suitable for making flaked rice (Aval) and puffed rice (pori)

Top


ASD 19

Parentage – Lalnakanda / IR 30
Duration (Days) – 127 (120 - 132)
Average yield (Kg/ha) – 5800
1000 grain weight (g) – 18.39
Grain type – Short, slender
Habit – Semi-dwarf, erect
Rice color – White
Special features - Resistant to blast and Brown Plant Hopper



Top

TPS 2




Parentage – IR 26 / CO 40
Duration (Days) – 125 - 130
Average yield (Kg/ha) – 4615
1000 grain weight (g) – 23.5
Grain type – Short Bold
Habit – Semi dwarf
Rice color – White
Special features - Moderately resistant to Brown Plant Hopper and leaf folder, Resistant to rice white tip nematode,Moderately resistant to blast and bacterial leaf blight

Top

TPS 3

Parentage – RP 31-492/LMN
Duration (Days) – 135 -140
Average yield (Kg/ha) – 5600
1000 grain weight (g) – 23.2
Grain type – Short Bold
Habit – Semi dwarf/erect
Rice color – White
Special features - Resistant to blast and Leaf folder



Top

White ponni


Parentage – Taichung 65/2/Mayang Ebos-80
Duration (Days) – 135 - 140
Average yield (Kg/ha) – 4500
1000 grain weight (g) – 16.4
Grain type – Medium slender
Habit – Medium tall
Rice color – White
Special features – Fine grain quality Rice, Resistant to Rice Tungro Virus, Moderately resistant to Brown Spot and Blast

Top

Ruling Varieties - Long Duration

ADT 44

Parentage – IR56 / OR142-99
Duration (Days) – 148
Average yield (Kg/ha) – 6214
1000 grain weight (g) – 23.9
Grain type – Short Bold
Habit – Medium Tall
Rice color – White
Special features – Resistant to blast, Green leaf hopper, Stem borer, Brown spot and Leaf Folder



Top


Ponmani(CR 1009)

Parentage – Pankaj x Jagannath
Duration (Days) – 155 - 160
Average yield (Kg/ha) – 5300
1000 grain weight (g) – 23.5
Grain type – Short bold
Habit – Erect
Rice color – White
Special features - Resistant to Brown plant hopper and high yield potential.

Top


BPT 5204

Parentage – GEB-24 x T(N)1 x Mahsuri
Duration (Days) – 150
Average yield (Kg/ha) – 6000
Grain type – Medium slender
Habit – Semi dwarf
Rice color – White
Special features – Resistant to Blast, suitable for Rainfed Shallow Low Land.



Top

PY 4(Jawahar)




Parentage – IR 8 X H4
Duration (Days) – 145 - 150
Average yield (Kg/ha) – 6200
1000 grain weight (g) – 24.8
Grain type – Long bold
Habit – Semi dwarf, erect
Rice color – White
Suitable season - Samba Season
Special features –Tolerant to Bacterial leaf blight, Brown spot, Rice Tungro virus, Sheath rot, leaf folder and stem borer

Top

Ruling Varieties - Hybrids


CORH 1

Parentage – IR-62829 A x IR 10198-66-2 R
Duration (Days) – 110-115
Average yield (Kg/ha) – 6000
Grain type – Medium slender
Habit – Semi dwarf
Rice color – White
Special features – moderately resistant to Sheahth Blight, Brown Spot, Green leaf hopper, Rice Tungro virus and Brown plant hopper



Top

CORH 2




Parentage – IR 58025 A/C 20R
Duration (Days) – 125
Average yield (Kg/ha) – 6100
1000 grain weight (g) – 23.77
Grain type – Medium
Habit – Semi dwarf
Rice color – White
Special features - High yield potential

Top


CORH 3

Parentage : TNAU CMS 2A / CB 87 R
Duration (days) : 110 – 115
Season : Kar / Kuruvai / Sornawari / Navarai
Yield (Kg/ha) : 6600
Grain type - Medium slender
Rice color - White
Special features - High yield potential, Tolerant to blast and Rice tungro virus, Resistant to Green leaf hopper and tolerant to White backed plant hopper and Brown Plant Hopper



Top

ADTRH 1




Parentage – IR 58025 A/ IR 66 R
Duration (Days) – 115
Average yield (Kg/ha) – 6400
1000 grain weight (g) – 23.8
Grain type – Long slender
Habit – Semi dwarf, erect
Rice color – White milky, scented
Special features – High yield potential

Top

New varieties of Paddy


CORH 3

Year of release - 2006
Parentage - TNAU CMS 2A / CB 87 R
Duration (days) - 110 – 115
Season - Kar / Kuruvai / Sornawari / Navarai
Yield (Kg/ha) - 6600
Grain type - Medium slender
Rice color - White
Special features - High yield potential, Tolerant to blast and Rice tungro virus, Resistant to Green leaf hopper and tolerant to White backed plant hopper and Brown Plant Hopper



Top

RMD(R) 1




Year of release - 2006
Parentage - Selection from TGR 75R
Duration (days) - 100-105
Season : September – October (Rainfed direct sowing)
Yield (Kg/ha) - 4000
Grain type - Semi dwarf
Rice color - Long bold white rice
Special features - Drought tolerant. Resistant to stem borer and leaf folder, Tolerant to bacterial leaf blight, blast and sheath rot

Top


TPS(R) 4

Year of release - 2006
Parentage - TS 29 / ASD 16
Duration (days) - 95
Season : Kannipoo (Jun-Sept.) - Kumbapoo (Oct.-Feb.)
Yield (Kg/ha) - 6000
Grain type - Medium bold
Rice color - White
Special features - Field tolerance to disease blast, brown spot and sheath blight and stem borer, leaf folder, Brown plant hopper and Green leaf hopper.



Top

CO(R) 48




Year of release - 2007
Parentage - CO 43 / ASD 19
Duration (days) - 130 – 135
Season - Thaladi
Yield (Kg/ha) - 9625
Grain type - Medium Slender
Rice color - White rice
Special features - Moderately resistant to stem borer and hoppers, blast, Rice tungro virus and sheath blight

Top


CO(R) 49

Year of release - 2008
Parentage - C 20 / RNR 52147
Duration (days) - 130-135
Season - Late Samba / Thaladi
Yield (Kg/ha) - 6286
Grain type - Medium slender
Rice color - White fine rice
Special features - Moderately resistant to blast and Rice Tungro Disease. Moderately resistant to yellow stem borer and Green Leaf Hopper



Top

PMK(R) 4 or Anna(R) 4




Year of release - 2009
Parentage - Pantdhan 10 / IET 9911
Duration (days) - 100 - 105
Season - Samba (September – October)
Yield - 3.7 tonnes / ha
Grain type - Semi dwarf, erect and non–lodging
Rice color - Long slender white rice
Special features - Drought tolerant

Top


CO(R) 50

Year of release - 2010
Parentage - CO 43 / ADT 38
Duration (days) - 130 -135
Season - Late Samba / Thaladi
Yield (Kg/ha) - 6338
Grain type - Medium slender
Rice color - White fine rice
Special features - Moderately resistant to blast, sheath blight, brown spot, bacterial leaf blight and rice tungro disease



Top

TRY 3




Year of release - 2010
Parentage - ADT 43 / Jeeraga Samba
Duration (days) - 135
Season - Samba / Late Samba / Thaladi
Yield (Kg/ha) - 5833 / ha
Grain type - Medium bold grain
Rice color - white rice
Special features - Resistant to leaf folder, stem borer, brown plant hopper, blast, brown spot, sheath rot and sheath blight

Top


ADT 49

Year of release - 2011
Parentage - CR 1009/Jeeragasamba
Duration - 130 to 137 days
Season - Late samba / Thaladi
Grain type - Medium slender
Rice color - white rice
Yield - 6173 kg/ha (10.5% over BPT 5204)
Highest yield obtained - 10,250 kg/ha
Recommended districts - All districts of Tamil Nadu except Virudhunagar, Ramnad, Sivaganga and The Nilgiris
Special features - 1000 grain weight - 14.0 g
Head rice yield - 71.3%
Non-sticky cooked rice Moderately resistant to blast, sheath rot, sheath blight and resistant to RTD under artificial conditions Moderately resistant to leaf folder and brown spot under field conditions



Top

CORH 4



Year of release - 2011
Parentage - COMS 23 A / CB 174 R
Duration - 130 – 135 days
Season - Late samba / Thaladi
Grain type - Medium slender
Rice color - White rice
Yield - 7348 kg/ha (14% over 27 P11)
Highest yield obtained - 11250 kg/ha
Recommended districts - Suitable for cultivation as transplanted Hybrid rice throughout Tamil Nadu except Virudhunagar, Ramanathapuram, Sivagangai and Nilgris districts
Special features - Medium duration hybrid Intermediate amylose content with high linear elongation ratio on cooking. Resistant to blast, brown spot and moderately resistant to WBPH, GLH, sheath blight, sheath rot and RTD.

Top

Traditional varieties of Paddy

Puzhuthi Kar Nel – Dry Land Rice

- Grown in Western Ghats bounded in Erode district under rain fed and garden land conditions.
- Direct sown crop.

- Crop duration is 125-130 days and the average height of the crop is 130 cm.
- Susceptible for lodging.
- Rice are bold and red coloured.

Top

Kalurundai

- Grown in Nagapattinam district
- Late Samba (Sep 15 – Feb 14) and Navarai (Dec 15 - Mar14) are the best suited season.
- Requires clay loam soil for better growth.
- Direct sown crop with duration of 120 days.
- It grows upto a height of 120 cm.
- It is resistant to drought, pest and disease as well as salinity.
- Colour of the grain is yellow with black colour faded stripes.
- Rice is bold sized with pale yellow colour.
- Straw obtained from this rice variety is used for thatching roof.

Top

Soolai kuruvai

- Grown in coastal villages in Sempodai, Periyakuthagai, Thopputhurai at Nagai District.
- Late samba (Sep 15-Feb 14) and Kuruvai (June 1-Aug 31) are the suitable season.
- Clay loam and coastal saline soils are well suitable.
- Grown in both irrigated or rain fed conditions.
- Crop duration is 130-140 days and grows upto a height of 110-120 cm.
- The feature of this variety is motta or bold and pale dirty brown color.
- Straw used for cattle feed and also used for thatching roof

Top

Kuzhi Vedichan

- Grown in Vellapalam and Keevalur taluk of Nagapattinam district
- Samba (July 15 – Jan 14) is the apt season for cultivating Kuzhi vedichan.
- It grows well in sandy clay soil and saline soil, so that it is very well suited for coastal areas
- It is resistant to salinity, drought and pest and diseases
- It is a short duration crop and the cropping cycle is 110 days
- Size of the grain is bold in nature with pale dirty white colour.
- The specific character of this variety is induced milk secretion in lactating women
- Even when the field is flooded and the crop lodges, it has an unique feature of tillering again.

Top

Val Sivappu

- Grown in Vellapalam and Keevalur taluk of Nagapattinam district.
- Late samba (Sep 15-Feb 14)
- Sandy clay and saline soils are most favourable soil
- Saline as well as drought tolerant.
- Crop duration is 145-150 days and grows upto 160cm in height
- Red in colour and also good taste
- Posterior end of each grain has a single bristle like structure which looks like a small bird's tail

Top

Vellai Kuruvi Kar

- Grown in Vellapalam and Keevalur taluk of Nagapattinam district.
- Samba (July15-Jan 14) and late samba (Sep 15-Feb 14) are the best suited seasons.
- Crop duration is 125-135 days out of which 25-30 days are nursery period.
- In upland conditions, direct sowing can be followed.
- This variety grows upto 100-120 cm.
- Both the grains as well as straw are pale yellow to dirty white colour.
- During tillering stage, rat damage will be more because the stem is tastier.

Top

Seeraga Samba

- Grown in Vellapalam and Keevalur taluk at Nagapattinam district
- 125-130 days, Samba variety and suited method is transplanting
- The nursery period is 30-35 days. The plant grows upto a height of 80.5 cm.
- The grain size is also very small and fine. It is highly used for making briyani on the special occasion.

Top

Pichavari

- Grown in Vellapalam and Keevalur taluk of Nagapattinam district.
- Tolerant to flood and drought and Thaladi crop.
- Direct sowing and transplanting system are used.
- Crop duration is 110-115 days

Top

Suran Kuruvai

- Grown in Vellapalam and Keevalur taluk of Nagapattinam district.
- Samba and late samba period.
- Crop duration is 130-135 days and the nursery period is 30-35 days.
- Rice is dark brown and coarse in texture.

Top

Vai Kunda

- Grown in Keevalur taluk of Nagapattinam district.
- Direct sowing and transplanting system.
- 145-150 days and nursery period is 35-40 days is inclusive.
- Tolerant to flood and drought.

Top

Kichedi samba

- Grown in Keevalur taluk of Nagapattinam district.
- Samba variety with 140 days.
- Rice colour is pale white.
- Special feature of this variety is resistant to pest and diseases.

Top

Chinna samba / Sada samba

- Grown in Keevalur taluk of Nagapattinam district.
- Samba variety with 140 days in which 30 days are nursery period.

- Grain is light green in colour during panicle initiation and milky stage, whereas in maturity stage it becomes pale yellow colour.

Top

Kuthiraival Samba

- Grown in Kampalathupatti village in Trichy district.
- Samba season (July 15-Jan 14).
- Crop duration is 140-150 days.
- Resistant to lodging and also resistant to pest and disease.
- Grain colour is yellowish, whereas rice is pure white in colour.

Top

Mara nel

- Grown in Thenkanikottai taluk of Krishnagiri district.
- Crop duration is 120-125 days and Late samba and Navarai(Summer) variety.
- Transplanting method is used for cultivation.
- The rice is red in colour and coarse in texture.
- Grains of this variety have a tough seed coat. So the grains do not germinate even if there is heavy rainfall at the time of harvest.
- Resistant to storage pest because of the hard seed coat.

Top

Kallimadayaan

- Grown in Iyerpalayam village of Perambalur district.
- Six months variety.
- It has good tillering ability with a plant height ranging from 110-120cm.
- The panicle length is 30 cm.
- Yields bold grains and has the potential to give two tonnes per acre.
- The colour of the grain is white.
- It is resistant to stem borer, ear head bug and also blight disease.
- Nursery period is 35-60 days.

Top

Karuppukavuni

- Grown in Anumandhakudi village in Sivagangai district.
- 150-170 days and Suitable season for growing this variety is September – January.
- Suitable to direct sowing.
- Needs only organic manures such as green and green leaf manure.
- Profuse tillering.
- The colour of the grain is black and it is only used for sweet making along with coconut milk and it is not suitable for meals preparation.
- Tillers are moist in nature and it fetches good weight.
- It requires lesser quantity of water comparatively and suitable for black cotton soil and red soil in dry lands.

Top

Chandikaar

- Grown in Anumandhakudi village in Sivagangai district.
- Season is September – January.
- The people often cut out the stem after three months of sowing to prevent lodging.
- It is suitable for black cotton soil, red soil and saline soil.
- It is suitable for direct sowing and transplanting.
- Resistant to paddy blight disease.

Top

Varappukkodanchan

- Grown in Chatrakudi village in Ramnad district.
- Suitable for rain fed condition, drought resistant and tall variety.
- The outer skin is black in colour and the rice is red in colour.
- Farmers believe it gives more energy compared to other fine rice varieties for working people and the straw yield is very higher.
- Three months variety and it is sown normally in the middle of Puratasi(September) and harvested in the month of Tai(January).

Top

Kuliparichan

- Grown in Pallabachery village in Ramnad district.
- Variety is suitable for coastal sandy soil and drought resistant and tall variety.
- The outer skin is slightly dark. The inner grain is bold and round and the cooked rice has good shelf-life.
- Three months variety and it is sown normally in the middle of Puratasi(September) and harvested in the month of Tai(January).

Top

Chitrakar

- Grown in Thirupullani tracts of Ramnad district.
- Peasants generally name all these local cultivars in the name of mattai or norungan.
- Suitable for coastal sandy soils and drought resistant and tall variety.
- The duration of the variety is 110 days. It is sown normally in the middle of September and harvested in the month of January.

Top

Sivappu Chitraikar

- Grown in Paramakudi tracts.
- Skin is slightly black in colour and the inner skin is red in colour.
- Drought resistant and tall variety and 110 days crop.
- Sown in the month of “Aavani” and harvested in the month of “Thai”.
- Farmers mainly rely on these traditional varieties rather than other varieties.
- “Rice kanji” of sivappu chitraikar is very tastier than other red rice varieties.

Top

Murungaikar

- Grown in Ramnad areas and it is suitable for even very low rainfall periods.
- Yield of murungaikar is 1000-1300 kg per acre in normal rain fed conditions
- The crop is sown normally in the month of Aavani (August) and harvested in Markazhi (January).
- Goat manure and Urea alone are applied generally to the Murungaikar field.

Top

Nootripathu

- Grown in Ramnad district.
- The outer skin is white in colour and the inner coat is red in colour.
- It is the drought resistant and tall variety.
- The rice kanji of this variety is good in taste and sown in “Aavani” and harvested in “Thai”.

- Some of the fields nearer to vaigai river basin, clay soil is found.
- The rainfed red rice varieties grow very well in clay soils rather than black and red soils.

Top

Ariyaan

- There are 4 types of Ariyaan rice varieties grown in sandy soils of Regunathapuram areas at Ramnad district. Viz, “Vellai Ariyaan”, Karuppu Ariyaan, Sivappu Ariyaan, and Vazhai Ariyaan.
- All Ariyaan varieties have awns. It grows very tall upto 5.5 to 6.5 feet. The duration of all the Ariyaan varieties is 120 days.
- They are sown in the month of Ayppasi (October-November) and harvested in the month of Thai(January).
- This variety is suitable for coastal sandy soils and drought resistant tall variety.
- It needs at least one rainfall shower in first 3 months for higher yield.
- The harvesting is done in three stages viz. top portion cut at first, middle portion cut at thereafter and bottom portion is cut at last.
- It is affected by leaf folder and stem borer.

Top

Sadaikkar

- It is available only in Aakadavalasai village in Ramanathapuram district.
- It has medicinal properties. It is used to cure wound injuries in leg and hand by applying over the wounded portion.
- Most of the local “maattu vaidyars” (Ethno Veterinary practicers) are using this rice for its medicinal value.
- The sowing season is September and harvested in February-March.
- The crop needs at least three showers in a month and it is not affected by pest and disease generally except leaf folder.
- It is grown in sandy soils, irrigation through well is not possible because the water will stagnate at surface level.

Top

Poongkar

- Grown in Raghunathapuram village at Ramnad district.
- If the season fails in “Puratasi” (September) another traditional early duration red rice variety of Poongkar is sown in the month of “Karthigai” (October) and it is harvested in Thai (January) but the yield is lowers.
- Suitable for coastal sandy soils, drought resistant and tall variety.
- The crop can be able to grow in very less rainfall condition and it is drought tolerant than kuliparichan and varppukodanchan


Top

Kuruvai kalanchium

- Grown generally in Ramnad district.
- The duration of the variety is 110 days and it is sown normally in the month of Aavani and harvested in the month of Thai.
- The seed rate is 40-50 kg per acre for direct sowing.
- The paddy straw yield from an acre is one tonnes.
- Rice kanji of Kuruvaikalanchiam is different than “sivappu chitraikar”. The cooked rice is very good in taste even after two days.

Top

Norungan



- Grown in Ramnad district with duration of 110 days.
- Sown in rain fed condition in sandy soils in the month of Ayppasi (October) as rainfed crop and it is harvested in the month of Thai (January).
- It is affected by leaf folder and stem borer, if there is no sufficient rain water


Top

Kalurundaikaar

- Grown in Paramakudi taluk of Ramnad district.
- Drought resistant and tall variety.
- In some fields 3 types of sorghum are grown as border crop. They are red, white and black.

Top

Thanga Samba



- Grown in Kanchipuram district.
- Late samba and samba period is the best season.
- Crop duration is 160-165 days where 40 days in nursery.
- The grain is golden colour and fine in texture.
- The ear head of this variety is very long.
- This variety of rice is extremely fine and long,it is used for the preparation of special dishes like pula.


Top

Neelan samba

- Grown in Sukkan kollai village in Kanchipuram district.
- Samba season is best suitable season and duration is 175-180 days.
- The rice is red in colour and coarse in texture.
- Highly suitable for water logging condition.
- Resistant to brown plant hopper and ear head bug.
- Suitable for lactating women.

Top

Vadan Samba




- Grown in Kanchipuram district.
- Samba season is best suitable and duration is 155-160 days.
- Direct sown and transplanting is used for growing this variety.
- The rice is dull white in colour and coarse in texture.
- Drought resistant.
- Resistant to pest and disease.
- Since the grains shatter, it is important to harvest them when they are still green.

Top


Kaliyan samba

- Grown in Sukkan kollai village at Kanchipuram district.
- Samba season is best suitable season and duration is 155-160 days out of which, 40-45 days will be nursery period.
- Transplanting is used for growing this variety.
- Rice is red in colour.
- It grows very tall upto the height of 5.5 feet.
- It can be grown in water logging and low lying area.
- Highly resistant to pest and disease.



Top

Samba Mozsanam




- Grown in Sukkan kollai village at Kanchipuram district.
- Samba season is best suitable season and the duration is 160-165 days (40-45 days nursery period).
- Dry sown and transplanting methods is used for growing this variety.
- The rice is red in colour and coarse in texture.
- This variety is also called Puzudikal, Erinel and Maduvu muzangi.
- It is suitable for growing in the vicinity of lakes.
- It is said that people travelled by boats and harvested the ‘Samba Mosanam’ in the lakes.
- The variety is good for preparing Aval (Flattened rice), idly and dosa.

Top


Kadikazhuthan

- Grown in Sukkan kollai village at Kanchipuram district.
- Samba season is best suitable season and the duration is 165 days.
- Transplanting system is used for growing this variety.
- The rice is white in colour and having the fine quality.
- It is also known as ‘Garudan samba’. The appearance of the grains is like the neck of the Brahmin kite(Garudan) i.e. it has a white ring and so called this name.
- It is suitable for making a specialty dish called ‘Puttu’ of South India.
- It grows well in areas which have good drainage facilities.



Top

Pisini




- Grown in Sukkan kollai village of Chengalpet taluk at Kanchipuram district.
- Samba season is best season and the duration of 120 days.
- The rice has red in colour and having the coarse quality.
- Highly suitable for making dosa, porridge and aval.
- Highly resistant to drought.
- Since the ear heads are found with awn it is important to take extra care while removing the grains.

Top


Thuyamalli

- Grown in Sukkan kollai village at Kanchipuram district.
- Samba season is best suitable season and the duration is 135-140 days.
- Direct sown and transplanting system is used for growing this variety.
- The rice is white in colour and having the fine quality.
- During the flowering stage the ear heads look like flowers.
- Highly resistant to pest and disease.



Top

Kalar palai



- Grown in Sukkan kollai village of Chengalpet taluk at Kanchipuram district.
- Navarai season is the best season and the duration is 120 days.
- Transplanting system is used for growing this variety.
- The rice is brown in colour and having the coarse quality.
- Resistant to pest and disease.
- Suitable for kalar nilam i.e. saline soil.


Top

Kudaivazhai

- Grown in Sukkan kollai village of Chengalpet taluk at Kanchipuram district.
- Late Samba season is best season and duration is 120-125 days.
- Transplanting is used for growing this variety.
- The rice has red in colour and having the coarse quality.
- The ear heads of this variety appear like open umbrellas and hence the name ‘kudaivazhai’.
- This variety is highly suitable for making dishes such as idly and Dosa.
- This variety is highly suitable for areas like water logging.
- This is a short duration variety; hence it can even be cultivated thrice a year.

Top

Kumvalai




- Grown in Sukkan kollai village of Chengalpet taluk at Kanchipuram district.
- Samba season is best season and duration is 128-130 days.
- Direct sown and transplanting is used for growing this variety.
- The rice is brown in colour and having the coarse quality.
- Straw is particularly preferred as a roofing material.
- It is suitable for sandy soil and also water logging area.

Top


Muttakar

- Grown in Edaiyur village of Thirukazhukundram taluk at Kanchipuram district.
- Navarai season is best season and duration of 120 days in which 30 days will be the nursery period.
- Wet sown and transplanting is used for growing this variety.
- The plant grows upto a height of 4.5 feet and it is a lodging type.
- The rice has red in colour and having the coarse quality.



Top

Sempalai



- Grown in Edaiyur village of Thirukazhukundram taluk at Kanchipuram district.
- Samba season is best season and duration of 100-105 days out of which 21-25 days are nursery period.
- Direct sown and transplanting system is used for growing this variety.
- The rice is dull white in colour and coarse quality.
- It is highly suitable for making ‘Pori’ (puffed rice).

Top

Kappakar

- Grown in Thakandarapuram village of Vandavasi taluk at Thiruvannamalai district.
- Samba season is best season and duration is 150-160 days in which 40 days will be the nursery period.
- The seed rate is 50 kg per acre.
- Dry sown and transplanting system is used for growing this variety.
- The rice is brown in colour.
- This variety is highly suitable for making idly and dosa.
- It gives high yield in clayey soil.
- It is suitable for growing in water logging areas as well as drought prone areas.

Top

Perunkar

- Grown in Thakandarapuram village of Vandavasi taluk at Thiruvannamalai district.
- Duration of the crop is 120 days and the seed rate for direct sown crop is 35 kg per acre and 40 kg per acre is used for transplanting system.
- The plant grows to a height of 4.5 feet.
- Tolerant to stem borer and ear head bug.
- Suitable for preparing idly and dosa.

Top

Kalanamak

- Kalanamak is one of the finest quality scented rices of India.
- It was quite popular in Himalayan Tarai of eastern Uttar Pradesh, India.
- Tall stature causing lodging, long duration, and low yield and grain quality.
- Ten germplasm lines were resistant to panicle blast.
- Similarly several germplasm lines performed well in Usar (saline-sodic) soil of pH 9.0 to 9.5.
- These selections offer great promise for cultivation in Usar soil.
- Cooked rice is fluffy, soft, non-sticky, sweet, and easily digestible with relatively longer shelf-life.
- In local market it earns higher price than Basmati rice, which is approximately 4 to 5 times higher than non-scented rice varieties.

Top

Copyright © TNAU | All Rights Reserved