FELLSWAY PHYSIOTHERAPY
INITIAL ASSESSMENT - LOWER LIMB

Patient: Denver Lang DOB: 1947-08-14
Date: 2024-04-15 Referring: ___, MD
Physiotherapist: ___, PT, DPT

Diagnosis: Bilateral knee osteoarthritis (M17.0), right > left

SUBJECTIVE
Knee pain for years, worse over the past 2 years. Right knee 6/10 on stairs and after 15 minutes of walking, left 3/10. Morning stiffness 15 minutes. No locking or giving way. Numbness in both feet from diabetic neuropathy; unsteady on uneven ground. One fall in 2019 (clavicle fracture) and an ankle sprain in 2021. Goal: walk 45 minutes with grandchildren and lose weight.

OBJECTIVE
Gait: antalgic on right, reduced knee flexion in swing.
ROM: right knee 0-115 degrees, left 0-125 degrees.
Strength: quadriceps 4-/5 right, 4/5 left; hip abductors 4-/5 both.
Effusion: trace right.
Outcome measures:

KOOS pain	52/100
30-second chair stand	8 repetitions (below norm for age)
Timed Up and Go	12.8 seconds
Single-leg stance	right 3 s, left 5 s

Monofilament: reduced sensation both feet.

ASSESSMENT
Knee osteoarthritis with quadriceps and hip weakness and reduced balance, increasing fall risk given neuropathy. Good candidate for progressive strengthening and a graded walking program.

PLAN
8 sessions over 10 weeks: quadriceps and hip strengthening, balance training, walking program starting at 10 minutes twice daily. Home exercise program issued. Supportive shoes advised. Cleared for supervised gym training with a trainer focused on low-impact strength work.

___, PT, DPT

Synthetic record of Denver Lang written for Stellis Health. Not a real person.