___ PRIMARY CARE ASSOCIATES
TELEPHONE ENCOUNTER

Name: ___ Unit No: ___
Date: 2024-09-19
Caller: Patient Clinician: ___, MD

Reason for Call:
Lightheadedness on standing for 2 weeks; dietitian letter and home BP log received.

History:
77 year old man with type 2 diabetes, hypertension, CABG 1997, on lisinopril 10 mg and hydrochlorothiazide 25 mg. Lost 5 kg since April with diet and exercise. Home standing readings 84/56 and 86/58 with dizziness; sitting average 97/62. No chest pain, palpitations, fainting, falls, black stools or vomiting.

Assessment:
Orthostatic lightheadedness, likely from blood pressure treatment that is now more than he needs after weight loss, with lower fluid intake. Low blood sugar unlikely on metformin alone.

Plan:
1. Stop hydrochlorothiazide 25 mg. Continue lisinopril 10 mg.
2. Fluids as advised by dietitian.
3. Rise slowly; sit on the bed edge for 1 minute in the morning.
4. Recheck home BP twice daily for 2 weeks and send readings.
5. Basic metabolic panel next week given diuretic change.
6. Seek urgent care for fainting, a fall or chest pain.

Patient agreed with plan.

___, MD

Synthetic record of Denver Lang written for Stellis Health. Not a real person.