Breast pathology report

Case TCGA-BH-A0HA

FINAL DIAGNOSIS: PART 1: BREAST, RIGHT, TOTAL MASTECTOMY -. A. B. INVASIVE LARGEST DUCTAL TUMOR (MASS CARCINOMAS #1): NOTTINGHAM (3). GRADE Il (TUBULE FORMATION 3, NUCLEAR PLEOMORPHISM. 2, SECOND LARGEST TUMOR (MASS #2): NO TINGHAM GRADE II (TUBULE FORMATION 3, NUCLEAR. MITOTIC ACTIVITY 1; TOTAL SCORE 6/9). c. PLEOMORPHISM 3, MITOTIC ACTIVITY 1; TOTAL SCORE 7/9). D. SMALLEST TUMOR (MASS #3): NOTTINGHAM GRADE II (TUBULE FORMATION 3, NUCLEAR. PLEOMORPHISM 2, MITOTIC ACTIVITY 1; TOTAL SCORE 6/9). is. INVASIVE TUMORS MEASURE 1.3 CM (SLIDE 1D), 0.8 CM. AND 0.5 CM l. E. DUCTAL CARCINOMA IN-SITU (DCIS), NUCLEAR GRADE 2, SOLID PATTERN. G. F. DCIS CONSTITUTES 10% OF THE TOTAL TUMOR MASS AND IS PRESENT ADMIXED WITH AND AWAY.

FROM THE INVASIVE COMPONENT. H.

LOBULAR CARCINOMA IN-SITU (LCIS) IS ALSO IDENTIFIED. I.

NO LYMPHOVASCULAR SPACE INVASION IS IDENTIFIED. J.

RESECTION MARGINS ARE NEGATIVE FOR CARCINOMA. K. INVASIVE CARCINOMA IS 0.5 CM FROM THE NEAREST ANTERIOR MARGIN. L. DUCTAL CARCINOMA IN-SITU IS 0.5 CM FROM THE NEAREST ANTERIOR MARGIN. M.

NIPPLE IS NEGATIVE FOR TUMOR. N.

NON-NEOPLASTIC BREAST TISSUE SHOWS NON-PROLIFERATIVE FIBROCYSTIC CHANGES. O.

BIOPSY SITE CHANGES. P. INVASIVE TUMOR CELLS (MASS #1) TESTED POSITIVE FOR ESTROGEN AND PROGESTERONE. RECEPTORS AND NEGATIVE FOR HER-2 AS PER PREVIOUS PATHOLOGY REPORT. PART 2: LYMPH NODE, RIGHT, SENTINEL NODE #1, BIOPSY -. NO METASTATIC MAMMARY CARCINOMA IDENTIFIED IN ONE LYMPH NODE (0/1). PART 3: BREAST, RIGHT, SENTINEL NODE #2, BIOPSY -. NO METASTATIC MAMMARY CARCINOMA IDENTIFIED IN TWO LYMPH NODES (0/2). PART 4: LYMPH NODE, RIGHT, SENTINEL NODE #3, BIOPSY -. NO METASTATIC MAMMARY CARCINOMA IDENTIFIED IN ONE LYMPH NODE (0/1). PART 5: BREAST, LEFT, TOTAL MASTECTOMY -. A. NIPPLE WITH NO EVIDENCE OF PAGET'S DISEASE. B.

BENIGN BREAST PARENCHYMA WITH NON-PROLIFERATIVE FIBROCYSTIC CHANGES.

CASE SYNOPSIS:

SYNOPTIC - PRIMARY INVASIVE CARCINOMA OF BREAST.

LATERALITY: Right.

PROCEDURE: Simple mastectomy. Site: breast, NOS C50-9 3/0/11 her. Not specified.

SIZE OF TUMOR: Maximum dimension invasive component: 1.3 cm.

MULTICENTRICITY/MULTIFOCALITY OF INVASIVE FOCI:

TUMOR AGGREGATE SIZE: Sum of the sizes of multiple invasive tumors: 2.6 cm. TUMOR TYPE (invasive component): Ductal adenocarcinoma, NOS.

NOTTINGHAM SCORE: Nuclear grade: 2. Tubule formation: 3. Mitotic activity score: 1. Total Nottingham score: 6. Nottingham grade (1, 2, 3): 2.

ANGIOLYMPHATIC INVASION:

DERMAL LYMPHATIC INVASION:

CALCIFICATION: TUMOR TYPE, IN SITU: Solid.

SURGICAL MARGINS INVOLVED BY INVASIVE COMPONENT: Distance of invasive tumor to closest margin: 5 mm.

SURG MARGINS INVOLVED BY IN SITU COMPONENT: Distance of in situ disease to closest margin: 5 mm. PAGET'S DISEASE OF NIPPLE:

LYMPH NODES POSITIVE: o.

LYMPH NODES EXAMINED: 4.

METHOD(S) OF LYMPH NODE EXAMINATION: H/E stain.

SENTINEL NODE METASTASIS:

NON-NEOPLASTIC BREAST TISSUE: FCD. T STAGE, PATHOLOGIC: pT1c. N STAGE, PATHOLOGIC: pNO(i-). M STAGE, PATHOLOGIC: pMX.

ESTROGEN RECEPTORS: positive.

PROGESTERONE RECEPTORS: positive. HER2/NEU: zero or 1+.

Text from TCGA-Reports (Kefeli et al., 2024), MIT License. De-identified TCGA pathology report.