CLARA Cardiovascular Clinical Question Bank

100 Selected Questions from the NLM Clinical Questions Collection

Research and Development Evaluation Material

Editable reviewer copy | 6 November 2026

Paste each CLARA-generated answer beneath its question, then ask the reviewer to enter one answer relevancy score in the box provided. The response paragraphs expand normally in Word.

Answer relevancy rating instructions

Read each question alongside its corresponding CLARA-generated answer. Rate how directly and completely the answer addresses the question and whether it includes unnecessary content.

1: Does not address the question.

2: Mostly unrelated; addresses only a small part.

3: Partially addresses the question, with substantial omissions.

4: Directly addresses most of the question, with minor omissions.

5: Directly and fully addresses the question without unnecessary content.

This researcher-developed scale measures clinician-rated answer relevancy separately from automated RAGAS Answer Relevancy. Review the rubric before scoring. Record clinical inaccuracies, safety concerns, and reasons for scores separately using the question number; relevance does not establish medical correctness.

Document scope and source attribution

This document contains 100 unique cardiovascular questions selected from 481 cardiovascular-tagged records imported from the NLM Clinical Questions Collection. It is a CLARA project compilation, not an official NLM publication or a clinically validated examination.

The National Library of Medicine (NLM, 2025) describes the collection as questions gathered from healthcare providers in clinical settings between 1991 and 2003. The collection is historical and is no longer expanded with new content.

Selection and presentation

Selection emphasized adult cardiovascular diagnosis, treatment, monitoring, prevention, and medication safety, with reduced repetition. This purposive selection is not a statistical sample or a validated ranking of clinical importance.

Question text uses the source general_question field when available, otherwise the original_question field. Only display whitespace is normalized. Each item includes its NLM ID and original collection record ID. The accompanying JSONL retains the original wording and archive provenance.

Evaluation status

Reference answers and evaluation scores are not included. CLARA responses will be reviewed using the answer relevancy rubric at the beginning of this document. Reference answers must be prepared from the relevant guideline PDFs and reviewed separately. Validation remains pending until the reviewer completes the final sign-off. Historical question wording is retained for traceability.

Source citation

National Library of Medicine. (2025). Clinical Questions Collection [Data set]. Data.gov.

https://catalog.data.gov/dataset/clinical-questions-collection

Source archive references and attribution details appear after Question 100.

Hypertension: diagnosis, treatment, and monitoring

1. What is the best way to decide whether to go by home blood pressures that are normal versus office blood pressures that are high?

Source: NLM | NQ000767 | IA-0767 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

2. How many elevated blood pressures are required to make the diagnosis of hypertension?

Source: NLM | NQ003926 | IC-0926 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

3. What are the indications for a 24-hour blood pressure monitor?

Source: NLM | NQ001878 | IB-0895 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

4. Does it make any difference if you take the blood pressure again in the same arm after a few seconds of the first reading? Do you get an accurate reading the second time?

Source: NLM | NQ001995 | IB-1012 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

5. Does it make any difference whether someone has their shirt on with a rolled up sleeve when you take their blood pressure?

Source: NLM | NQ001994 | IB-1011 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

6. What antihypertensives should be used according to what comorbid conditions are present?

Source: NLM | NQ002571 | IB-1621 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

7. What two- and three-drug combinations are reasonable for hypertension?

Source: NLM | NQ002609 | IB-1659 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

8. What is the drug of choice to treat hypertension in someone with chronic renal failure?

Source: NLM | NQ002167 | IB-1186 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

9. What are good drugs to treat hypertension in the elderly?

Source: NLM | NQ002720 | IB-1775 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

10. What antihypertensives are safe to use when the patient wants to get pregnant?

Source: NLM | NQ003606 | IC-0606 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

11. What antihypertensives are safe while breast feeding?

Source: NLM | NQ001548 | IB-0562 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

12. What does the workup for secondary causes of hypertension consist of?

Source: NLM | NQ002966 | IB-2114 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

13. What are the options for ruling out renal artery stenosis in a patient with refractory hypertension?

Source: NLM | NQ002801 | IB-1857 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

14. If a patient is on three medications for hypertension which is not controlled, should a fourth medication be added or should a substitution be made?

Source: NLM | NQ003776 | IC-0776 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

15. What are effective techniques to aid medication compliance in patients with hypertension?

Source: NLM | NQ000047 | IA-0047 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

16. What is the recommended approach to an angiotensin-converting-enzyme-inhibitor-induced cough in terms of evaluation and treatment?

Source: NLM | NQ000805 | IA-0805 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

17. Do nonsteroidal anti-inflammatory drugs cause hypertension? If so, how commonly?

Source: NLM | NQ002714 | IB-1767 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

18. Are oral decongestants contraindicated in patients with hypertension?

Source: NLM | NQ002103 | IB-1122 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

19. Should a low diastolic blood pressure which causes an increase in the pulse pressure be treated?

Source: NLM | NQ003715 | IC-0715 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

20. What testing and what treatment is indicated for a patient with orthostatic hypotension which may be exacerbating coronary artery spasm?

Source: NLM | NQ003275 | IC-0275 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

21. What is the definition of malignant hypertension (criteria for diagnosis)?

Source: NLM | NQ004046 | IC-1046 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

22. How do you use labetalol to acutely lower blood pressure?

Source: NLM | NQ002537 | IB-1586 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

23. How often should patients with stable hypertension and hyperlipidemia be seen?

Source: NLM | NQ000696 | IA-0696 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

24. What was the outcome of the controversy about whether sodium in the diet aggravates hypertension?

Source: NLM | NQ001612 | IB-0627 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

25. What class of antihypertensives have the least effect on male sexual function?

Source: NLM | NQ000374 | IA-0374 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

Heart failure: assessment and treatment

26. What are New York Heart Association classes?

Source: NLM | NQ000225 | IA-0225 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

27. In a patient with congestive heart failure, how can you tell whether they have systolic or diastolic dysfunction?

Source: NLM | NQ001583 | IB-0597 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

28. What is a normal left ventricular ejection fraction?

Source: NLM | NQ002170 | IB-1189 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

29. What would cause global hypokinesis on an echocardiogram (ejection fraction 40%)? What kinds of cardiomyopathy are there?

Source: NLM | NQ002725 | IB-1780 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

30. What is the best way to determine whether the recent onset of dyspnea in a patient with chronic congestive heart failure is due to the heart failure or something else?

Source: NLM | NQ001065 | IA-1065 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

31. What drugs should patients with heart failure be on unless contraindicated, besides diuretics?

Source: NLM | NQ002726 | IB-1781 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

32. When should you use carvedilol or other beta blockers (for example, metoprolol) for congestive heart failure?

Source: NLM | NQ002536 | IB-1585 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

33. How do you treat congestive heart failure when you also have chronic renal failure?

Source: NLM | NQ001514 | IB-0528 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

34. Are angiotensin converting enzyme inhibitors contraindicated in hypertensive patients with a creatinine over a certain level?

Source: NLM | NQ001731 | IB-0747 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

35. How do you treat refractory congestive heart failure?

Source: NLM | NQ001494 | IB-0508 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

36. How do you treat diastolic dysfunction when seen on echocardiogram?

Source: NLM | NQ002883 | IB-2030 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

37. Is there any treatment that improves survival in patients with diastolic dysfunction?

Source: NLM | NQ003528 | IC-0528 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

38. When should a potassium-sparing diuretic be used instead of potassium supplementation?

Source: NLM | NQ000376 | IA-0376 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

39. In a patient with alcoholic cardiomyopathy, is it OK to drink any alcohol at all?

Source: NLM | NQ003413 | IC-0413 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

40. What causes jugular venous distension besides congestive heart failure?

Source: NLM | NQ002439 | IB-1488 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

Lipids and cardiovascular prevention

41. What is the best treatment for hypercholesterolemia in an elderly (mid-70's) patient?

Source: NLM | NQ000391 | IA-0391 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

42. At what age should screening cholesterol testing be discontinued?

Source: NLM | NQ000368 | IA-0368 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

43. In a patient status post coronary angioplasty, should elevated lipids be treated initially with diet only or should drugs be used as initial therapy?

Source: NLM | NQ001071 | IA-1071 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

44. What is the recommended management for an elderly woman with elevated cholesterol (total cholesterol approximately 290, low-density-lipoprotein (LDL) cholesterol approximately 210)?

Source: NLM | NQ000850 | IA-0850 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

45. If someone is taking lovastatin and they get sore muscles but have a normal CPK (creatine phosphokinase), do you still need to worry about myositis from lovastatin?

Source: NLM | NQ001950 | IB-0967 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

46. How high can you let the creatine phosphokinase get in someone on Mevacor (lovastatin) before you have to stop the drug?

Source: NLM | NQ002082 | IB-1101 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

47. What types of fat are good for preventing coronary disease?

Source: NLM | NQ002399 | IB-1445 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

48. Who should receive primary prophylaxis against vascular events with aspirin?

Source: NLM | NQ003529 | IC-0529 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

49. Should a young woman with a strong family history of coronary disease at a young age be screened for coronary disease?

Source: NLM | NQ003445 | IC-0445 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

50. Is it okay to use the nicotine patch in someone with known coronary artery disease?

Source: NLM | NQ001751 | IB-0767 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

Coronary disease, chest pain, and cardiac testing

51. What is the best way to determine whether chest pain is due to coronary disease?

Source: NLM | NQ000509 | IA-0509 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

52. What are the indications for immediate evaluation of a patient who gives a history of angina?

Source: NLM | NQ000553 | IA-0553 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

53. What are the possible causes of chest pain in women, and what is the best way to distinguish among the possibilities?

Source: NLM | NQ000587 | IA-0587 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

54. How do you treat an acute myocardial infarction in the emergency room? What things should be done?

Source: NLM | NQ002808 | IB-1864 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

55. Should patients who are status post myocardial infarction be placed on beta blockers, aspirin, and angiotensin-converting-enzyme inhibitors indefinitely?

Source: NLM | NQ002804 | IB-1860 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

56. How do you interpret Troponin levels?

Source: NLM | NQ002187 | IB-1206 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

57. What is the difference between ischemia, injury, and infarction on the electrocardiogram?

Source: NLM | NQ002286 | IB-1309 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

58. What is the relative accuracy and cost of various stress tests for coronary artery disease?

Source: NLM | NQ003532 | IC-0532 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

59. Why should you do a pharmacologic stress echocardiogram rather than an exercise stress echocardiogram in a patient with left bundle branch block?

Source: NLM | NQ003747 | IC-0747 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

60. When should a patient who is status post coronary bypass surgery or other procedure for coronary artery disease have followup exercise testing?

Source: NLM | NQ003521 | IC-0521 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

61. Should a patient with typical chest pain and a negative treadmill 3 years ago have further testing?

Source: NLM | NQ003019 | IC-0019 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

62. What is the best cardiac evaluation to do in a patient following a coronary artery bypass graft, who has terrible lungs and can't do a treadmill?

Source: NLM | NQ000925 | IA-0925 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

Arrhythmias and conduction disorders

63. What is the recommended management of new onset atrial fibrillation?

Source: NLM | NQ000847 | IA-0847 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

64. What drug should be used for chemical cardioversion of atrial fibrillation?

Source: NLM | NQ001956 | IB-0973 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

65. How do you differentiate ventricular tachycardia from supraventricular tachycardia with aberrancy?

Source: NLM | NQ002006 | IB-1023 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

66. How do you treat ventricular tachycardia?

Source: NLM | NQ002917 | IB-2065 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

67. What causes Torsades de Pointes and how do you treat it? Overdrive with external pacemaker?

Source: NLM | NQ001710 | IB-0726 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

68. When should you be concerned about frequent premature ventricular contractions in an adult?

Source: NLM | NQ002455 | IB-1504 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

69. What are the possible causes of syncope, and what is the best way to distinguish among the possibilities?

Source: NLM | NQ000505 | IA-0505 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

70. Should a young woman with anxiety and palpitations have a Holter monitor?

Source: NLM | NQ003689 | IC-0689 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

71. What should you do with extreme bradycardia (35 beats per minute) in an asymptomatic adult?

Source: NLM | NQ002681 | IB-1734 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

72. What are the indications for a pacemaker in a patient with intermittent third-degree heart block?

Source: NLM | NQ000612 | IA-0612 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

73. What are the side effects of amiodarone?

Source: NLM | NQ002192 | IB-1211 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

74. Could verapamil, a beta blocker, and digoxin cause enough bradycardia to produce weakness and dizziness?

Source: NLM | NQ000319 | IA-0319 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

Stroke, transient ischemic attack, and carotid disease

75. What is the recommended evaluation in a patient with transient ischemic attacks?

Source: NLM | NQ000032 | IA-0032 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

76. In a patient with transient ischemic attacks but no heart murmur, is an echocardiogram indicated?

Source: NLM | NQ000033 | IA-0033 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

77. In a patient with atrial fibrillation, what are the possible causes of a dizzy episode besides a transient ischemic attack, and what is the best way to distinguish among the possibilities?

Source: NLM | NQ000663 | IA-0663 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

78. Is Plavix (clopidogrel) effective in preventing transient ischemic attacks? What are the indications?

Source: NLM | NQ002518 | IB-1567 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

79. When does the maximal swelling and clinical deterioration occur after an intracerebral hemorrhage?

Source: NLM | NQ002477 | IB-1526 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

80. What are the indications for carotid endarterectomy?

Source: NLM | NQ003423 | IC-0423 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

81. What is the recommended evaluation and treatment of patients with a carotid bruit?

Source: NLM | NQ000089 | IA-0089 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

82. What are the ultrasound (duplex doppler) recommendations for surveillance after carotid endarterectomy?

Source: NLM | NQ003366 | IC-0366 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

Anticoagulation and thromboembolic disease

83. What are the indications for starting Coumadin in a patient with chronic atrial fibrillation?

Source: NLM | NQ000939 | IA-0939 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

84. What is the best way to decide whether a patient with atrial fibrillation at high risk of falling should be on Coumadin?

Source: NLM | NQ000845 | IA-0845 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

85. How should you anticoagulate someone in atrial fibrillation before you cardiovert them?

Source: NLM | NQ002078 | IB-1097 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

86. How long should you keep someone anticoagulated after a successful cardioversion?

Source: NLM | NQ002064 | IB-1083 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

87. How should you suspend anticoagulants for surgery? Should you use heparin, low-molecular weight heparin, or nothing and for how long?

Source: NLM | NQ002172 | IB-1191 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

88. What would cause the protime to fluctuate from time to time on the same dose of Coumadin?

Source: NLM | NQ000903 | IA-0903 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

89. How do you dose Vitamin K if you want to reverse the effects of warfarin because the prothrombin time is too long?

Source: NLM | NQ002608 | IB-1658 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

90. What is the dose of Lovenox and do you need to alter the dose of the elderly or if there is renal insufficiency?

Source: NLM | NQ003302 | IC-0302 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

91. What is the reference to that algorithm for diagnosing a pulmonary embolus and what is the strategy?

Source: NLM | NQ002242 | IB-1265 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

92. How do you treat a deep-vein thrombosis and pulmonary embolus during pregnancy and how do you monitor the patient on heparin or low-molecular-weight heparin?

Source: NLM | NQ003308 | IC-0308 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

Peripheral vascular disease and broader cardiovascular assessment

93. How can you distinguish the leg pain that comes from intermittent claudication as opposed to diabetic neuropathy?

Source: NLM | NQ000332 | IA-0332 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

94. How do you interpret different ankle/brachial blood pressure indices? What's bad, what's normal, is there a cutoff ratio for considering surgery?

Source: NLM | NQ001111 | IB-0019 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

95. What are the 5 P's of acute arterial occlusion?

Source: NLM | NQ002643 | IB-1694 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

96. Is it reasonable to get periodic ultrasounds to rule out an abdominal aortic aneurysm in someone with a family history of abdominal aortic aneurysm?

Source: NLM | NQ002297 | IB-1320 | Collection IB

CLARA response:

Clinician answer relevancy score (1-5): ☐

97. How often should a patient have an ultrasound or computerized tomography (CT) to follow a thoracic aortic aneurysm?

Source: NLM | NQ000715 | IA-0715 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

98. What is the recommended followup for a patient with an incidentally noted Type B aortic dissection on chest computerized tomogram?

Source: NLM | NQ003455 | IC-0455 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

99. Should asymptomatic patients get screening ultrasound evaluations for carotid artery stenosis, abdominal aortic aneurysms, or peripheral vascular disease?

Source: NLM | NQ003367 | IC-0367 | Collection IC

CLARA response:

Clinician answer relevancy score (1-5): ☐

100. What are the indications for an echocardiogram in a young adult with a Grade I or II murmur?

Source: NLM | NQ000191 | IA-0191 | Collection IA

CLARA response:

Clinician answer relevancy score (1-5): ☐

References and source attribution

National Library of Medicine. (2025). Clinical Questions Collection [Data set]. Data.gov.

https://catalog.data.gov/dataset/clinical-questions-collection

National Library of Medicine. (n.d.). IOWA Collection IA questions [XML data set]. Clinical Questions Collection. https://data.lhncbc.nlm.nih.gov/public/Clinical-Questions/Collection-IA-1095-Questions.zip

National Library of Medicine. (n.d.). IOWA Collection IB questions [XML data set]. Clinical Questions Collection. https://data.lhncbc.nlm.nih.gov/public/Clinical-Questions/Collection-IB-1892-Questions.zip

National Library of Medicine. (n.d.). IOWA Collection IC questions [XML data set]. Clinical Questions Collection. https://data.lhncbc.nlm.nih.gov/public/Clinical-Questions/Collection-IC-1062-Questions.zip

Attribution and traceability

All selected questions are drawn from Collections IA, IB, and IC. Collection ID was included in the candidate import but contributes no items to this shortlist. Item-level source labels identify the corresponding XML records. The 2025 date above refers to the catalog-reported dataset update, not the year each clinical question was asked.

The Data.gov catalog identifies the dataset license as the Open Data Commons Open Database License (ODbL). Source attribution is retained; the selection and organization are specific to the CLARA project. Catalog and download links were checked on 2 November 2026.

The citations identify the origins of the questions. They are not clinical evidence supporting an answer. References for eventual evaluation answers must point to the actual guideline evidence used.

Validation statement - complete after review

By signing below, I confirm that I have reviewed the submitted dataset, selected questions, and accompanying CLARA-generated answers for their stated research purpose. My validation is limited to the materials reviewed and is subject to my recorded comments and required corrections.

Validated by: Johnrey T. Pepito

Team Manager

New Marketlink Pharmaceutical Corporation

Health & Wellness Segment

Signature: _____ Date: _____