Greater Western Hospital

Discharge Summary

55 Federation Street, Western Plains NSW 2830 | Tel: (02) 6883 1000 | Fax: (02) 6883 1001
Western NSW Local Health District | NSW Health

Patient Name:	**Ryan Christopher Stewart**	Medical Record No.:	**545161**
Date of Birth:	25/06/1977 (Age 48)	Medicare No.:	5877 95436 2
Address:	309 King Cl, Leichhardt NSW 2040	Next of Kin:	Stefan Stewart (Father) 0456 764 302
GP:	Dr Zara Jones, Eastside General Practice	Allergies:	**Aspirin (asthma)**

ADMISSION DETAILS

Date Admitted:	04/04/2025 via Emergency Dept	Date Discharged:	09/04/2025
Ward:	6C (Respiratory)	Length of Stay:	5 days
Admitting Clinician:	Dr O. Lin (Advanced Trainee, Respiratory Medicine)	Consultant:	A/Prof S. Goldstein (Staff Specialist)
Discharge Destination:	Home (independent)	Discharge Mode:	Patient transport

PRESENTING COMPLAINT

Mr Stewart presented with sudden-onset right-sided pleuritic chest pain and progressive dyspnoea over 24 hours. Recent 14-hour flight a three-day history prior. SpO2 91% on arrival.

PRINCIPAL DIAGNOSIS

Acute bilateral pulmonary embolism (ICD-10-AM: I26.99)

ADDITIONAL DIAGNOSES

1. Recent long-haul travel
2. Hormonal contraception (where applicable)

HOSPITAL COURSE

Admitted with bilateral PE confirmed on CTPA. Anticoagulation with apixaban initiated. Respiratory support continued. Discharged Day 4 with 6-month course of apixaban. Outpatient thrombophilia screen arranged.

MEDICATIONS AT DISCHARGE

Medication	Dose / Route / Frequency	Notes
Apixaban	10mg PO BD	NEW - first 7 days
Apixaban	5mg PO BD	NEW - from Day 8 for 6 months
Paracetamol	1g PO QID PRN	for pain

KEY INVESTIGATIONS

Bloods on admission (04/04/2025): D-dimer 8,400 ng/mL (H), Troponin I 24 ng/L (H), BNP 180 pg/mL (H), Lactate 1.6 mmol/L.

Imaging: CTPA: bilateral pulmonary emboli involving right main pulmonary artery and left lower lobe segmental branches. Mild RV strain. Lower limb Doppler: no DVT.

FOLLOW-UP PLAN

SYNTHETIC TRAINING DOCUMENT - NOT FOR CLINICAL USE
Generated for MEDISCAN Document AI training | Batch ID: AUS-TRN-0048
All patient data is fictional. Any resemblance to real individuals is coincidental.

1. Continue apixaban for 6 months minimum.
2. GP review at 1 week.
3. Respiratory clinic at 4 weeks.
4. Outpatient thrombophilia screen.

ALERTS AND PRECAUTIONS

ALLERGY: Aspirin (asthma)

Anticoagulation: bleeding risk discussed. Avoid contact sports.

PREPARED BY

Dr Olivia Lin, Advanced Trainee, Respiratory Medicine

Reviewed and approved by: A/Prof Steven Goldstein, Staff Specialist, Respiratory Medicine

Date: 09/04/2025

Dr O. Lin (Advanced Trainee) A/Prof S. Goldstein (Staff Specialist)

SYNTHETIC TRAINING DOCUMENT - NOT FOR CLINICAL USE
Generated for MEDISCAN Document AI training | Batch ID: AUS-TRN-0048
All patient data is fictional. Any resemblance to real individuals is coincidental.