

## A Review on the Antioxidative and Prooxidative Properties of Luteolin

Haixia Xu<sup>1</sup>, Becky S. Linn<sup>2</sup>, Yingmei Zhang<sup>1,2</sup>, and Jun Ren<sup>1,2</sup>

<sup>1</sup>Department of Cardiology and Shanghai Institute of Cardiovascular Diseases, and Department of Cardiology, Zhongshan Hospital, Fudan University, Shanghai 210032, China; <sup>2</sup>School of Pharmacy, University of Wyoming College of Health Sciences, Laramie, WY 82071, USA

Correspondence: ren.jun@zs-hospital.sh.cn (J.R.)

*Xu H et al. Reactive Oxygen Species* 7(21):136–147, 2019; ©2019 Cell Med Press

<http://dx.doi.org/10.20455/ros.2019.833>

(Received: December 11, 2018; Revised: February 15, 2019; Accepted: February 16, 2019)

**ABSTRACT** | Luteolin is a natural flavonoid compound mainly present as glycosides in vegetables, fruits, and herbs including carrot, cabbage, artichoke, tea, celery, and apple. Evidence has revealed that luteolin possesses profound biological properties such as antimutagenic, antitumorigenic, antioxidative, immunomodulatory, anticarcinogenic, antibacterial, antiapoptotic, and anti-inflammatory capacities. In particular, luteolin helps preserve the oxidation and antioxidation balance, the disruption of which plays a crucial role in the pathophysiologic processes of various diseases including cardiovascular disorders, cancers, and neurodegenerative diseases. This mini-review briefly summarizes the established beneficial effect of luteolin with regard to the maintenance of balance between pro- and anti-oxidation. We will provide an overview of luteolin and its therapeutic application in cardiovascular diseases, cancers, and neurodegenerative diseases.

**KEYWORDS** | Antioxidative activity; Luteolin; Oxidative stress; Reactive oxygen species; Redox balance

**ABBREVIATIONS** | ATP, adenosine triphosphate; ER, endoplasmic reticulum; HCC, hepatocellular carcinoma; IMM, inner mitochondrial membrane; LDL, low-density lipoprotein; MAPK, mitogen-activated protein kinase; MPTP, mitochondrial permeability transition pore; NOX, NADPH oxidase; OMM, outer membrane of mitochondria; ROS, reactive oxygen species; SOD, superoxide dismutase

### CONTENTS

1. Introduction
2. Basic Chemical Structure and Function of Luteolin
3. Antioxidative Properties of Luteolin
4. Prooxidative Properties of Luteolin
5. Application of Luteolin in Cardiovascular Diseases
6. Application of Luteolin in Cancer
7. Application of Luteolin in Neurodegenerative Diseases
8. Application of Luteolin in Other Diseases
9. Summary and Conclusion

## 1. INTRODUCTION

Reactive oxygen species (ROS) are commonly defined as a class of small and highly reactive chemical species resulting from incomplete one-electron reduction of oxygen (e.g., superoxide, hydroxyl radicals, and peroxides such as hydrogen peroxide) [1, 2]. In physiological settings, ROS are produced through both non-enzymatic and enzyme-catalyzed reactions. Electrons are transmitted through respiratory chain reactions to complete oxidative phosphorylation in mitochondria [3, 4]. An electrochemical gradient is established across the inner mitochondrial membrane (IMM) in this process for the production of adenosine triphosphate (ATP). Cellular ROS are commonly generated through mitochondrial respiratory chain in most tissues. Steady-state ROS are maintained at nontoxic levels through various antioxidant and repair enzymes [2, 5, 6]. The delicate balance between oxidative stress and antioxidant defenses appears to play an essential role in cell survival, proliferation, and cell death [7, 8]. In pathological conditions, accumulation of ROS can result in the oxidation and damage of proteins, DNA, and lipids, resulting in pathological sequelae such as oxidative stress, inflammation, apoptosis, and autophagy [2]. On the other hand, cells generate non-enzymatic or enzymatic anti-oxidative defensive factors including superoxide dismutase (SOD), catalase, glutathione, thioredoxin, catalase, and peroxidases to scavenge ROS and combat oxidative stress [2, 9]. This oxidative/anti-oxidative balance plays a crucial role in the physiological processes of various organisms and disruption of this balance can lead to a number of diseases including cardiovascular disorders, cancers, and neurodegenerative diseases [2, 4].


Flavonoids are a class of hydroxy derivatives found ubiquitously in various kinds of vegetables, fruits, and herbs. It has been well known that fruit and vegetable intake is closely related to risk reduction of various forms of cancers and cardiovascular diseases [10]. While this protection seems to be of primary benefit from  $\beta$ -carotene and ascorbate, flavonoid ingredients should not be discounted. Luteolin (3',4',5,7-tetrahydroxyflavone) is a natural, soluble flavone found in high concentrations in carrot, cabbage, artichoke, tea, celery, and apple (**Figure 1**). Accumulating evidence has depicted an important role for luteolin in an array of biological processes such as cardiovascular regulation, antitu-

tagenic, antitumorigenic, antioxidative, immunologic, anticarcinogenic, antibacterial, and antiapoptotic activities, as well as anti-inflammatory properties [11]. Epidemiological research has validated that luteolin intake may play a vital role in the reduced prevalence of certain chronic diseases, while prospective clinical studies are still warranted to further consolidate its clinical benefit [12, 13]. This mini-review summarizes recent research discoveries on luteolin concerning primarily its regulatory role in the balance of antioxidative and prooxidative functions and its protective effects in cardiovascular disorders and other chronic diseases.

## 2. BASIC CHEMICAL STRUCTURE AND FUNCTION OF LUTEOLIN

Luteolin is a natural flavonoid compound found mostly in the form of glycosides, among various medicinal plants (**Figure 1**). These plants have higher content of whole leaves of blue orchid, honeysuckle, and perilla [12]. It was originally isolated from leaves, stems, and branches of the genus *Resedaceae* and *Reseda odorata* L. Levels of luteolin are generally lower than some other flavonols such as quercetin and kaempferol [11]. However, the levels may be much higher in some plants including sage, artichokes, parsley, wild carrots, thyme, and peanut hulls [11]. The median consumption of luteolin is approximately 0.01–0.2 mg/day through daily diet [14]. Luteolin is generally absorbed through intestinal mucosa after oral administration. For example, with a quantity of 14.3 mg/kg luteolin taken orally, a maximal plasma concentration ( $C_{max}$ ) may be achieved within  $1.02 \pm 0.22$  h, with a  $C_{max}$  of  $1.97 \pm 0.15$   $\mu$ g/ml, and a half-life of  $4.94 \pm 1.2$  h [15].

Since flavonoids possess large  $\pi$  bonds and super delocalization, their oxygen atoms are highly coordinated to form coordination complexes with ease. Moreover, the efficacy of diverse flavonoids varies according to its glycosidic moieties, hydroxylation pattern, conjugation between the aromatic rings, and methoxy groups [16]. For instance, the ability to scavenge peroxy radicals of luteolin significantly exceeds kaempferol, which attributes to the 3',4'-catechol structure in the B-ring. Consequently, lipid peroxidation inhibition of luteolin is greatly enhanced, although luteolin and kaempferol share identical hydroxyl configurations [17–19].



**FIGURE 1.** The chemical structure of luteolin and its role in maintaining the redox balance between anti-oxidative and prooxidative properties in protecting against various pathological processes. The major dietary sources of luteolin are also shown in the scheme. See text for detailed discussion on how luteolin protects against disease processes via its redox properties.

Luteolin is one of the most effective flavonoids, given its well-characterized structure and pharmacological properties [11]. Nonetheless, the pharmacokinetics of luteolin is less clear. It is reported that luteolin is hydrolyzed and absorbed in small intestines, where it undergoes hydrolysis to yield luteolin aglycone, and methyl- and sulfate conjugates [20]. Additionally, the absorption of luteolin may vary considerably among foods, due to the heterogeneity of sugars, dosages, vehicles of administration, gender, and microbial population of the colon [16]. Furthermore, bioavailability requires a quantitative estimate for its storage, after biotransformation to glucuronide. Although *in vivo* studies have revealed the beneficial effect of luteolin following intraperitoneal injection or oral intake, molecules of luteolin glycosides are cleaved during the process of uptake and are unable to be detected in the plasma. To this end, results from most *in vitro* studies should be considered with special caution.

### 3. ANTIOXIDATIVE PROPERTIES OF LUTEOLIN

Cumulative evidence has suggested that ROS play a crucial role in the pathogenesis of various diseases including cardiovascular disorders, neurodegenerative diseases, and cancer. The production and aggregation of ROS cause DNA damage, protein cross-linking, and lipid peroxidation, which contribute to mutagenesis, carcinogenesis, cellular aging [21], destabilization of cell membranes [17], and oxidation of low-density lipoprotein (LDL) [22–24]. These oxidative stress conditions are accompanied with the loss of mitochondrial transmembrane potential and the opening of mitochondrial permeability transition pores (MPTP) in the IMM. The increased opening of MPTPs leads to reduced ATP synthesis [25]. Simultaneously, the loss of mitochondrial transmembrane potential may result in the disruption of the outer membrane of mitochondria (OMM). Consequently,

the external leakage of apoptogenic factors, including cytochrome c and caspases, into the cytosol results in cell death [26].

In the past few decades, cellular oxidative and antioxidative balance has become an important factor for the maintenance of physiological functions of humans and other organisms. Novel cytoprotective treatment strategies geared towards the oxidative/antioxidative balance may effectively improve the pathological processes. Screening for drugs or compounds, such as luteolin, with a known oxidative/antioxidative balancing property or a multi-target activity would aid in providing an effective regimen for various diseases including cardiovascular diseases, neurodegenerative disorders, and cancer.

A plethora of experimental evidence has revealed the antioxidative properties for luteolin. As shown in **Figure 2**, luteolin was demonstrated to suppress oxidative damage and lipid peroxidation, and loss of antioxidant enzymes including catalase and SOD. It was also shown that luteolin attenuated lipopolysaccharide-induced pulmonary injury via decreasing NF- $\kappa$ B and mitogen-activated protein kinase (MAPK) expression [27]. Furthermore, luteolin possessed the highest DNA-protective effect against  $H_2O_2$  [28]. Radio-protective studies revealed that luteolin possessed a remarkable protective property against gamma-ray-induced DNA damage [29]. In addition, an investigation in human melanoma HMB-2 cells demonstrated DNA protective potential of luteolin in scavenging free radicals and reducing damage caused by  $H_2O_2$  in a dose-dependent manner. It was shown that luteolin decreased the incidence of chromosomal aberrations more effectively than quercetin [30]. Research has suggested that ROS generation drastically increased when human bronchial epithelial cells (BEAS-2B) were exposed to Cr(VI) after a short time. Luteolin could reduce lipid peroxidation, NADPH oxidase (NOX) activation, and glutathione depletion in a dose-dependent manner (**Figure 2**). Further studies also indicated that luteolin protected against carcinogenesis through scavenging ROS and impacting multiple cell signaling pathways related to ROS [31].


It has been found that luteolin can form complexes with metal ions such as magnesium, iron, and zinc, and has potent free radical scavenging capacity. The ability to chelate metal ions of luteolin attributes to the possession of an *o*-di-OH catechol group in the B ring which is strongly associated with its antioxi-

dative property [11]. Furthermore, luteolin may protect or enhance intracellular antioxidative enzymes, such as catalase, SOD, glutathione reductase, and glutathione *S*-transferase [32]. It was found that mice treated with luteolin exhibited significantly lowered incidence of rectal crypt lesions and upregulated levels of catalase, SOD, glutathione reductase, and glutathione *S*-transferase during the azoxymethane-induced colon carcinogenesis [33]. Moreover, luteolin may interfere with the oxidative production of some cellular components, probably through the inhibition of corresponding enzymes [34].

#### 4. PROOXIDATIVE PROPERTIES OF LUTEOLIN

Luteolin primarily plays a preventative role in human nutrition and diseases due to its antioxidative properties through mechanism summarized in **Figure 2**. Meanwhile, luteolin may possess a prooxidative activity in protecting against some cancers. In normal conditions, luteolin as an antioxidant has weak prooxidative abilities and cannot undergo redox cycling. Some phytochemicals have displayed potent prooxidative action, leading to an oxidative stress environment that cannot be tolerated by cancer cells, ultimately resulting in subsequent cell death [35]. Similarly, the aforementioned study has found that luteolin also exerts its prooxidative effect on cholangiocarcinoma cells, lung cancer cells, glioblastoma cells, and melanoma cells [36–39]. This may mean that it exhibits selective cytotoxicity to cancer cells with no effect on normal cells in preclinical testing. Previous studies have revealed that luteolin plays its proapoptotic role by disrupting the oxidative/antioxidative balance, which leads to severe ROS accumulation, the loss of mitochondrial membrane potential, and endoplasmic reticulum (ER) stress, activating mitochondrial apoptotic pathways and causing cell death (**Figure 2**) [26].

The mechanism of the prooxidative capacity of luteolin is thought to be related to the number of hydroxyl groups [16, 40]. Hanasaki and colleagues once reported that a series of mono- and di-hydroxy flavonoids had no detectable prooxidative capacity, while the compounds with multiple hydroxyl groups, especially in the B-ring, could significantly increase the production of hydroxyl radicals in the Fenton system [41]. Other studies revealed that luteolin may



**FIGURE 2.** Possible mechanisms contributing to luteolin-mediated regulation of the balance of antioxidative and prooxidative functions and its protective role in cardiovascular and other chronic diseases. SERCA denotes sarco(endo)plasmic reticulum  $\text{Ca}^{2+}$ -ATPase. See text for detailed discussion.

potentiate the increase of ROS generation and the activation of caspase-9 and caspase-3, accompanied by the release of cytochrome c and the decreased Bcl-2 and Bcl-XL proteins in KKU-100 cholangiocarcinoma cells [26] (also see **Figure 2**). More in-depth work is needed to further define the mechanism behind the prooxidative properties of luteolin.

In addition to its participation in redox regulation, luteolin may exert an array of additional biological functions, including cardiovascular regulation, anti-inflammatory activity, antibacterial effect, antitumor action, and immune regulation. It is known that luteolin may suppress macrophage activation and inhibit the inflammation in endothelial cells [42, 43]. An independent study revealed that luteolin inhibited niacin-induced flush [44, 45], and increased insulin sensitivity in endothelial cells [42]. Luteolin also retarded peroxidation of low-density lipoprotein (LDL) in diet-induced obesity [46], as well as prevented against high fat-diet-induced cognitive deficits in mice [47].

## 5. APPLICATION OF LUTEOLIN IN CARDIOVASCULAR DISEASES

Cardiovascular diseases are known to be a widespread public health problem and a leading cause of morbidity and mortality. Considerable evidence re-

vealed that oxidative stress plays a vital role in the pathogenesis and development of coronary heart disease, heart failure, and cardiac arrhythmias. The importance of antioxidants in the diet in the intervention and prevention of cardiovascular diseases has aroused considerable interest over the last few years. Several studies have shown that luteolin may protect against cardiovascular disorders via acting as an antioxidative agent.

A recent epidemiological study, involving individuals aged 65–99 years old with a 10-year follow-up, suggests that luteolin may have important influence on the reduced risk of acute myocardial infarction [48]. Another study, involving a cohort of 2,748 Finnish men and 2,385 Finnish women, showed that subjects with very low consumptions of flavonoids displayed higher risks of coronary heart disease. It is thought that luteolin might afford the protective effect through protection against oxidation of LDL [49]. In another population study of over 9000 Finnish residents with a 28-year follow-up, it was reported that the risk of cardiovascular disease risk could be related to apple intake, but not intake of quercetin which is the main flavonol component of apples [50]. A United States and European population cohort study also reported that the intake of flavonoids is strongly related to a lower incidence of mortality of coronary heart disease [51]. Another double-blind, randomized, placebo-controlled study showed that



the polyphenolic compound, pycnogenol, could improve the endothelial function in coronary heart disease [52]. Based on the aforementioned studies, it appears that nutritional intake of luteolin would contribute to the protective effect against cardiovascular disease due to its capacity of inhibiting lipid peroxidation, chelating redox-active metals, and attenuating other processes involving ROS.

Data from preclinical studies were consistent with the above-mentioned epidemiological studies. Xia and colleagues reported that luteolin may attenuate the development of atherosclerosis by protecting endothelial cells from TNF- $\alpha$ -induced oxidative stress and inflammatory response through NF- $\kappa$ B and MAPK signaling. This research revealed that luteolin decreased the production of ROS and the expression of NOX4 and p22<sup>phox</sup> expression. While the exact mechanism needs further investigation, luteolin treatment led to down-regulation of NOX4 and decreased BCL-2, ICAM-1, and VCAM-1 [53]. In another study, luteolin was found to possess potent protective actions in cardiomyocytes subjected to simulated ischemia/reperfusion injury. Several mechanisms, including inhibiting oxidative stress, improving cardiomyocyte contractile function, and activating PI3K/Akt pathways, were speculated to participate in luteolin-afforded cardiac benefits [54]. It was also found that luteolin exerted protection against the prooxidant pyrogallol-induced superoxide accumulation and subsequent apoptosis in cardiac capillary endothelial H5V and human umbilical vein endothelial cells. It was suggested that luteolin might be capable of up-regulating Bcl-2 expression and suppressing the expression of caspase-3 and caspase-8, and reducing apoptosis via MAPK signaling pathway [55].

Based on the above discussion, luteolin appears to play a cardioprotective role. The protective effect may stem from the maintenance of oxidative/antioxidative balance and the interaction with mitochondrial function by suppressing the phosphorylation of JNK and p38 MAPK and up-regulating the expression of MnSOD [56]. In addition to the antioxidative action, luteolin also promotes sarco(endo)plasmic reticulum Ca<sup>2+</sup>-ATPase (SERCA) activity and improves systolic or diastolic function during cardiac ischemia/reperfusion injury and heart failure in rats [57, 58]. Several mechanisms have been suggested for luteolin-induced cardiac responses, including phosphorylation of protein kinase B

(Akt), phospholamban, and p38 MAPK [59, 60]. Thus, luteolin could be considered as a candidate for therapeutics against cardiovascular diseases.

## 6. APPLICATION OF LUTEOLIN IN CANCER

Investigation has been ongoing for several decades regarding the biological functions of ROS and their potential involvement in the development and disease progression of cancer [61]. Cancer cells are known to be in a more active metabolic condition and undergo increased oxidative stress, which is possibly related to the uncontrolled cell proliferation and metabolic regulation dysfunction. Since the overproduction of ROS disrupts the balance between oxidation and antioxidative defense systems, it is possible that the apoptotic induction of cancer cells is one of the direct approaches to attenuating the development and progressing of cancers [62].

The prooxidative properties of flavonoids have been observed in various cell types [63, 64]. A recent study demonstrated that both luteolin and kaempferol could induce apoptosis in cells isolated from hepatocellular carcinoma (HCC), but not normal cells. The apoptosis of HCC cells was subsequent to the reduction of cytochrome c and caspase-3 activation. Luteolin was suggested as a potential candidate for the complementary treatment of HCC [36]. Another research also showed that luteolin inhibited cancer cell growth and induced cell death in HepG2 hepatocarcinoma cells via the release of ROS and AMPK-NF- $\kappa$ B signaling [65].

Another study also found that luteolin played a prooxidative and proapoptotic role in KKU-100 cholangiocarcinoma cells [66], which could be attributed to its ability to promote intracellular ROS production and subsequently activating the mitochondria-mediated apoptotic cell death. These effects were related to the suppression of Nrf2, heme oxygenase-1 (HO-1), and gamma-glutamylcysteine ligases, mitochondrial depolarization, release of cytochrome c, and the decrease of Bcl-2 and Bcl-xl [26]. In addition, the mechanism of ROS accumulation induced by luteolin has also been investigated in lung cancer cells, which seem to be involved in the decrease of SOD activity [37]. A proteomic study in human hepatoma Huh-7 cells revealed that several proteins were associated with the anti-cancer effects of luteolin. Interestingly, these proteins included

peroxiredoxin 6 (PRDX6) and prohibitin, which are implicated in ROS metabolism and apoptosis [66].

A study examined the sensitivity to the ionizing radiation in non-small cell lung cancer cells and revealed a radio-sensitizing role for luteolin due to its induction of apoptotic cell death through p38-ROS-caspase cascade, an effect independent of activation of p53 and phosphatase and tensin homolog (PTEN) [67]. Furthermore, luteolin was found to promote ROS accumulation, which led to ER stress in human melanoma cells through up-regulated expression of ER stress-associated factors, including protein kinase RNA-like ER kinase, activating transcription factor (ATF) 6, phosphorylation of eukaryotic translation initiation factor 2 $\alpha$ , C/EBP homologous protein (CHOP), and cleaved caspase-12 [39]. Work in cultured glioblastoma cells and in vivo suggested similar activities of luteolin through an ERK-eIF2 $\alpha$ -ATF4-CHOP-caspase 12-mediated mechanism [38]. These studies indicated that luteolin promoted ROS accumulation to cause apoptosis, ER stress, and mitochondrial dysfunction in various carcinoma cells. To this end, luteolin may have the potential to become an antitumor drug through its prooxidative characteristics.

## 7. APPLICATION OF LUTEOLIN IN NEURODEGENERATIVE DISEASES

Under normal metabolic conditions, there are levels of ROS generated in the mitochondria which are rapidly scavenged by enzymatic and non-enzymatic antioxidants. The maintenance of oxidative and antioxidative balance prevails in various tissues and cells [68]. Nonetheless, the capacity of scavenging ROS in the brain is relatively weak, making the tissue more sensitive to oxidant stress [69, 70]. In certain pathological conditions, the cerebral tissue is susceptible to attack of increased ROS accumulation, which inflicts the peroxidation of membrane polyunsaturated fatty acids and proteins and also increases membrane permeability [71]. Therefore, the increased oxidative stress may be one of the major pathogenic causes in the development and progression of neurodegenerative diseases, including cerebral ischemia/reperfusion injury, Alzheimer's disease, and Parkinson's disease, as well as aging [72, 73]. The antioxidative property of luteolin provides a basis for its use in treating neurodegenerative diseases.

Some studies have shown a great capacity of luteolin to protect neurons against oxidative stress. Luteolin could exert protective effects on oxidative attack-related events including increased formation of endogenous free radicals, loss of mitochondrial membrane potential, and mitochondrial dysfunction [74]. Under chronic cerebral hypo-perfusion conditions, luteolin administration also protected rats against the cognitive dysfunction and increased the neuron survival via decreasing intracellular ROS accumulation and up-regulation of Nrf2-ARE pathway [75]. Another research also suggested that luteolin protected against the high-fat diet-induced cognitive dysfunction in mice possibly through its antioxidative function [47]. Additionally, luteolin treatment improved the cognitive function and neuronal damage in both cortex and hippocampus due to its antioxidative activity in a streptozotocin-induced diabetes model [76].

Alzheimer's disease is a complicated disease associated with multiple genetic and environmental factors. In addition to the formation of senile plaques and neurofibrillary tangles due to amyloid- $\beta$  and tau proteins, a growing body of evidence suggests that oxidative stress and mitochondrial dysfunction are possibly involved in Alzheimer's disease. Zhou and colleagues reported that luteolin efficiently attenuated zinc-induced tau hyperphosphorylation and decreased oxidative stress in Alzheimer's disease. Luteolin was considered as a nontoxic and highly bioavailable dietary flavonoid for protecting against chronic neurological damage induced by zinc toxicity [77, 78].

## 8. APPLICATION OF LUTEOLIN IN OTHER DISEASES

A number of studies have indicated that ultraviolet light induces generation of harmful ROS burst, contributing to the development of human skin carcinoma. Considering luteolin's capacity to penetrate into deep skin layers, it may be useful in counteracting the oxidative stress as well as affording anti-inflammatory effects in skin carcinoma [34]. Another potential protective effect of luteolin was shown in children with autism, where a formulation containing luteolin significantly improved children's attention and behavior [79]. It should be mentioned that luteolin is marketed and advertised as a dietary supple-

ment in humans. Moreover, luteolin is a potential antiangiogenic treatment for retinal neovascularization, which is associated with its antioxidative capacity to decrease ROS and subsequently proangiogenic function of vascular endothelial growth factor [80].

## 9. SUMMARY AND CONCLUSION

Luteolin exerts various beneficial biological actions through its antioxidative properties. These effects are likely attributable to a maintenance of oxidative and antioxidative balance. In order to ensure the safety and efficacy of luteolin, more research is warranted prior to the full validation of luteolin as an antioxidative agent. The beneficial effects of luteolin, either alone or in combination with another medicinal compound, should be directed to treat diseases associated with profound oxidative stress. It is likely that luteolin may possess a therapeutic value for multiple pathological conditions.

## ACKNOWLEDGMENTS

This work was not supported by any grants. The authors declare no conflicts of interest.

## REFERENCES

1. Pelicano H, Carney D, Huang P. ROS stress in cancer cells and therapeutic implications. *Drug Resist Updat* 2004; 7(2):97–110.
2. Siasos G, Tsigkou V, Kosmopoulos M, Theodosiadis D, Simantiris S, Tagkou NM, et al. Mitochondria and cardiovascular diseases-from pathophysiology to treatment. *Ann Transl Med* 2018; 6(12):256. doi: 10.21037/atm.2018.06.21.
3. Thomas JJ, Ren J. Obstructive sleep apnoea and cardiovascular complications: perception versus knowledge. *Clin Exp Pharmacol Physiol* 2012; 39(12):995–1003. doi: 10.1111/1440-1681.12024.
4. Ren J, Pulakat L, Whaley-Connell A, Sowers JR. Mitochondrial biogenesis in the metabolic syndrome and cardiovascular disease. *J Mol Med (Berl)* 2010; 88(10):993–1001. doi: 10.1007/s00109-010-0663-9.
5. Zhou H, Wang S, Hu S, Chen Y, Ren J. ER-mitochondria microdomains in cardiac ischemia-reperfusion injury: a fresh perspective. *Front Physiol* 2018; 9:755. doi: 10.3389/fphys.2018.00755.
6. Zhang Y, Whaley-Connell AT, Sowers JR, Ren J. Autophagy as an emerging target in cardiorenal metabolic disease: From pathophysiology to management. *Pharmacol Ther* 2018; 191:1–22. doi: 10.1016/j.pharmthera.2018.06.004.
7. Wientjes FB, Segal AW. NADPH oxidase and the respiratory burst. *Semin Cell Biol* 1995; 6(6):357–65.
8. Vignais PV. The superoxide-generating NADPH oxidase: structural aspects and activation mechanism. *Cell Mol Life Sci* 2002; 59(9):1428–59.
9. Scherz-Shouval R, Elazar Z. Regulation of autophagy by ROS: physiology and pathology. *Trends Biol Sci* 2011; 36(1):30–8. doi: 10.1016/j.tibs.2010.07.007.
10. Havsteen BH. The biochemistry and medical significance of the flavonoids. *Pharmacol Ther* 2002; 96(2–3):67–202.
11. Seelinger G, Merfort I, Schempp CM. Anti-oxidant, anti-inflammatory and anti-allergic activities of luteolin. *Planta Med* 2008; 74(14):1667–77. doi: 10.1055/s-0028-1088314.
12. López-Lázaro M. Distribution and biological activities of the flavonoid luteolin. *Mini Rev Med Chem* 2009; 9(1):31–59.
13. Gates MA, Tworoger SS, Hecht JL, De Vivo I, Rosner B, Hankinson SE. A prospective study of dietary flavonoid intake and incidence of epithelial ovarian cancer. *Int J Cancer* 2007; 121(10):2225–32.
14. Wang L, Lee IM, Zhang SM, Blumberg JB, Buring JE, Sesso HD. Dietary intake of selected flavonols, flavones, and flavonoid-rich foods and risk of cancer in middle-aged and older women. *Am J Clin Nutr* 2009; 89(3):905–12. doi: 10.3945/ajcn.2008.26913.
15. Weng CJ, Yen GC. Flavonoids, a ubiquitous dietary phenolic subclass, exert extensive in vitro anti-invasive and in vivo anti-metastatic activities. *Cancer Metastasis Rev* 2012; 31(1–2):323–51. doi: 10.1007/s10555-012-9347-y.
16. Heim KE, Tagliaferro AR, Bobilya DJ. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *J Nutr*



- Biochem* 2002; 13(10):572–84.
17. Mora A, Payá M, Ríos JL, Alcaraz MJ. Structure-activity relationships of polymethoxyflavones and other flavonoids as inhibitors of non-enzymic lipid peroxidation. *Biochem Pharmacol* 1990; 40(4):793–7.
18. Dugas AJ, Castañeda-Acosta J, Bonin GC, Price KL, Fischer NH, Winston GW. Evaluation of the total peroxyl radical-scavenging capacity of flavonoids: structure-activity relationships. *J Nat Prod* 2000; 63(3):327–31.
19. Ratty AK, Das NP. Effects of flavonoids on nonenzymatic lipid peroxidation: structure-activity relationship. *Biochem Med Metab Biol* 1988; 39(1):69–79.
20. Day AJ, DuPont MS, Ridley S, Rhodes M, Rhodes MJ, Morgan MR, et al. Deglycosylation of flavonoid and isoflavonoid glycosides by human small intestine and liver beta-glucosidase activity. *FEBS Lett* 1998; 436(1):71–5.
21. Sastre J, Pallardó FV, Viña J. Mitochondrial oxidative stress plays a key role in aging and apoptosis. *IUBMB Life* 2000; 49(5):427–35.
22. Khan MA, Baseer A. Increased malondialdehyde levels in coronary heart disease. *J Pak Med Assoc* 2000; 50(8):261–4.
23. Kawanishi S, Hiraku Y, Oikawa S. *Mutat Res* 2001; 488(1):65–76.
24. Takabe W, Niki E, Uchida K, Yamada S, Satoh K, Noguchi N. Oxidative stress promotes the development of transformation: involvement of a potent mutagenic lipid peroxidation product, acrolein. *Carcinogenesis* 2001; 22(6):935–41.
25. Kim JS, He L, Lemasters JJ. Mitochondrial permeability transition: a common pathway to necrosis and apoptosis. *Biochem Biophys Res Commun* 2003; 304(3):463–70.
26. Kittiratphatthana N, Kukongviriyapan V, Prawan A, Senggunprai L. Luteolin induces cholangiocarcinoma cell apoptosis through the mitochondrial-dependent pathway mediated by reactive oxygen species. *J Pharm Pharmacol* 2016; 68(9):1184–92. doi: 10.1111/jphp.12586.
27. Kuo MY, Liao MF, Chen FL, Li YC, Yang ML, Lin RH, et al. Luteolin attenuates the pulmonary inflammatory response involves abilities of antioxidation and inhibition of MAPK and NFκB pathways in mice with endotoxin-induced acute lung injury. *Food Chem Toxicol* 2011; 49(10):2660–6. doi: 10.1016/j.fct.2011.07.012.
28. Romanová D, Vachálková A, Cipák L, Ovesná Z, Rauko P. Study of antioxidant effect of apigenin, luteolin and quercetin by DNA protective method. *Neoplasma* 2001; 48(2):104–7.
29. Shimoi K, Masuda S, Furugori M, Esaki S, Kinae N. Radioprotective effect of antioxidative flavonoids in gamma-ray irradiated mice. *Carcinogenesis* 1994; 15(11):2669–72.
30. Horváthová K, Chalupa I, Sebová L, Tóthová D, Vachálková A. Protective effect of quercetin and luteolin in human melanoma HMB-2 cells. *Mutat Res* 2005; 565(2):105–12.
31. Pratheeshkumar P, Son YO, Divya SP, Roy RV, Hitron JA, Wang L, et al. Luteolin inhibits Cr(VI)-induced malignant cell transformation of human lung epithelial cells by targeting ROS mediated multiple cell signaling pathways. *Toxicol Appl Pharmacol* 2014; 281(2):230–41. doi: 10.1016/j.taap.2014.10.008.
32. Leung HW, Kuo CL, Yang WH, Lin CH, Lee HZ. Antioxidant enzymes activity involvement in luteolin-induced human lung squamous carcinoma CH27 cell apoptosis. *Eur J Pharmacol* 2006; 534(1–3):12–8.
33. Ashokkumar P, Sudhandiran G. Protective role of luteolin on the status of lipid peroxidation and antioxidant defense against azoxymethane-induced experimental colon carcinogenesis. *Biomed Pharmacother* 2008; 62(9):590–7. doi: 10.1016/j.biopha.2008.06.031.
34. Yan M, Liu Z, Yang H, Li C, Chen H, Liu Y, et al. Luteolin decreases the UVA-induced autophagy of human skin fibroblasts by scavenging ROS. *Mol Med Rep* 2016; 14(3):1986–92. doi: 10.3892/mmr.2016.5517.
35. Wang J, Yi J. Cancer cell killing via ROS: to increase or decrease, that is the question. *Cancer Biol Ther* 2008; 7(12):1875–84.
36. Seydi E, Salimi A, Rasekh HR, Mohsenifar Z, Pourahmad J. Selective cytotoxicity of luteolin and kaempferol on cancerous hepatocytes obtained from rat model of hepatocellular carcinoma: involvement of ROS-mediated mitochondrial targeting. *Nutr Cancer* 2018; 70(4):594–604. doi: 10.1080/01635581.2018.1460679.
37. Ju W, Wang X, Shi H, Chen W, Belinsky SA, Lin Y. A critical role of luteolin-induced reactive oxygen species in blockage of tumor necrosis factor-activated nuclear factor-kappaB

- pathway and sensitization of apoptosis in lung cancer cells. *Mol Pharmacol* 2007; 71(5):1381–8.
38. Wang Q, Wang H, Jia Y, Pan H, Ding H. Luteolin induces apoptosis by ROS/ER stress and mitochondrial dysfunction in glioblastoma. *Cancer Chemother Pharmacol* 2017; 79(5):1031–41. doi: 10.1007/s00280-017-3299-4.
  39. Kim JK, Kang KA, Ryu YS, Piao MJ, Han X, Oh MC, et al. Induction of endoplasmic reticulum stress via reactive oxygen species mediated by luteolin in melanoma cells. *Anticancer Res* 2016; 36(5):2281–9.
  40. Cao G, Sofic E, Prior RL. Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships. *Free Radic Biol Med* 1997; 22(5):749–60.
  41. Hanasaki Y, Ogawa S, Fukui S. The correlation between active oxygens scavenging and antioxidative effects of flavonoids. *Free Radic Biol Med* 1994; 16(6):845–50.
  42. Deqiu Z, Kang L, Jiali Y, Baolin L, Gaolin L. Luteolin inhibits inflammatory response and improves insulin sensitivity in the endothelium. *Biochimie* 2011; 93(3):506–12. doi: 10.1016/j.biochi.2010.11.002.
  43. Ando C, Takahashi N, Hirai S, Nishimura K, Lin S, Uemura T, et al. Luteolin, a food-derived flavonoid, suppresses adipocyte-dependent activation of macrophages by inhibiting JNK activation. *FEBS Lett* 2009; 583(22):3649–54. doi: 10.1016/j.febslet.2009.10.045.
  44. Kalogeromitros D, Makris M, Chliva C, Aggelides X, Kempuraj D, Theoharides TC. A quercetin containing supplement reduces niacin-induced flush in humans. *Int J Immunopathol Pharmacol* 2008; 21(3):509–14.
  45. Papaliodis D, Boucher W, Kempuraj D, Theoharides TC. The flavonoid luteolin inhibits niacin-induced flush. *Brit J Pharmacol* 2008; 153(7):1382–7. doi: 10.1038/sj.bjp.0707668.
  46. Xu N, Zhang L, Dong J, Zhang X, Chen YG, Bao B, et al. Low-dose diet supplement of a natural flavonoid, luteolin, ameliorates diet-induced obesity and insulin resistance in mice. *Mol Nutr Food Res* 2014; 58(6):1258–68. doi: 10.1002/mnfr.201300830.
  47. Liu Y, Fu X, Lan N, Li S, Zhang J, Wang S, et al. Luteolin protects against high fat diet-induced cognitive deficits in obesity mice. *Behav Brain Res* 2014; 267:178–88. doi: 10.1016/j.bbr.2014.02.040.
  48. Marniemi J, Alanen E, Impivaara O, Seppänen R, Hakala P, Rajala T, et al. Dietary and serum vitamins and minerals as predictors of myocardial infarction and stroke in elderly subjects. *Nutr Metab Cardiovasc Dis* 2005; 15(3):188–97.
  49. Knekt P, Jarvinen R, Reunanen A, Maatela J. Flavonoid intake and coronary mortality in Finland: a cohort study. *BMJ* 1996; 312(7029):478–81.
  50. Knekt P, Kumpulainen J, R Jr, Rissanen H, Heliövaara M, Reunanen A, et al. Flavonoid intake and risk of chronic diseases. *Am J Clin Nutr* 2002; 76(3):560–8.
  51. Peterson JJ, Dwyer JT, Jacques PF, McCullough ML. Associations between flavonoids and cardiovascular disease incidence or mortality in European and US populations. *Nutr Rev* 2012; 70(9):491–508. doi: 10.1111/j.1753-4887.2012.00508.x.
  52. Enseleit F, Sudano I, Périat D, Winnik S, Wolfrum M, Flammer AJ, et al. Effects of pycnogenol on endothelial function in patients with stable coronary artery disease: a double-blind, randomized, placebo-controlled, cross-over study. *Eur Heart J* 2012; 33(13):1589–97. doi: 10.1093/eurheartj/ehr482.
  53. Xia F, Wang C, Jin Y, Liu Q, Meng Q, Liu K, et al. Luteolin protects HUVECs from TNF- $\alpha$ -induced oxidative stress and inflammation via its effects on the Nox4/ROS-NF- $\kappa$ B and MAPK pathways. *J Atheroscler Thromb* 2014; 21(8):768–83.
  54. Zhang RQ, Li DY, Xu TD, Zhu SS, Pan HJ, Fang F, et al. Antioxidative effect of luteolin pretreatment on simulated ischemia/reperfusion injury in cardiomyocyte and perfused rat heart. *Chin J Integr Med* 2017; 23(7):518–27.
  55. He D, Ma X, Chen Y, Cai Y, Ru X, Bruce IC, et al. Luteolin inhibits pyrogallol-induced apoptosis through the extracellular signal-regulated kinase signaling pathway. *The FEBS J* 2012; 279(10):1834–43. doi: 10.1111/j.1742-4658.2012.08558.x.
  56. Xu T, Li D, Jiang D. Targeting cell signaling and apoptotic pathways by luteolin: cardioprotective role in rat cardiomyocytes

- following ischemia/reperfusion. *Nutrients* 2012; 4(12):2008–19. doi: 10.3390/nu4122008.
57. Zhu S, Xu T, Luo Y, Zhang Y, Xuan H, Ma Y, et al. Luteolin enhances sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase activity through p38 MAPK signaling thus improving rat cardiac function after ischemia/reperfusion. *Cell Physiol Biol* 2017; 41(3):999–1010. doi: 10.1159/000460837.
  58. Fang F, Li D, Pan H, Chen D, Qi L, Zhang R, et al. Luteolin inhibits apoptosis and improves cardiomyocyte contractile function through the PI3K/Akt pathway in simulated ischemia/reperfusion. *Pharmacology* 2011; 88(3–4):149–58. doi: 10.1159/000330068.
  59. Hu W, Xu T, Wu P, Pan D, Chen J, Chen J, et al. Luteolin improves cardiac dysfunction in heart failure rats by regulating sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase 2a. *Sci Rep* 2017; 7:41017. doi: 10.1038/srep41017.
  60. Wu X, Xu T, Li D, Zhu S, Chen Q, Hu W, et al. ERK/PP1a/PLB/SERCA2a and JNK pathways are involved in luteolin-mediated protection of rat hearts and cardiomyocytes following ischemia/reperfusion. *PloS One* 2013; 8(12):e82957. doi: 10.1371/journal.pone.0082957.
  61. Sabharwal SS, Schumacker PT. Mitochondrial ROS in cancer: initiators, amplifiers or an Achilles' heel? *Nat Rev Cancer* 2014; 14(11):709–21. doi: 10.1038/nrc3803.
  62. Idelchik M, Begley U, Begley TJ, Melendez JA. Mitochondrial ROS control of cancer. *Semin Cancer Biol* 2017; 47:57–66. doi: 10.1016/j.semcancer.2017.04.005.
  63. Skibola CF, Smith MT. Potential health impacts of excessive flavonoid intake. *Free Radic Biol Med* 2000; 29(3–4):375–83.
  64. Matsuo M, Sasaki N, Saga K, Kaneko T. Cytotoxicity of flavonoids toward cultured normal human cells. *Biol Pharm Bull* 2005; 28(2):253–9.
  65. Hwang JT, Park OJ, Lee YK, Sung MJ, Hur HJ, Kim MS, et al. Anti-tumor effect of luteolin is accompanied by AMP-activated protein kinase and nuclear factor-kappaB modulation in HepG2 hepatocarcinoma cells. *Int J Mol Med* 2011; 28(1):25–31. doi: 10.3892/ijmm.2011.667.
  66. Yoo DR, Jang YH, Jeon YK, Kim JY, Jeon W, Choi YJ, et al. Proteomic identification of anti-cancer proteins in luteolin-treated human hepatoma Huh-7 cells. *Cancer Lett* 2009; 282(1):48–54. doi: 10.1016/j.canlet.2009.02.051.
  67. Cho HJ, Ahn KC, Choi JY, Hwang SG, Kim WJ, Um HD, et al. Luteolin acts as a radiosensitizer in non-small cell lung cancer cells by enhancing apoptotic cell death through activation of a p38/ROS/caspase cascade. *Int J Oncol* 2015; 46(3):1149–58. doi: 10.3892/ijo.2015.2831.
  68. Kehrer JP. Free radicals as mediators of tissue injury and disease. *Crit Rev Toxicol* 1993; 23(1):21–48.
  69. Kim HC, Jhoo WK, Kim WK, Suh JH, Shin EJ, Kato K, et al. An immunocytochemical study of mitochondrial manganese-superoxide dismutase in the rat hippocampus after kainate administration. *Neurosci Lett* 2000; 281(1):65–8.
  70. Lewén A, Matz P, Chan PH. Free radical pathways in CNS injury. *J Neurotrauma* 2000; 17(10):871–90.
  71. López E, Arce C, Oset-Gasque MJ, Cañadas S, González MP. Cadmium induces reactive oxygen species generation and lipid peroxidation in cortical neurons in culture. *Free Radic Biol Med* 2006; 40(6):940–51.
  72. Casetta I, Govoni V, Granieri E. Oxidative stress, antioxidants and neurodegenerative diseases. *Curr Pharm Des* 2005; 11(16):2033–52.
  73. Li H, Repa JJ, Valasek MA, Beltroy EP, Turley SD, German DC, et al. Molecular, anatomical, and biochemical events associated with neurodegeneration in mice with Niemann-Pick type C disease. *J Neuropathol Exp Neurol* 2005; 64(4):323–33.
  74. Zhao G, Yao-Yue C, Qin GW, Guo LH. Luteolin from *Purple Perilla* mitigates ROS insult particularly in primary neurons. *Neurobiol Aging* 2012; 33(1):176–86. doi: 10.1016/j.neurobiolaging.2010.02.013.
  75. Xu J, Wang H, Ding K, Zhang L, Wang C, Li T, et al. Luteolin provides neuroprotection in models of traumatic brain injury via the Nrf2-ARE pathway. *Free Radic Biol Med* 2014; 71:186–95. doi: 10.1016/j.freeradbiomed.2014.03.009.
  76. Liu Y, Tian X, Gou L, Sun L, Ling X, Yin X. Luteolin attenuates diabetes-associated cognitive decline in rats. *Brain Res Bull* 2013; 94:23–9. doi: 10.1016/j.brainresbull.2013.02.001.
  77. Zhou F, Chen S, Xiong J, Li Y, Qu L. Luteolin reduces zinc-induced tau phosphorylation at

- Ser262/356 in an ROS-dependent manner in SH-SY5Y cells. *Biol Trace Elem Res* 2012; 149(2):273–9. doi: 10.1007/s12011-012-9411-z.
78. Kwon Y. Luteolin as a potential preventive and therapeutic candidate for Alzheimer's disease. *Exp Gerontol* 2017; 95:39–43. doi: 10.1016/j.exger.2017.05.014.
79. Theoharides TC, Asadi S, Panagiotidou S. A case series of a luteolin formulation (NeuroProtek®) in children with autism spectrum disorders. *Int J Immunopathol Pharmacol* 2012; 25(2):317–23.
80. Park SW, Cho CS, Jun HO, Ryu NH, Kim JH, Yu YS, et al. Anti-angiogenic effect of luteolin on retinal neovascularization via blockade of reactive oxygen species production. *Invest Ophthalmol Vis Sci* 2012; 53(12):7718–26. doi: 10.1167/iovs.11-8790.