

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 December 2002 (12.12.2002)

PCT

(10) International Publication Number
WO 02/099090 A1

- (51) International Patent Classification⁷: **C12N 9/00**
- (21) International Application Number: PCT/AU02/00739
- (22) International Filing Date: 7 June 2002 (07.06.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
PR 5527 7 June 2001 (07.06.2001) AU
- (71) Applicant (for all designated States except US): **JOHNSON & JOHNSON RESEARCH PTY LTD** [AU/AU]; Level 4, 1 Central Avenue, Eveleigh, New South Wales 1430 (AU).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **SUN, Lun-Quan** [AU/AU]; 8 Banksia Street, EASTWOOD, New South Wales 2122 (AU). **WANG, Li** [AU/AU]; 8 Banksia Street, EASTWOOD, New South Wales 2122 (AU). **TURNER, Rachel, Jane** [AU/AU]; 54 Bellevue Street, GLEBE, New South Wales 2037 (AU). **SARAVOLAC, Edward, George** [CA/AU]; 30 Allen Street, Leichhardt, New South Wales 2040 (AU). **DASS, Crispin, Rajnish** [FJ/AU]; 35/1-9 Terrace Road, DULWHICH HILL, New South Wales 2203 (AU).
- (74) Agent: **BLAKE DAWSON WALDRON PATENT SERVICES**; Level 39, 101 Collins Street, Melbourne, Victoria 3000 (AU).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: BCL-2 DNAZYMES

(57) **Abstract:** The present invention provides DNazymes which specifically cleaves mRNA transcribed from a member of the bcl-2 gene family selected from the group consisting of bcl-2, bcl-xl, bcl-w, bfl-1, brag-1, Mcl-1 and A1. The DNazymes comprise (a) a catalytic domain that has the nucleotide sequence GGCTAGCTACAACGA (SEQ ID NO.1) and cleaves mRNA at any purine: pyrimidine cleavage site at which it is directed, (b) a binding domain contiguous with the 5' end of the catalytic domain, and (c) another binding domain contiguous with the 3' end of the catalytic domain. The binding domains are complementary to, and therefore hybridise with, the two regions immediately flanking the purine residue of the cleavage site within the bcl-2 gene family mRNA, at which DNzyme-catalysed cleavage is desired. Each binding domain is at least six nucleotides in length, and both binding domains have a combined total length of at least 14 nucleotides.



WO 02/099090 A1

bcl-2 DNAZYMES

FIELD OF THE INVENTION

5 The present invention relates to DNAzymes targeted to bcl-2 gene family members and their use in cancer therapy. This invention further relates to use of these DNAzymes to treat and/or inhibit onset of human cancers. The DNAzymes accomplish this end by cleaving mRNA transcribed from members of the bcl-2 gene family thereby provoking apoptosis of cancer cells directly and/or increasing the
10 sensitivity of cancer cells to chemotherapeutics.

BACKGROUND OF THE INVENTION

Apoptosis and Bcl-2 gene family

15 Apoptosis is a complex process resulting in the regulated destruction of a cell, which plays a major role in normal development, cellular response to injury and carcinogenesis(Ellis et al., 1991). It has been suggested that an apoptotic component either contributes to, or accounts for, many human disease pathologies including cancer, viral infection and some neurological disorders (Ashkenazi and Dixit, 1998;
20 Vocero-Akbani et al., 1999; Yakovlev et al., 1997).

 The Bcl-2 family of proteins are among the most studied molecules in the apoptotic pathway. Bcl-2 gene was first identified in B-cell lymphomas where the causal genetic lesion has been characterised as a chromosomal translocation (t (14:18)) which places the Bcl-2 gene under the control of the immunoglobulin promoter. The
25 resulting overexpression of Bcl-2 retards the normal course of apoptotic cell death that otherwise maintains B-cell homeostasis, resulting in B-cell accumulation and follicular lymphoma (Adams and Cory, 1998). This observation showed that cancers do not strictly arise from unrestrained cell proliferation, but could also be due to insufficient apoptotic turnover. In addition to follicular lymphomas, Bcl-2 levels are elevated in a
30 broad range of other human cancers, indicating that this molecule may have a role in raising the apoptotic threshold in a broad spectrum of cancerous disorders.

 The Bcl-2 gene family has at least 16 members involved in the apoptosis pathway. Some genes in this family are apoptosis inducers, including, bax, bak, bcl-Xs, bad, bid, bik and hrk, and others, such as bcl-2, bcl-XL, bcl-w, bfl-1, brag-1,
35 Mcl-1 and A1 are apoptosis suppressors (Reed, 1998). Bcl-2 family members have been suggested to act through many different mechanisms, including pore formation

in the outer mitochondrial membrane, through which cytochrome *c* (Cyt *c*) and other intermembrane proteins can escape; and heterodimerization between pro- and anti-apoptotic family members (Reed, 2000).

It has been suggested that a decrease in Bcl-2 levels or the inhibition of Bcl-2 activity might provoke apoptosis or at least sensitise cells to apoptotic death. In the absence of a clearly defined biochemical mechanism of action or activity for this family of cell-death regulatory proteins (for which conventional inhibitors could therefore be developed), gene therapy and antisense approaches have become a reasonable alternative. For example, an 18-mer all-phosphorothioate Bcl-2 antisense oligodeoxynucleotide (ODN), G-3139 that targets the first six codons of the human Bcl-2 open reading frame, has shown very promising results in both preclinical and clinical studies (Jansen et al., 1998; Waters et al., 2000). This antisense molecule binds to the Bcl-2 mRNA blocking translation of the mRNA into Bcl-2 protein and targeting the message for RNase H-mediated degradation. The resultant decrease in bcl-2 levels in the treated cells alters the balance between pro-apoptotic and anti-apoptotic family members in favour of pro-apoptotic members resulting in apoptosis.

Using a similar strategy, antisense oligonucleotides to another member of the bcl-2 gene family bcl-xL has also been shown to be active in down-regulation of the bcl-xL expression, leading to an increased chemosensitivity in a range of cancer cells (Zangemeister-Wittke et al., 2000).

Catalytic DNA (DNAzyme)

In human gene therapy, antisense nucleic acid technology has been one of the major tools of choice to inactivate genes whose expression causes disease and is thus undesirable. The anti-sense approach employs a nucleic acid molecule that is complementary to, and thereby hybridizes with, a mRNA molecule encoding an undesirable gene. Such hybridization leads to the inhibition of gene expression.

Anti-sense technology suffers from certain drawbacks. Anti-sense hybridization results in the formation of a DNA/target mRNA heteroduplex. This heteroduplex serves as a substrate for RNase H-mediated degradation of the target mRNA component. Here, the DNA anti-sense molecule serves in a passive manner, in that it merely facilitates the required cleavage by endogenous RNase H enzyme. This dependence on RNase H confers limitations on the design of anti-sense molecules regarding their chemistry and ability to form stable heteroduplexes with their target mRNA's. Anti-sense DNA molecules also suffer from problems associated with non-specific activity and, at higher concentrations, even toxicity.

As an alternative to anti-sense molecules, catalytic nucleic acid molecules have shown promise as therapeutic agents for suppressing gene expression, and are widely discussed in the literature (Haseloff and Gerlach 1988; Breaker 1994; Koizumi et al 1993; Kashani-Sabet et al 1992; Raillard et al 1996; and Carmi et al 1998) Thus, unlike a
5 conventional anti-sense molecule, a catalytic nucleic acid molecule functions by actually cleaving its target mRNA molecule instead of merely binding to it. Catalytic nucleic acid molecules can only cleave a target nucleic acid sequence if that target sequence meets certain minimum requirements. The target sequence must be complementary to the hybridizing regions of the catalytic nucleic acid, and the target
10 must contain a specific sequence at the site of cleavage.

Catalytic RNA molecules ("ribozymes") are well documented (Haseloff and Gerlach 1988; Symonds 1994; and Sun et al 1997), and have been shown to be capable of cleaving both RNA (Haseloff and Gerlach 1988) and DNA (Raillard et al 1996) molecules. Indeed, the development of *in vitro* selection and evolution techniques has
15 made it possible to obtain novel ribozymes against a known substrate, using either random variants of a known ribozyme or random-sequence RNA as a starting point (Pan 1997; Tsang and Joyce 1996; and Breaker 1994).

Ribozymes, however, are highly susceptible to enzymatic hydrolysis within the cells where they are intended to perform their function. This in turn limits their
20 pharmaceutical applications.

Recently, a new class of catalytic molecules called "DNAzymes" was created (Breaker and Joyce 1995; Santoro and Joyce 1997). DNAzymes are single-stranded, and cleave both RNA (Breaker (1994; Santoro and Joyce 1997) and DNA (Carmi et al 1998). A general model for the DNAzyme has been proposed, and is known as the "10-23"
25 model. DNAzymes following the "10-23" model, also referred to simply as "10-23 DNAzymes", have a catalytic domain of 15 deoxyribonucleotides, flanked by two substrate-recognition domains of seven to nine deoxyribonucleotides each. *In vitro* analyses show that this type of DNAzyme can effectively cleave its substrate RNA at purine:pyrimidine junctions under physiological conditions (Santoro and Joyce 1998).

Several groups have examined the activity of DNAzymes in biological systems. DNAzyme molecules targeting c-myc were found to suppress SMC proliferation after serum stimulation (Sun *et al*/1997). Two studies have explored the activity and specificity of DNAzymes targeting the *bcr-abl* fusion in Philadelphia chromosome positive leukemia cells ; Wu *et al.*, 1999). The activity of these DNAzymes compared
30 favourably with previous work with hammerhead ribozymes and antisense oligonucleotides (Gewirtz *et al.*, 1998).

More recently a 10-23 DNAzyme targeting the transcription factor Egr-1 has been shown to inhibit smooth muscle cell proliferation in cell culture and neointima formation in the rat carotid artery damaged by ligation injury or balloon angioplasty (Santiago *et al.*, 1999). Suppression of Egr-1 was also monitored at the RNA and
5 protein level in treated smooth muscle cells by northern and western blot analysis respectively. This was the first evidence of DNAzyme efficacy in vivo, and furthermore the activity displayed by this anti-Egr-1 molecule could potentially find application in various forms of cardiovascular disease such as restenosis.

10 SUMMARY OF THE INVENTION

The present inventors have determined that the level of expression of bcl-2 gene family members can be inhibited by DNAzymes.

Accordingly in a first aspect the present invention consists in a DNAzyme
15 which specifically cleaves mRNA transcribed from a member of the bcl-2 gene family, the DNAzyme comprising (a) a catalytic domain that has the nucleotide sequence GGCTAGCTACAACGA (SEQ ID No.1) and cleaves mRNA at any purine:pyrimidine cleavage site at which it is directed, (b) a binding domain contiguous with the 5' end of the catalytic domain, and (c) another binding domain contiguous with the 3' end of the
20 catalytic domain, wherein the binding domains are complementary to, and therefore hybridise with, the two regions immediately flanking the purine residue of the cleavage site within the bcl-2 gene family mRNA, at which DNAzyme-catalysed cleavage is desired, and wherein each binding domain is at least six nucleotides in length, and both binding domains have a combined total length of at least 14
25 nucleotides.

This invention also provides a method to enhance the sensitivity of malignant or virus infected cells to therapy by modulating expression level of a member of the bcl-2 gene family using catalytic DNA.

It is preferred that the bcl-2 gene family member is selected from the group
30 consisting of bcl-2, bcl-xl, bcl-w, bfl-1, brag-1, Mcl-1 and A1. It is particularly preferred that the bcl-2 gene family member is bcl-2 or bcl-xl.

BRIEF DESCRIPTION OF THE FIGURES

35 **Figure 1:** "10-23" DNAzyme (PO-DNAzyme) and its phosphorothioate modified version (PS-DNAzyme). Panel A contain illustration for 10-23 DNAzyme.

Watson-Crick interactions for DNAzyme-substrate complex is represented by generic ribonucleotides (N) in the target (top) and the corresponding DNAzyme (N) in the arms of the DNAzyme (bottom). The defined sequence in the loop joining the arms and spanning a single unpaired purine at the RNA target site of the model represents the conserved catalytic motif. Panel B shows a chemically modified version of the DNAzyme. * represents a phosphorothioate linkage.

Figure 2: Stability of phosphorothioate-modified DNAzyme oligonucleotides in human serum. DNAzymes with 1, 3, or 5 phosphorothioate linkages at each arm were incubated with fresh human serum and sampled at various time points. From each sample, intact oligonucleotides were extracted by phenol and ³²P-labelled using polynucleotide kinase. The labelled reactions were subjected to a gel electrophoresis. Percentage of intact oligos is calculated from: intensity at various time points/intensity at 0 time point x 100, as measured by PhosphoImage.

Figure 3: TMP-mediated DNAzyme transfection of PC3 cells. 2 µM FITC-labelled DNAzyme was complexed with TMP at a charge ratio of 0, 1, 3, 5, 10 and 20. The result from FACS analysis are represented.

Figure 4: Chemosensitization of PC3 cells by Bcl-xL DNAzyme. PC3 cells were treated with DNAzyme/TMP complex for 4 hours. The medium was then replaced with fresh DMEM containing 10% FBS and 5 µM Carboplatin and further incubated for 72 hours. MTS assays were performed for cell proliferation of all the samples. % cell death is derived from the percentage of OD490 from the Carboplatin-treated samples of that from untreated PC3 cells.

Figure 5: Chemosensitization of PC3 tumour cells in human xenograph mouse model (PC3) by anti-bcl-xL. Nude mice bearing established, subcutaneously growing PC3 tumour xenograft either remained untreated (saline) or were treated with DNAzyme oligo, Taxol or DNAzyme + Taxol. DNAzyme DT882 was delivered using an osmotic pump and Taxol was administrated via i.p. route weekly. Tumour size was measured at the time points indicated.

Figure 6: Chemosensitization of MDA-MB231 human xenograph breast cancer mouse model by anti-bcl-xL DNAzyme. Nude mice bearing established, subcutaneously growing MDA-MB231 tumour xenograft either remained untreated

(saline) or were treated with DNAzyme oligo, Taxol or DNAzyme + Taxol. DNAzyme DT882 was delivered using an osmotic pump and Taxol was administrated via i.p. route weekly. Tumour size was measured at the time points indicated.

- 5 **Figure 7:** Chemosensitization of MDA-MB231 human xenograph breast cancer mouse model by anti-bcl-2 DNAzyme. Nude mice bearing established, subcutaneously growing MDA-MB231 tumour xenograft either remained untreated (saline) or were treated with DNAzyme oligo, Taxol or DNAzyme + Taxol. DNAzyme DT912 was delivered using an osmotic pump and Taxol was administrated via i.p.
10 route weekly. Tumour size was measured at the time points indicated.

- Figure 8:** Western analysis of Bcl-2 expression level I in MDA-MB 231 tumors. Bcl-2 expression levels were determined by densitometry analysis of western blots of protein extracts of tumors removed from groups of 6 mice after 15 days of treatment.
15 The relative bcl-2 expression was calculated based on the ratio of Bcl-2 to β -actin levels.

- Figure 9:** Chemosensitization of human prostate tumour cells in xenograph mouse model by anti-bcl-2 DNAzyme. Nude mice bearing established, subcutaneously
20 growing PC3 tumour xenograft either remained untreated (saline) or were treated with DNAzyme oligo, Taxol or DNAzyme + Taxol. DNAzyme DT912 was delivered using an osmotic pump and Taxol was administrated via i.p. route weekly. Tumour size was measured at the time points indicated.

- 25 **Figure 10:** Chemosensitization of human melanoma tumour cells in human xenograph mouse model (518A2) by anti-bcl-2 DNAzyme SCID mice bearing established, subcutaneously growing 518A2 tumour xenograft either remained untreated (saline) or were treated with DNAzyme oligo, DTIC or DNAzyme + DTIC. DNAzyme DT912 was delivered using an osmotic pump and DTIC was administrated
30 via i.p. route weekly. Tumour size was measured at the time points indicated and the fold of tumor growth was plotted in the figure.

DETAILED DESCRIPTION OF THE INVENTION

In a first aspect the present invention consists in a DNAzyme which specifically
5 cleaves mRNA transcribed from a member of the bcl-2 gene family selected from the
group consisting of bcl-2, bcl-xl, bcl-w, bfl-1, brag-1, Mcl-1 and A1, the DNAzyme
comprising (a) a catalytic domain that has the nucleotide sequence
GGCTAGCTACAACGA (SEQ ID NO.1) and cleaves mRNA at any purine:pyrimidine
10 cleavage site at which it is directed, (b) a binding domain contiguous with the 5' end of
the catalytic domain, and (c) another binding domain contiguous with the 3' end of the
catalytic domain, wherein the binding domains are complementary to, and therefore
hybridise with, the two regions immediately flanking the purine residue of the
cleavage site within the bcl-2 gene family mRNA, at which DNAzyme-catalysed
15 cleavage is desired, and wherein each binding domain is at least six nucleotides in
length, and both binding domains have a combined total length of at least 14
nucleotides.

This invention also provides a method to enhance the sensitivity of malignant
or virus infected cells to therapy by modulating expression level of a member of the
bcl-2 gene family selected from the group consisting of bcl-2, bcl-xl, bcl-w, bfl-1,
20 brag-1, Mcl-1 and A1 using catalytic DNA (see Table 6).

In a preferred embodiment the DNAzyme is 29 to 39 nucleotides in length.

It is preferred that the bcl-2 gene family member is bcl-2 or bcl-xl. Where the
bcl-2 gene family member is bcl-2 it is preferred that the DNAzyme is selected from
those set out in Table 1. Where the bcl-2 gene family member is bcl-xl it is preferred
25 that the DNAzyme is selected from those set out in Table 2.

Where the DNAzyme cleaves bcl-2 mRNA it is further preferred that the
DNAzyme cleaves bcl-2 mRNA at position 455, 729, 1432, 1806 or 2093 (SEQ ID NO.2).
It is particularly preferred that the sequence of the DNAzyme is as set out in SEQ ID
NO 24, 45, 53, 55 or 57.

30 Where the DNAzyme cleaves bcl-xl mRNA it is further preferred that the
DNAzyme cleaves bcl-xl mRNA at position 126, 129 or 135 (SEQ ID NO.3). It is
particularly preferred that the sequence of the DNAzyme is as set out in SEQ ID NO
82, 83 or 84.

The present invention comprehends DNAzyme compounds capable of
35 modulating expression bcl-2 gene family members, in particular human bcl-2 and
bcl-xL genes. These genes inhibit apoptosis and therefore inhibitors of these genes,

particularly specific inhibitors of bcl-2 and bcl-xL such as the DNAzyme compounds of the present invention are desired as promoters of apoptosis.

More specifically, this application provides a set of DNAzymes which specifically cleaves mRNA of the bcl-2 and bcl-xL genes, comprising (a) a catalytic
5 domain that has the nucleotide sequence GGCTAGCTACAACGA (SEQ ID NO.1) and cleaves mRNA at any purine:pyrimidine cleavage site at which it is directed, (b) a binding domain contiguous with the 5' end of the catalytic domain, and (c) another binding domain contiguous with the 3' end of the catalytic domain, wherein the binding domains are complementary to, and therefore hybridise with, the two regions
10 immediately flanking the purine residue of the cleavage site within the mRNA of the bcl-2 and bcl-xL genes, respectively, at which DNAzyme-catalysed cleavage is desired, and wherein each binding domain is at least six nucleotides in length, and both binding domains have a combined total length of at least 14 nucleotides.

As used herein, "DNAzyme" means a DNA molecule that specifically
15 recognizes a distinct target nucleic acid sequence, which can be either pre-mRNA or mRNA transcribed from the target genes. The instant DNAzyme cleaves RNA molecules, and is of the "10-23" model, as shown in Figure 1, named so for historical reasons. This type of DNAzyme is described in Santoro *et al*/1997. The RNA target sequence requirement for the 10-23 DNAzyme is any RNA sequence consisting of
20 NNNNNNNR*YNNNNNNN, NNNNNNNR*YNNNNNN or NNNNNNR*YNNNNNNN, where R*Y is the cleavage site, R is A or G, Y is U or C and N is any of G, U, C, or A.

Within the parameters of this invention, the binding domain lengths (also referred to herein as "arm lengths") can be any permutation, and can be the same or
25 different. In the preferred embodiment, each binding domain is nine nucleotides in length.

In this invention, any contiguous purine:pyrimidine nucleotide pair within mRNA transcribed from a member of the bcl-2 gene family selected from the group consisting of bcl-2, bcl-xL, bcl-w, bfl-1, brag-1, Mcl-1 and A1 can serve as a cleavage
30 site. In the preferred embodiment, purine:uracil is the purine:pyrimidine cleavage site.

As used herein the term "specifically cleaves" refers to a DNAzyme which cleaves mRNA, particularly *in vivo*, transcribed from the specified gene such that the activity of the gene is modulated.

35 Targeting a DNAzyme compound to a particular nucleic acid is generally a multistep process. The process usually begins with the identification of a nucleic acid

sequence whose function is to be modulated. This may be, for example, cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the preferred targets are members of the bcl-2 gene family, in particular the nucleic acids encoding bcl-2 and bcl-xL. The targeting process also includes determination of sites within these genes for the DNAzyme catalytic activity to occur such that the desired effect, eg., detection or modulation of the proteins, will result. Within the context of the present invention, the preferred target sites are determined by a multiplex *in vitro* selection method and cell-based screening assays.

10 In applying DNAzyme-based treatments, it is important that the DNAzymes be as stable as possible against degradation in the intracellular milieu. One means of accomplishing this is by phosphorothioate modifications at both ends of the DNAzymes. Accordingly, in the preferred embodiment, two phosphorothioate linkages are introduced into both the 5' and 3' ends of the DNAzymes. In addition to 15 phosphorothioate modification, the DNAzymes can contain other modifications. These include, for example, the 3'-3' inversion at the 3' end, N3'-P5' phosphoramidate linkages, peptide-nucleic acid linkages, and 2'-O-methyl. These are well known in the art (Wagner 1995).

The DNAzymes of the present invention can be utilised for diagnostics, 20 therapeutics, and prophylaxis and as research reagents and kits. For therapeutics, an animal, preferable a human, suspected of having a disease or disorder which can be treated by modulation the expression of a member of the bcl-2 gene family, in particular bcl-2 and bcl-xL, is treated by administering DNAzyme compounds in accordance with this invention.

25 The DNAzyme compounds of this invention are useful for research and diagnostics, because these compounds hybridise to and cleave nucleic acids encoding bcl-2 and bcl-xL, enabling the assays to be easily constructed to exploit this fact. The means for the detection include, for example, conjugation of a fluorophore and a quencher to the substrate of the DNAzymes.

30 The present invention also includes pharmaceutical compositions and formulations, which comprise the DNAzyme compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. The administration can be topical, pulmonary, oral or 35 parenteral.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powders or oily bases, thickeners and the like may be necessary or desirable.

5 Composition and formulations for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules satchels or tablets.

10 The DNAzymes of the present invention can be used to increase the susceptibility of tumour cells to anti-tumour therapies such as chemotherapy and radiation therapy.

 Accordingly in certain embodiments of this invention there are provided liposomes and other compositions containing (a) one or more DNAzyme compounds of the invention and (b) one or more chemotherapeutic agents which function by a non-hybridisation mechanism. Examples of such chemotherapeutic agents include, 15 but are not limited to, anticancer drugs such as taxol, daunorubicin, dacitinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil, floxuridine, methotrexate, colchicine, vincristine, vinlastin, etoposide, cisplatin. See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed., Berkow et al eds., 20 1987, Rahway, N.J., pp 1206-1228.

 The formulation of the therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is 25 effected or diminution of the disease state is achieved. Optimal dosing schedules can be determined from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. In general, dosage is from 0.01 µg to 100 g per kg of body weight and may be given daily, weekly, monthly or yearly.

30 In a further aspect the present invention consists in a method of treating tumours in a subject, the method comprising administering to the subject a composition comprising the DNAzyme of the first aspect of the present invention.

 In a preferred embodiment the composition further comprises a chemotherapeutic agent.

35 Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated

element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

All publications mentioned in the specification are herein incorporated by reference.

5 Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority
10 date of each claim of this application.

In order that the nature of the present invention may be more clearly understood preferred forms thereof will be described with reference to the following Examples.

15 EXAMPLES

Example 1

Identification of Cleavable sites in the bcl-2 and bcl-xL mRNA.

20 Two genes, Bcl-2 and Bcl-XL were chosen as DNAzyme targets for the treatment of cancers. These genes both belong to the Bcl-2 family and both are apoptosis repressors. Their products are found in elevated levels in many cancer types including malignant melanoma, ovarian cancer, lymphoma and prostate cancer.

25 Identification of Cleavage sites in bcl-2 mRNA for DNAzyme Design:

A partial bcl-2 cDNA clone was generated from cellular RNA, which contained 31 bp of the 5' UTR, 720 bp of ORF and 2.2 kb of the 3' UTR sequences. By scanning the mRNA corresponding to the bcl-2 clone, 210 potential AU and GU cleavage sites were identified and these sites were further subjected to two thermodynamic analyses.
30 The first analysis was on the thermodynamic stability of the enzyme-substrate heteroduplex as predicted by the hybridisation free energy (Sugimoto et al., 1995, Cairns et al., 1999). DNA enzymes with the greatest heteroduplex stability indicated by a low free energy of hybridisation (calculated using the nearest neighbour method), was often found to have the greatest kinetic activity. The selection parameters
35 included a cut-off value of $-\Delta G^\circ$ kcal/mol of less than 25. The second analysis was to examine if the arms of the DNAzyme had a high hairpin melting temperature (T_m),

thus to avoid any intramolecular bonds (Cairns et al., 1999; Santoro & Joyce 1998). After completion of these analyses, 55 (fifty-five) DNAzymes were designed and synthesised for in vitro multiplex selection. The sequence of these DNAzymes is set out in Table 1.

5

Table 1. Summary list of bcl-2 DNAzymes.

Unmodified ¹	SEQ. ID NO	2/2 PS ³	Sequence	-ΔG° kcal/mol	Activity <i>In vitro</i> ²	Activity In Cells ⁴
DT564	7		cgtagcgccaggctagctacaacgaattcccag	28.8		
DT565	8		tcccggtaggctagctacaacgacgtaccctg	28.6		
DT566	9	DT891	tcatactaggctagctacaacgactcccgggt	26	+	NO
DT567	10		cgcatcccaggctagctacaacgacgtagccc	30.7		
DT568	11		tctccgcaggctagctacaacgacccactcgt	31.6		
DT569	12		gcgcccacaggctagctacaacgactcccgc	33		
DT570	13		cggcgcccaggctagctacaacgaatctccgc	33.8		
DT571	14		aggagaaggaggctagctacaacgagcccggcgc	28.9		
DT572	15		gcggctgtaggctagctacaacgagggcggtgt	30.4		
DT573	16		gtcccgggaggctagctacaacgagcggtgta	31.2		
DT574	17	DT892	tcctggcgaggctagctacaacgacgggtccc	31.7	+++	NO
DT575	18	DT893	caggtaggaggctagctacaacgacgggtgag	27.7	+++	NO
DT576	19	DT894	ggtggaccaggctagctacaacgaagggtggcac	27.8	+++	NO
DT577	20		gcctggacaggctagctacaacgactcggcgaa	27.6		
DT578	21		ctgcctggaggctagctacaacgaatctcggcg	28.1		
DT579	22		cctccaccaggctagctacaacgacgtggcaaa	27.6		
DT580	23		gctcctccaggctagctacaacgacccgtggc	31.4		
DT581	24	DT895	cccagttcaggctagctacaacgacccgtccct	32.8	+	YES
DT582	25		aggccacaaggctagctacaacgacctcccca	32.3		
DT583	26		agaaggccaggctagctacaacgaaatctccc	27		
DT584	27		cacacatgaggctagctacaacgacccaccgaa	25.1		
DT585	28		ccacacacaggctagctacaacgagacccacc	29.5		
DT586	29		ctccacacaggctagctacaacgaatgaccca	27.7		
DT587	30		ctctccacaggctagctacaacgaacatgaccc	26.3		
DT588	31		cccggtaggctagctacaacgagctctccac	29.6		
DT589	32		ggggcgacaggctagctacaacgactcccgggt	30.7		

Unmodified ¹	SEQ. ID NO	2/2 PS ³	Sequence	$-\Delta G^\circ$ kcal/mol	Activity <i>In vitro</i> ²	Activity In Cells ⁴
DT590	33		caggggagcaggctagctacaacgaatctcccg	29.6		
DT591	34		tgtgtccaggctagctacaacgacagggcgga	27.5		
DT592	35		acagggcagaggctagctacaacgagttgtccac	26.7		
DT593	36	DT896	agtcattcaggctagctacaacgaaggcgatg	25.4	+	NO
DT594	37	DT897	actcagtcaggctagctacaacgaccacagggc	26.7	+	NO
DT595	38		tatcctggaggctagctacaacgaccaggtgtg	25.5		
DT596	39	DT898	cctccgttaggctagctacaacgacctggatcc	28.6	+	NO
DT597	40		acaaaggcaggctagctacaacgacccagcctc	27.4		
DT598	41	DT899	ggggccgtaggctagctacaacgaagtccaca	28.8	+	YES
DT599	42		gaggccgcaggctagctacaacgagctggggcc	33.4		
DT600	43	DT900	aagctcccaggctagctacaacgacagggccaa	28.4	+	YES
DT601	44	DT901	ccagggtgaggctagctacaacgagcaagctcc	28.2	+	NO
DT602	45	DT902	agataggcaggctagctacaacgaccaggtga	25.3	+	YES
DT603	46	DT903	tggcccaggctagctacaacgaaggcaccca	31.3	+	NO
DT604	47	DT904	ttgacttcaggctagctacaacgattgtggccc	25.7	+	NO
DT605	48	DT905	gggcaggcaggctagctacaacgagttgacttc	26.5	+++	YES
DT606	49		ggagccacaggctagctacaacgagaagcggtg	26.5		
DT607	50	DT906	cccaatgaggctagctacaacgacaggtcctt	27.5	+++	NO
DT608	51		agggaggcaggctagctacaacgaggactccc	28.5		
DT609	52	DT907	ttctcccaggctagctacaacgacaggtatgc	27.5	+++	NO
DT610	53	DT908	ttttcccaggctagctacaacgacgtgtcct	27.9	+	YES
DT611	54	DT909	gcggcctgaggctagctacaacgagcttgggt	31.1	+++	NO
DT612	55	DT910	ccctgttgaggctagctacaacgacatccctgg	28.4	+++	YES
DT613	56	DT911	tggctcccaggctagctacaacgagctccacgt	31.4	+++	NO
DT614	57	DT912	cacagccaaggctagctacaacgagtgccatgt	26.7	+++	YES
DT615	58	DT913	accccataggctagctacaacgatccacact	30.1	+++	NO
DT616	59	DT914	cagggttaggctagctacaacgactcaccttc	25.6	+++	NO
DT617	60	DT915	gccagggaggctagctacaacgagaggaaacc	27.1	+++	NO
DT618	61	DT916	tgctggtcaggctagctacaacgattgccatct	26.5	+++	NO

2. In the process of in vitro selection, twenty-six active DNazymes were identified (26/55).

3. The in vitro selected DNAzymes were further chemically modified using two phosphorothioate linkages at both ends and renamed as indicated (Wagner 1995).
4. The modified DNAzymes were subjected to cell-based assay in which the bcl-2 protein level was measured by Western blots. Eight DNAzymes were shown active in
- 5 down-regulation of Bcl-2 protein.

Identification of Cleavage sites in bcl-xL mRNA for DNAzyme Design:

As for the bcl-2 DNAzyme selection, total of 26 DNAzymes were designed and synthesised for the bcl-xL mRNA, based on the sequence scanning, and $-\Delta^\circ G/T_m$ analyses. The sequence of these DNAzymes is set out in Table 2.

Table 2. Summary list of bcl-xL DNAzymes.

Unmodified ¹	SEQ. ID NO	2/2 PS ³	Sequence	$-\Delta^\circ G$ kcal/mol	<i>In vitro</i> Activity ²	Activity In cells ⁴
DT673	62	DT861	aagagttcaggctagctacaacgatcactacct	21.70	+	
DT674	63	DT862	tttaccacaggctagctacaacgacccggaaga	27.70	+	
DT675	64	DT863	cccagtttaggctagctacaacgacccatcccg	30.10	+	
DT676	65	DT864	acaatgcgaggctagctacaacgacccagttta	23.60		
DT677	66	DT865	aggccacaaggctagctacaacgagcgacccca	30.40		
DT678	67	DT866	aaaaggccaggctagctacaacgaaatgcgacc	22.70		
DT679	68	DT867	ttccacgcaggctagctacaacgaagtgcctcg	30.90		
DT680	69	DT868	cgctttccaggctagctacaacgagcacagtgc	27.60	+	
DT681	70	DT869	cctgtctaggctagctacaacgagctttccac	25.80	++	
DT682	71	DT870	atacctgcaggctagctacaacgactcctgtc	25.60	+++	
DT683	72	DT871	tcaccaataggctagctacaacgactgcattc	22.50	++	
DT684	73	DT872	actaccaaggctagctacaacgaacctgcattc	24.80		
DT685	74	DT873	tccgactcaggctagctacaacgacaatacctg	23.20		
DT686	75	DT874	gcgatccgaggctagctacaacgatcaccaata	24.40	+	
DT687	76	DT875	aagctgcgaggctagctacaacgacccgactcac	25.40	+	
DT688	77	DT876	aagtggccaggctagctacaacgaccaagtgc	27.10	++	++
DT689	78	DT877	aggtggtcaggctagctacaacgatcaggttaag	22.80	++	++
DT690	79	DT879	tctcctggaggctagctacaacgaccaaggctc	27.10	+++	
DT691	80	DT880	acaaaagtaggctagctacaacgacccagccgc	25.20		
DT692	81	DT881	gtctggtcaggctagctacaacgattccgactg	25.40	+++	+

Unmodified ¹	SEQ. ID NO	2/2 PS ³	Sequence	$-\Delta^\circ G$ kcal/mol	<i>In vitro</i> Activity ²	Activity In cells ⁴
DT693	82	DT882	ttttataaggctagctacaacgaaggatggg	18.90	+	+++
DT694	83	DT883	acatttttaggctagctacaacgaataggat	17.40	+	+
DT695	84	DT884	tctgagacaggctagctacaacgattttataat	16.80	+	+
DT696	85	DT885	gctctgagaggctagctacaacgaattttata	19.40	+++	
DT697	86	DT886	agtcaaccaggctagctacaacgacagctccg	27.50	++	
DT698	87	DT887	gtggctccaggctagctacaacgatcaccgcgg	30.50		

1. After screening the 926-bp bcl-xL cDNA clone, 81 potential cleavable AU or GU sites were found and these sites were subjected to thermodynamic analyses. Based on the threshold of -25 kcal/mol as selection criteria, twenty-six DNAzymes were synthesized for *in vitro* cleavage selection (26/81).
2. In the process of *in vitro* selection, eighteen active DNAzymes were identified (18/26).
3. All twenty-six DNAzymes were further chemically modified using two phosphorothioate linkages at both ends and renamed as indicated.
4. The modified DNAzymes were subjected to cell-based assay in which the bcl-xL protein level was measured by Western blots. Six DNAzymes were shown active in down-regulation of Bcl-xL protein (6/26).

Example 2

Multiplex Selection of Active DNAzymes *in vitro*:

In order to efficiently select active DNAzymes, *in vitro* selection was performed using a multiplex method, which enables a pool of DNAzymes to be screened for their ability to access and cleave RNA substrate under simulated physiological conditions (Cairns et al., 1999). The DNAzymes (0nM, 5nM, 50nM and 500nM) and RNA substrate (400nM) were pre-equilibrated separately for 10 min at 37°C in equal volumes of 50mM Tris-HCL, pH 7.5 10mM MgCl₂, 150mM NaCl and 0.01% SDS. Reaction was initiated by mixing the DNAzymes and substrate together. After 1 hr the reaction was stopped by extraction in 100μl phenol/chloroform and recovered by ethanol precipitation.

The primers for bcl-2 cleavage detection are:

5'-cacagcattaaacattgaacag-3' (SEQ ID NO.90)

5'-tggaactttttttgtcagg-3' (SEQ ID NO.91)

5'-tcctcacgttcccagccttc-3' (SEQ ID NO.92)
 5'- cagacattcgagaccacac-3' (SEQ ID NO.93)
 5'-cagtattgggagttgggggg-3' (SEQ ID NO.94)
 5'- ccaactcttttctcccacc-3' (SEQ ID NO.95)
 5'-cgacgttttgctgaagactg-3' (SEQ ID NO.96)
 5'- cagggccaaactgagcagag-3' (SEQ ID NO.97)
 5'-atctccccagttcacc-3' (SEQ ID NO.98)
 5'- ggatgcggctgtatgggg-3' (SEQ ID NO.99); and
 5'-aggccacgtaaagcaactctc-3' (SEQ ID NO.100).

10

The primers for bcl-xL cleavage detection are:

5'-cgggttctctggtggca-3' (SEQ ID NO.101)
 5'-ccttcggctctcggctg-3' (SEQ ID NO.102)
 5'-ccgccgaaggagaaaaag-3' (SEQ ID NO.103); and
 5'-gcctcagtcctgttctctcc-3' (SEQ ID NO.104).

15

Primer extension was then performed with Superscript II reverse transcriptase. In this reaction 4pmol of labelled primer was combined with 300nmol of RNA and denatured at 90°C for 2 min. The primer was then allowed to anneal slowly between 65°C-45°C before adding the first strand buffer, dithiothreitol, deoxynucleotides and enzyme. This mix (20µl) was incubated at 45°C for 1hr, before being stopped by placing the reaction on ice. Samples were placed in an equal volume of stop buffer and then run on a 6% polyacrylamide gel. Sequencing was performed by primer extension on the double stranded cDNA template in the presence of chain terminating dideoxynucleotides (ddNTP)(Sambrook et al., 1989). The sequence was used as a guide to attribute cleavage bands to specific DNazymes. The relative cleavage strength of each DNzyme was determined by intensity of the cleavage products. DNazymes were ranked according to their cleavage ability at lowest concentration (5nM). *In vitro* selection of bcl-2 DNazymes was achieved by incubating Bcl-2 DNazymes with its RNA substrate for 60 minutes in the presence of 10 mM Mg²⁺ at 37°C. Primer extension was performed using the sequence-specific primers along the bcl-2 mRNA. The reactions were analysed alongside with DNA sequencing on a polyacrylamide gel. *In vitro* selection of bcl-xL DNazymes was achieved by incubating Bcl-xL DNazymes with its RNA substrate for 60 minutes in the presence of 10 mM Mg²⁺ at 37°C. Primer extension was performed using the sequence-specific

20

25

30

35

primers along the bcl-xL mRNA. The reactions were analysed alongside with DNA sequencing on a polyacrylamide gel.

Example 3

5

Porphyrin-mediated DNzyme uptake in cancer cells

To test the selected DNzymes in cell culture systems, a prostate cancer cell line PC3 was initially used to examine their efficacy in down-regulation of bcl-2 and bcl-xL gene expression and impact on cellular functions. To facilitate delivery of
10 DNzyme oligonucleotides into cells, a cationic porphyrin, tetra meso-(4-methylpyridyl) porphyrine (TMP), was used as a transfection reagent for intracellular delivery (Benimetskaya et al., 1998).

Chemical modification of DNzymes:

15 To increase DNzyme stability in cells, two phosphorothioate linkages were incorporated into each of the arms in DNzymes (PS-Dz)(Wagner et al. 1995). This has been shown to increase the DNzyme stability significantly in human serum, while there was no marked effect on the DNzyme cleavage activity (Figure 2).

20 DNzyme transfection efficiency:

1.2 x 10⁶ cells were seeded in a 100-mm culture dish and incubated at 37°C, 5% CO₂ overnight. The cells were transfected with an FITC-labelled DNzyme that was complexed with TMP at a charge ratio of 3 (+/-). The transfected cells were analysed using FACS and fluorescent microscopy. As shown in Figure 3, a more efficient
25 delivery was observed when POS-Dz was complexed with TMP, compared with normal phosphodiester DNzyme (PO-Dz). In addition, nuclear delivery of the DNzymes (FITC-labelled) was evident.

Example 4

30

Suppression of bcl-2 and bcl-xL expression in cancer cells.

From the *in vitro* multiplex selection, 26 DNzymes against bcl-2 (26/55) and 16 DNzymes against bcl-xL (16/26) were shown to be efficient cleavers of their corresponding substrates. The modified version of these molecules were then tested
35 for their ability to down regulate the bcl-2 and bcl-xL expression in cells. The assays were performed in PC3 cells (a prostate cancer cell line). The cells were transfected

with 2 μ M DNAzyme complexed with TMP at a charge ratio of 3. After overnight incubation, cells were subject to either protein (Western blot) or RNA (Ribonuclease protection assay) analyses (Sambrook et al 1989).

5 Effect of Bcl-2 DNAzymes on bcl-2 expression in PC3 cells:

All 26 DNAzymes were tested in transfection assay for their activity by Western blots. Five out of 26 DNAzymes showed a consistent inhibitory effect on the bcl-2 protein level (Table 3). The effect of bcl-2 DNAzymes on expression of the bcl-2 gene family was determined by transfecting five active DNAzymes into PC3 cells (2 μ M). DT907 was used as an inactive DNAzyme control. Antibodies to Bcl-2, Bcl-xL, Bax and β -actin were used respectively to detect the corresponding proteins. While TMP alone and inactive DNAzyme control did not show any effect, all the five DNAzymes suppressed Bcl-2 level significantly. These DNAzymes had no effect on either other members of the bcl-2 gene family such as Bcl-xL and Bax, or house
15 keeping gene β -actin.

Table 3: Active bcl-2 DNAzymes identified in Western analyses.

DNAzyme	DNAzyme sequence	Target sites*
DT895	Cccagttcaggctagctacaacgaccggtcct	455
DT902	Agataggcaggctagctacaacgaccaggggtga	729
DT908	Ttttcccaggctagctacaacgacgctgtcct	1432
DT910	Ccctgttgaggctagctacaacgacatccctgg	1806
DT912	Cacagccaaggctagctacaacgagtgccatgt	2093

* indicates the cleavage site on human bcl-2 mRNA sequence.

20

Effect of bcl-xL DNAzymes on the bcl-xL expression:

After screening all 16 DNAzymes, three DNAzymes, DT882, DT883 and DT884, exhibited a very strong inhibitory effect on bcl-xL protein expression (Table 4). Suppression of bcl-xL protein level by bcl-xL DNAzymes was determined by
25 transfecting three active DNAzymes into PC3 cells (2 μ M). DT867 and 880 were used as inactive DNAzyme controls; and DT888 as an antisense control. Antibodies to Bcl-2 and β -actin were used respectively to detect the corresponding proteins. DT 880 and DT867 were inactive DNAzymes in this screening. The effect in PC3 cells was further confirmed using an RNase protection assay (RPA) of bcl-xL DNAzyme. In the RNase
30 protection assay, DNAzymes were complexed with TMP at a charge ratio of 3 and

transfected into PC3 cells. Cellular RNA was extracted from the transfected cells and used for RPA analysis. Apoptosis related riboprobe set was generated from a Pharmingen kit.

5

Table 4: Active bcl-xL DNAszymes identified in Western analyses.

DNAszyme	DNAszyme sequence	Target sites*
DT882	Ttttataaggctagctacaacgaagggatggg	126
DT883	Acatttttaggctagctacaacgaaatagggat	129
DT884	Tctgagacaggctagctacaacgattttataat	135

* indicates the cleavage site on human bcl-xL mRNA sequence.

10 *Example 5*

Bcl-2 and bcl-xL specific DNAszyme-mediated effect on cell cycle

Following the test of the DNAszymes in Western and RPA assays, some of the active molecules were further examined for their effect on cell cycle as an indication of apoptotic response. Two most active DNAszymes were chosen in FACS assay. These were DT 895 (a bcl-2 DNAszyme) and DT882 (a bcl-xL DNAszyme). In the assay, same transfection procedure as in Western assay was used, except those cells were subject to PI staining after the overnight incubation with the DNAszymes. Table 5 clearly showed that there was a substantial increase in sub G1 population in the DNAszyme treated cells (DT895 12.82% and DT882 23.17% respectively), indicating that the cells treated with anti-bcl-2 and bcl-xL DNAszymes were provoked to undergo apoptosis.

Table 5. Cell cycle analysis of DNAszyme-transfected PC3 cells.

Treatment	% Sub-G1 population
PC3	0.63
TMP	2.2
Bcl-2 DNAszyme 895	12.82
Bcl-xL DNAszyme 882	23.17
Inactive control	1.62

*Example 6***Effect of the Bcl-xL DNzyme on Cytochrome C Release**

Cytochrome c is a well-characterised mobile electron transport protein essential to energy conversion in all-aerobic organisms. In mammalian cells, this highly conserved protein is normally localised to the mitochondrial intermembrane space. More recent studies have identified cytosolic cytochrome c as a factor necessary for activation of apoptosis. During apoptosis, cytochrome c is translocated from the mitochondrial membrane to the cytosol, where it is required for activation of caspase-3 (CPP32). It has been reported that the translocation of cytochrome c can be blocked by overexpression of Bcl-2 or Bcl-xL. Based on this, the measurement of CytoC release from cells would be an ideal assay to determine the bcl-xL DNzyme effect on the early events of apoptosis caused by down-regulation of bcl-xL. After transfection of PC3 cells with bcl-xL DNzymes, the proteins from the cytoplasmic fraction were extracted and subjected to Western analysis. Studies by the applicants determined that Bcl-xL DNzyme-mediated down-regulation of bcl-xL and increased release of Cytochrome C. In these studies PC3 cells were transfected with 2 μ M DNzyme complexed with TMP. Western analyses were performed using the antibodies to Bcl-xL and Cytochrome C. DNzyme-mediated reduction of bcl-xL in PC3 cells led to an increased release of CytoC. This result not only confirmed previous data from cell cycle analysis, but also validated the specificity of the DNzyme against apoptotic pathway in PC3 cells.

Example 7

25

Chemosensitization of PC3 Cells with Anti-bcl-xL DNzymes

The Bcl-xL protein has been shown in a number of cell lines to be a potent protector of cellular apoptosis induced by anti-neoplastic agents. Thus an efficient DNzyme that decreased Bcl-xL expression in PC3 cells would sensitise them to the effect of cytotoxic therapy. To test this, cell survival was measured using MTS assays in PC3 cells treated with either DNzyme alone or DNzyme plus anti-cancer agents such as Carboplatin. The result in Figure 4 demonstrated that the anti-bcl-xL DNzyme DT882 sensitised PC3 cells to Carboplatin treatment at 5 μ M. This sensitization led to an increase of cell death from 17% when only Carboplatin was used, to about 50% cell death when the DNzyme and Carboplatin were combined.

*Example 8***Use of Anti-bcl-2 and bcl-xL DNazymes in Other Tumour Cell Lines**

High level expression of Bcl-2 and Bcl-xL has been found in various types of
5 cancers. In addition to the efficacy of the DNazymes shown in Prostate cancer cell
lines (PC3 and DU145), further cell-based assays were performed to explore the
therapeutic potential of the anti-bcl-2 and bcl-xL DNazymes in vivo. Several cell lines
of various cancer types have been used to validate the DNzyme efficacy in the
different settings. These are T24, bladder cancer; HCT116, colon cancer; and A549,
10 lung carcinoma.

To analyse inhibition of Bcl-2 expression in different tumour cells by bcl-2
DNazymes, T24 (bladder), A549 (lung) and HCT116 (colon) cells were treated with 2
 μ M DNzyme complexed with TMP at a charge ratio of 3. After 24 hours post
transfection, the cellular protein was extracted and immunoblotted with bcl-2
15 antibody or β -actin antibody. Inhibition of Bcl-xL expression in different tumour cells
by bcl-xL DNzyme was also investigated using T24 (bladder), A549 (lung) and
HCT116 (colon) cells treated with 2 μ M DNzyme complexed with TMP as described
and immunoblotted with bcl-xL antibody or β -actin antibody. These studies show that
both anti-bcl-2 and anti-bcl-xL DNazymes reduced the level of their respective gene
20 expression in all the cell lines tested.

*Example 9.***Chemosensitization in human tumour xenograph models by anti-bcl-xL DNzyme
25 DT882.**

In order to demonstrate that down-regulation of the bcl-2 gene family results in
Chemosensitization of tumour cells to anticancer drugs, murine models with human
PC3 prostate cancer and MDA-MB-231 breast cancer xenograph were used to
determine if the sensitivity to the chemotherapeutic is enhanced.

30 In the experiments, four groups of mice (8 mice per group) (Saline, DNzyme,
Taxol, Taxol + DNzyme) were employed At day 1: acclimatised nude male Balb/C
athymic mice were injected with 1×10^6 tumor cells suspended in 0.1 ml Matrigel in the
right hind leg under methoxyfluorane anesthesia. Tumour growth is measured twice
weekly using digital callipers and tumour volume is calculated using the $(l \times w \times h \times$
35 $\pi/6)$ formula. When tumours reach an average volume of 100-200 mm³, an Alzet
osmotic pump, which were used as a delivery vehicle for DNzyme oligonucleotides

in tumour bearing mice, was surgically implanted in the peritoneum of the mouse via the abdominal route. The Alzet model1002 pump is a capsule shaped pump (1.5 x 0.6 cm) and delivers a total volume of 0.5 ml at a rate of 0.25 μ l/hr over a period of 14 days. The pump was filled with a saline solution containing DNAzyme

- 5 oligonucleotide, which resulted in a dose rate of 12.5 mg/kg/d. Some mice will receive 25 mg/kg Taxol by intraperitoneal route in a 200 μ l injection once weekly post-surgery for the duration of the study.

As shown in Figures 5 and 6, combination of DNAzyme and Taxol treatment markedly inhibited both PC3 and MDA-MB-231 tumour growth compared with the
10 groups of DNAzyme alone or Taxol alone.

Example 10

Chemosensitization in human tumour xenograph models by anti-bcl-2 DNAzyme 15 DT912.

As described in Example 9, both prostate and breast cancer models were also used in testing the bcl-2 DNAzyme efficacy. In addition, a human melanoma model (518A2) was further used to determine the effect of the treatment of bcl-2, combined with Dacarbazine (DTIC), on the tumor growth. As shown in figures 7, 9 and 10, the
20 anti-bcl-2 DNAzyme DT912 could sensitise all three tumours to chemotherapeutic treatment and this effect was closely related to the down-regulation of the bcl-2 protein level (Figure 8).

Example 11

25

Accessibility and efficacy of antisense oligonucleotides cannot be correlated to DNAzyme targeting.

The protooncogene c-myc plays an important role in proliferation and differentiation of haematopoietic cells. C-myc protein levels vary according to the
30 level of differentiation of normal haematopoietic cells with low protein expression detected in terminally differentiated cells. In leukemia cells where there is rapid proliferation of myeloid precursors, c-myc has often been found to be overexpressed. In the literatures, it has been shown that use of antisense oligonucleotides could inhibit the c-myc expression in vitro and led to suppress leukemia development.
35 Against same regions targeted by antisense oligonucleotides, DNAzymes were designed and tested in leukemia cell cultures.

In the experiments, K562 cells were transfected with 2 μ M oligo complexed with TMP at a charge ratio (+/-) of 5 on Days 0 and 1. Cellular proteins were extracted on Day 2 and analysed by Western using a monoclonal antibody to c-Myb. Inhibition of c-Myb protein expression by antisense and DNAzymes was determined by Western blot analysis. Two antisense oligonucleotides DT860 (gtgccggggtcttcgggc,) (SEQ ID NO. 105) and DT1019 (gctttgcgatttctg;)(SEQ ID NO.106), consistently showed efficacy in inhibiting c-Myb protein expression, while none of the DNAzymes corresponding to these sites could effectively reduce c-Myb protein levels

This example clearly demonstrates that the different structures and conformations of oligonucleotides and DNAzymes results in these two molecules having different accessibility to their target RNA. Thus, the effect of the one type of agent is not a predictor of the activity of another type.

Table 6: Sequence ID Nos and description.

SEQUENCE ID NO.	Description	Database Accession Number
1	DNAzyme catalytic domain	
2	Bcl-2 CDS	M14745
3	Bcl-xL CDS	Z23115
4	Bcl-w gene	NM_004050
5	Bfl-1 gene	U27467
6	Mcl-1 gene	AF147742
7-61	Bcl-2 DNAzymes	
62-87	Bcl-xL DNAzymes	
88	Bcl-2 A1 gene	NM_004049
89	BRAG-1 gene	S82185
90-100	bcl-2 cleavage detection primers	
101-104	bcl-xL cleavage detection primers	
105	Antisense oligonucleotide	
106	Antisense oligonucleotide	

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as
5 illustrative and not restrictive.

References

- Adams, J. M., and Cory, S. (1998). The Bcl-2 protein family: arbiters of cell survival.
5 Science 281, 1322-6.
- Ashkenazi, A., and Dixit, V. M. (1998). Death receptors: signaling and modulation.
Science 281, 1305-8.
- 10 Benimetskaya, L., Takle, G. B., Vilenchik, M., Lebedeva, I., Miller, P., and Stein, C. A.
(1998). Cationic porphyrins: novel delivery vehicles for antisense
oligodeoxynucleotides. *Nucleic Acids Res* 26, 5310-7.
- Breaker RR and Joyce GF (1994) A DNA enzyme that cleaves RNA. *Chem Biol* 1:223-
15 229.
- Breaker RR and Joyce GF (1995) A DNA enzyme with Mg(2⁺)-dependent RNA
phosphoesterase activity. *Chem Biol* 2:655-660.
- 20 Cairns, M. J., Hopkins, T. M., Witherington, C., Wang, L., and Sun, L. Q. (1999). Target
site selection for an RNA-cleaving catalytic DNA [see comments]. *Nat Biotechnol* 17,
480-6.
- Carmi N, Balkhi SR and Breaker RR. (1998) Cleaving DNA with DNA. *Proc Natl Acad*
25 *Sci USA*. 95:2233-2237.
- Ellis, R. E., Yuan, J. Y., and Horvitz, H. R. (1991). Mechanisms and functions of cell
death. *Annu Rev Cell Biol* 7, 663-98.
- 30 Gewirtz, A. M., Sokol, D. L., and Ratajczak, M. Z. (1998). Nucleic acid therapeutics:
state of the art and future prospects. *Blood* 92, 712-36.
- Haseloff J and Gerlach WL (1988) Simple RNA enzymes with new and highly specific
endoribonuclease activities *Nature* 334:585-591.
35

Jansen, B., Schlagbauer-Wadl, H., Brown, B. D., Bryan, R. N., van Elsas, A., Muller, M., Wolff, K., Eichler, H. G., and Pehamberger, H. (1998). Bcl-2 antisense therapy chemosensitizes human melanoma in SCID mice. *Nat Med* 4, 232-4.

- 5 Kashani-Sabet M, Funato T et al (1992) Reversal of malignant phenotype by an anti-ras ribozyme. *Antisense Res Dev* 2:3-15.

Koizumi M, Kamiya H and Ohtsuka E (1993) Inhibition of c-Ha-ras gene expression by hammerhead ribozymes containing a stable C(UUCG)G hairpin loop. *Biol Pharm Bull*
10 16:879-883.

Pan T (1997) Novel and variant ribozymes obtained through in vitro selection. *Curr Opin Chem Biol.* 1:17-25.

- 15 Raillard SA and Joyce GF (1996) Targeting sites within HIV-1 cDNA with a DNA-cleaving ribozyme. *Biochemistry.* 35(36):11693-701

Reed, J. C. (1998). Bcl-2 family proteins. *Oncogene* 17, 3225-36.

- 20 Reed, J. C. (2000). Mechanisms of Apoptosis. *Am J Pathol* 157, 1415-1430.

Sambrook J., Fritsch E.F. and Maniatis T., 1989 *Molecular Cloning, a laboratory manual* 2nd ed. Cold Spring Harbour Laboratory Press.

- 25 Santiago, F. S., Lowe, H. C., Kavurma, M. M., Chesterman, C. N., Baker, A., Atkins, D. G., and Khachigian, L. M. (1999). New DNA enzyme targeting Egr-1 mRNA inhibits vascular smooth muscle proliferation and regrowth after injury [published erratum appears in *Nat Med* 5(12): 1438. *Nat Med* 5, 1264-9.

- 30 Santoro, S. W., and Joyce, G. F. (1997). A general purpose RNA-cleaving DNA enzyme. *Proc Natl Acad Sci U S A* 94, 4262-6.

Santoro SW and Joyce GF (1998) Mechanism and utility of an RNA-cleaving DNA enzyme. *Biochemistry* 37:13330-13342.

Sugimoto, N., Nakano, S., Katoh, M., Matsumura, A., Nakamuta, H., Ohmichi, T., Yoneyama, M., and Sasaki, M. (1995). Thermodynamic parameters to predict stability of RNA/DNA hybrid duplexes. *Biochemistry* 34, 11211-6.

- 5 Sun, L. Q., Cairns, M. J., Gerlach, W. L., Witherington, C., Wang, L., and King, A. (1999). Suppression of smooth muscle cell proliferation by a c-myc RNA-cleaving deoxyribozyme. *J Biol Chem* 274, 17236-41.

- 10 Tsang J and Joyce GF (1996) Specialization of the DNA-cleaving activity of a group I ribozyme through in vitro evolution. *J Mol Biol.* 262:31-42.

- 15 Vocero-Akbani, A. M., Heyden, N. V., Lissy, N. A., Ratner, L., and Dowdy, S. F. (1999). Killing HIV-infected cells by transduction with an HIV protease-activated caspase-3 protein [see comments]. *Nat Med* 5, 29-33.

- 20 Warashina, M., Kuwabara, T., Nakamatsu, Y., and Taira, K. (1999). Extremely high and specific activity of DNA enzymes in cells with a Philadelphia chromosome. *Chem Biol* 6, 237-50.

Wagner, R.W. (1995) *Nature Med* 1:1116-1118.

- 25 Waters, J. S., Webb, A., Cunningham, D., Clarke, P. A., Raynaud, F., di Stefano, F., and Cotter, F. E. (2000). Phase I clinical and pharmacokinetic study of bcl-2 antisense oligonucleotide therapy in patients with non-Hodgkin's lymphoma [see comments]. *J Clin Oncol* 18, 1812-23.

- 30 Wu, Y., Yu, L., McMahon, R., Rossi, J. J., Forman, S. J., and Snyder, D. S. (1999). Inhibition of bcr-abl oncogene expression by novel deoxyribozymes (DNAzymes). *Hum Gene Ther* 10, 2847-57.

Yakovlev, A. G., Knoblach, S. M., Fan, L., Fox, G. B., Goodnight, R., and Faden, A. I. (1997). Activation of CPP32-like caspases contributes to neuronal apoptosis and neurological dysfunction after traumatic brain injury. *J Neurosci* 17, 7415-24.

- Zangemeister-Wittke, U., Leech, S. H., Olie, R. A., Simoes-Wust, A. P., Gautschi, O., Luedke, G. H., Natt, F., Haner, R., Martin, P., Hall, J., Nalin, C. M., and Stahel, R. A. (2000). A novel bispecific antisense oligonucleotide inhibiting both bcl-2 and bcl-xL expression efficiently induces apoptosis in tumour cells [In Process Citation]. Clin
- 5 Cancer Res 6, 2547-55.

CLAIMS:

1. A DNzyme which specifically cleaves mRNA transcribed from a member of the bcl-2 gene family selected from the group consisting of bcl-2, bcl-xl, bcl-w, bfl-1,
5 brag-1, Mcl-1 and A1, the DNzyme comprising (a) a catalytic domain that has the nucleotide sequence GGCTAGCTACAACGA and cleaves mRNA at any purine:pyrimidine cleavage site at which it is directed, (b) a binding domain contiguous with the 5' end of the catalytic domain, and (c) another binding domain contiguous with the 3' end of the catalytic domain, wherein the binding domains are
10 complementary to, and therefore hybridise with, the two regions immediately flanking the purine residue of the cleavage site within the bcl-2 gene family mRNA, at which DNzyme-catalysed cleavage is desired, and wherein each binding domain is at least six nucleotides in length, and both binding domains have a combined total length of at least 14 nucleotides.
- 15 2. A DNzyme according to claim 1 wherein the DNzyme is 29 to 39 nucleotides in length.
3. A DNzyme according to claim 1 or claim 2 wherein the bcl-2 gene family member is bcl-2 or bcl-xl.
4. A DNzyme according to any one of claims 1 to 3 wherein is selected from
20 those listed in SEQ ID NOS. 7 to 61.
5. A DNzyme according to any one of claims 1 to 4 wherein the DNzyme cleaves bcl-2 mRNA at position 455, 729, 1432, 1806 or 2093.
6. A DNzyme according to any one of claims 1 to 5 wherein the sequence of the DNzyme sequence as set out in SEQ ID NOS 24, 45, 53, 55 or 57.
- 25 7. A DNzyme according to any one of claims 1 to 3 wherein is selected from those listed in SEQ ID NOS. 62 to 87.
8. A DNzyme according to any one of claims 1 to 3 or 7 wherein the DNzyme cleaves bcl-xl mRNA at position 126, 129 or 135.
9. A DNzyme according to any one of claims 1 to 3 or 7 or 8 wherein the
30 sequence of the DNzyme sequence as set out in SEQ ID NOS 82, 83 or 84.
10. A DNzyme according to any one of claims 1 to 9 wherein 1 to 6 phosphorothioate linkages are introduced into each of the 5' and 3' ends of the DNzymes.
11. A DNzyme according to any one of claims 1 to 10 wherein the DNzyme
35 comprises at least one modification selected from the group consisting of 3'-3'

inversion, N3'-P5' phosphoramidate linkages, peptide-nucleic acid linkages, and 2'-O-methyl.

12. A pharmaceutical compositions, the composition comprising a pharmaceutically acceptable carrier and at least one DNAzyme according to any one of claims 1 to 11.

13. A pharmaceutical composition according to claim 13 wherein the composition further comprises at least one chemotherapeutic agent selected from the group consisting of taxol, daunorubicin, dacitinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil, floxuridine, methotrexate, colchicine, vincristine, vinlastin, etoposide and cisplatin.

14. A method of treating tumours in a subject, the method comprising administering to the subject a composition according to claim 12 or 13.

15. A method of enhancing the sensitivity of malignant or virus infected cells to therapy, the method comprising modulating expression level of a member of the bcl-2 gene family selected from the group consisting of bcl-2, bcl-xl, bcl-w, bfl-1, brag-1, Mcl-1 and A1 using a DNAzyme according to any one of claims 1 to 12 or a pharmaceutical composition according to claim 12 or 13.

16. A method of treating tumours in a subject, the method comprising administering to the subject a first composition comprising at least one DNAzyme according to any one of claims 1 to 12 and a second composition comprising an anticancer agent.

17. A method as claimed in claim 16 in which the anticancer agent is selected from the group consisting of taxol, daunorubicin, dacitinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil, floxuridine, methotrexate, colchicine, vincristine, vinlastin, etoposide and cisplatin.

1/10




Figure 1. "10-23" DNAzyme and its phosphorothioate modified version (PS-DNAzyme).

2/10




Figure 2. Stability of phosphorothioate-modified DNAzyme oligonucleotides in human serum.

3/10




Figure 3. TMP-Mediated DNAzyme Transfection of PC3 Cells.

4/10




Figure 4. Chemosensitization of PC3 cells to Carboplatin by Bcl-xL DNazyme.

5/10




Figure 5.

Effect of bcl-xL DNAzyme on MDA-MB 231 Breast Cancer Mouse Model





Figure 6.

Effect of bcl-2 DNazyme on MDA-MB 231 Breast Cancer Mouse Model



* p > 0.05 vs Taxol

Figure 7.

Bcl-2 Protein Expression in MDA-MB-231 Tumor




Figure 8.

Effect of bcl-2 DNAzyme on PC3
Prostate Cancer Mouse Model




Figure 9.

Effect of bcl-2 DNzyme on 518A2 Melanoma Mouse Model




Figure 10.

1/23

SEQUENCE LISTING

<110> Johnson & Johnson Research Pty Ltd
 <120> bcl-2 DNazymes
 <130> WJP13107942
 <160> 104
 <170> PatentIn version 3.1

<210> 1
 <211> 15
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> DNzyme catalytic domain
 <400> 1
 ggctagctac aacga

15

<210> 2
 <211> 6030
 <212> DNA
 <213> Homo sapiens
 <300>
 <308> M14745
 <309> 1993-04-27
 <313> (1)..(6030)
 <400> 2

gttggccccc	gttacttttc	ctctgggaaa	tatggcgcac	gctggggagaa	caggggtacga	60
taaccgggag	atagtgatga	agtacatcca	ttataagctg	tcgcagaggg	gctacgagtg	120
ggatgcggga	gatgtgggcg	ccgcgcccc	ggggggccgc	cccgcgccg	gcatcttctc	180
ctcgcagccc	gggcacacgc	cccatacagc	cgcaccccg	gacccggctc	ccaggacctc	240
gccgctgcag	accccggtc	cccccggc	cgcccgggg	cctgcgctca	gcccgggtgc	300
acctgtggtc	cacctgaccc	tccgccagc	cggcgacgac	ttctcccgcc	gctaccgccc	360
cgacttcgcc	gagatgtcca	ggcagctgca	cctgacgccc	ttcaccgcgc	ggggacgctt	420
tgccacgggt	gtggaggagc	cttccagga	cggggtgaac	tgggggagga	ttgtggcctt	480
ctttgagttc	ggtgggggtca	tgtgtgtgga	gagcgtcaac	cgggagatgt	cgccccctgt	540
ggacaacatc	gcccctgtgga	tgactgagta	cctgaaccgg	cacctgcaca	cctggatcca	600
ggataacgga	ggctgggatg	cctttgtgga	actgtacggc	cccagcatgc	ggcctctgtt	660
tgattttctc	tggtctgtct	tgaagactct	gctcagtttg	gccctgggtg	gagcttgcat	720
caccctgggt	gcctatcttg	gccacaagt	aagtcaacat	gcctgcccc	aacaaatatt	780
caaaagggtc	actaaagcag	tagaaataat	atgcattgtc	agtgatgttc	catgaaacaa	840
agctgcaggc	gttttaagaa	aaaataacac	acataataac	atcacacaca	cagacagaca	900
cacacacaca	caacaattaa	cagtcttcag	gcaaaacgtc	gaatcagcta	tttactgcca	960
aagggaata	tcattttatt	tttacattat	taagaaaaaa	agattttatt	atttaagaca	1020
gtcccatcaa	aactcctgtc	tttgaaaatc	cgaccactaa	ttgccaagca	ccgcttcctg	1080
tggtccacc	tggtgttct	gtgcctgtaa	acatagattc	gctttccatg	ttgttgcccg	1140
gatcaccatc	tgaagagcag	acggatggaa	aaaggacctg	atcattgggg	aagctggcct	1200
tctggctgct	ggaggctggg	gagaaggtgt	tcattcactt	gcatttcttt	gccctggggg	1260
ctgtgatatt	aacagaggga	gggttcctgt	ggggggaagt	ccatgcctcc	ctggcctgaa	1320
gaagagactc	tttgcatatg	actcacatga	tgcatacctg	gtgggaggaa	aagagttggg	1380
aacttcagat	ggacctagta	cccactgaga	tttccacgcc	gaaggacagc	gatgggaaaa	1440
atgcccttaa	atcataggaa	agtatttttt	taagctacca	attgtgccga	gaaaagcatt	1500
ttagcaattt	atacaatatt	atccagtacc	ttagccctg	attgtgtata	ttcatatatt	1560
ttggatacgc	accccccaac	tcccaatact	ggctctgtct	gagtaagaaa	cagaatcctc	1620
tggaacttga	ggaagtgaac	atttcggtga	cttccgcac	aggaaggcta	gagttaccca	1680
gagcatcagg	ccgccacaag	tgctgtcttt	taggagaccg	aagtcgcgag	aacctgcctg	1740
tgctccagct	tggaggcctg	gtcctggaac	tgagccgggg	ccctcactgg	cctcctccag	1800
ggatgatcaa	cagggcagtg	tggtctccga	atgtctggaa	gctgatggag	ctcagaattc	1860
cactgtcaag	aaagagcagt	agaggggtgt	ggctgggccc	gtcaccctgg	ggccctccag	1920
gtaggcccg	tttcacgtgg	agcatgggag	ccacgacctt	tcttaagaca	tgtatcactg	1980
tagaggggaag	gaacagaggc	cctgggccc	tcctatcaga	aggacatgg	gaaggctggg	2040
aacgtgagga	gaggcaatgg	ccacggccca	ttttggctgt	agcacatggc	acgttggtct	2100
tgtggccttg	gccacactgt	gagtttaaag	caaggcttta	aatgactttg	gagagggtca	2160

caaatacctaa	aagaagcatt	gaagtggaggt	gtcatggatt	aattgacccc	tgtctatgga	2220
attacatgta	aaacattatc	ttgtcactgt	agtttggttt	tatttgaaaa	cctgacaaaa	2280
aaaaagttcc	agggtgtgaa	tatggggggt	atctgtacat	cctggggcat	taaaaaaaaa	2340
atcaatgggtg	gggaactata	aagaagtaac	aaaagaagtg	acatcttcag	caaataaact	2400
aggaaatttt	ttttctttcc	agtttagaat	cagccttgaa	acattgatgg	aataactctg	2460
tggcattatt	gcattatata	ccatttatct	gtattaactt	tggaatgtac	tctgttcaat	2520
gtttaatgct	gtggttgata	tttcgaaagc	tgctttaaaa	aaatacatgc	atctcagcgt	2580
ttttttgttt	ttaattgtat	ttagttatgg	cctatacact	atthgtgagc	aaaggtgatc	2640
gttttctgtt	tgagattttt	atctcttgat	tcttcaaaaag	cattctgaga	aggtagagata	2700
agccctgagt	ctcagctacc	taagaaaaac	ctggatgtca	ctggccactg	aggagctttg	2760
tttcaaccaa	gtcatgtgca	tttccacgtc	aacagaattg	tttattgtga	cagttatatac	2820
tggtgtccct	ttgacctgtt	ttcttgaagg	tttctctgtc	cctgggcaat	tccgcattta	2880
attcatggta	ttcaggatta	catgcatgtt	tggtttaaacc	catgagattc	attcagttta	2940
aaatccagat	ggcaaatgac	cagcagattc	aaatctatgg	tggtttgacc	tttagagagt	3000
tgctttacgt	ggcctgtttc	aacacagacc	cacccagagc	cctcctgccc	tccttccgcg	3060
ggggctttct	catgctgtgc	cttcagggtc	ttcctgaaat	gcagtgggtgc	ttacgctcca	3120
ccaagaaagc	aggaaacctg	tggtatgaag	ccagacctcc	cgggcgggccc	tcagggaaca	3180
gaatgatcga	acctttgaat	gattctaat	tttaagcaaa	atattatttt	atgaaagggt	3240
tacattgtca	aagtgatgaa	tatggaatat	ccaatcctgt	gctgctatcc	tgccaaaaatc	3300
atthtaattg	agtcagtttg	cagtatgtct	cacgtggtaa	gatcctccaa	gctgcttttag	3360
aagtaacaat	gaagaacgtg	gacgctttta	atataaagcc	tgthttgtct	tctgttggtg	3420
ttcaaacggg	attcacagag	tatttgaaaa	atgtatatat	attaagagggt	cacggggggt	3480
aattgtctgg	tggtctgctt	ttgctgtggg	gtthttgttac	ctggttttaa	taacagtaaa	3540
tgtgcccagc	ctcttgccc	cagaactgta	cagtattgtg	gctgcacttg	ctctaagagt	3600
agttgatgtt	catttttctt	tattgtttaa	aacatgttag	aagcaatgaa	tgatatataa	3660
agcctcaact	agtcattttt	ttctctctct	ctthtttttt	attatatcta	attatttttgc	3720
agttgggcaa	cagagaacca	ttcctatttt	gtattgaaga	gggattcaca	tctgcatctt	3780
aactgtctct	tatgaatgaa	aaaacagctc	tctgtatgta	ctcctcttta	cactggccag	3840
ggtcagagtt	aaatagagta	tatgcacttt	ccaaattggg	gacaagggtc	ctaaaaaaag	3900
cccaaaaagg	agaagaacat	ctgagaacct	ctcggccct	cccagtcctc	cgctgcacaa	3960
atactccgca	agagaggcca	gaatgacagc	tgacagggtc	tatggccatc	gggtcgtctc	4020
cgaagatttg	gcaggggagc	aaaactctgg	caggcttaag	atthtgaata	aagtcacaga	4080
atcaagggaag	cacctcaatt	tagttcaaac	aagacgcca	cattctctcc	acagctcact	4140
tacctctctg	tggtcagatg	tggtcttcca	tttatatgtg	atctttgttt	tattagtaaa	4200
tgcttatcat	ctaaagatgt	agctctggcc	cagtgggaaa	aattaggaag	tgattataaa	4260
tcgagaggag	ttataataat	caagattaaa	tgtaataaat	cagggcaatc	ccaacacatg	4320
tctagctttc	acctccagga	tctattgagt	gaacagaatt	gcaaatagtc	tctatttgta	4380
attgaactta	tcctaaaaca	aatagtttat	aaatgtgaac	ttaaactcta	attaattcca	4440
actgtacttt	taaggcagtg	gctgttttta	gactttctta	tcacttatag	ttagtaatgt	4500
acacctactc	tatcagagaa	aaacaggaaa	ggctcgaaat	acaagccatt	ctaaggaaat	4560
tagggagttc	gttgaaattc	tattctgata	ttattctgtg	gtgtcttttg	cagcccagac	4620
aaatgtgggt	acacactttt	taagaaatac	aattctacat	tgtaagctt	atgaagggtc	4680
caatcagatc	tttatttgta	ttcaatttgg	atctttcagg	gatttttttt	ttaaaattat	4740
atgggacaaa	ggacatttgt	tggaggggtg	ggagggagga	acaattttta	aatataaaac	4800
attcccaagt	ttggatcagg	gagttggaag	ttttcagaat	aaccagaact	aagggtatga	4860
aggacctgta	ttggggtcga	tgtgatgcct	ctgcgaagaa	ccttgtgtga	caaatgagaa	4920
acattttgaa	gtttgtggta	cgacctttag	attccagaga	catcagcatg	gctcaaagtg	4980
cagctccgtt	tggcagtgca	atggtataaa	gttcaagctg	gatatgtcta	tggtgttatg	5040
aaacaataaa	tgtgcagttt	taactaacag	gatatttaat	gacaaccttc	tggttggttg	5100
ggacatctgt	ttctaaatgt	ttattatgta	caatacagaa	aaaaatttta	taaaattaaag	5160
caatgtgaaa	ctgaattgga	gagtgataat	acaagtcctt	tagtcttacc	cagtgaatca	5220
ttctgttcca	tgtctttgga	caacctatgac	cttggaacat	catgaaatat	gcactctcact	5280
ggatgcaaaag	aaaatcagat	ggagcatgaa	tggtactgta	cgggttcac	tggtactgccc	5340
cagaaaaata	acttcaagca	aacatcctat	caacaacaag	gttgttctgc	ataccaagct	5400
gagcacagaa	gatgggaaca	ctggtggagg	atggaaaagg	tcgctcaatc	aagaaaaattc	5460
tgagactatt	aataaataag	actgtagtgt	agatactgag	taaatccatg	cacctaaacc	5520
ttttggaaaa	tctgccgtgg	gccctccaga	tagctcattt	cattaaagttt	ttccctccaa	5580
ggtagaattt	gcagagtgga	cagtggattg	catttctttt	ggggaagctt	tcttttggtg	5640
gtttttgttt	ttataccttc	ttaagttttc	aaccaagggt	tgcttttgtt	ttgagttact	5700
gggtttattt	ttgtttttaa	taaaaataag	tgtacaataa	gtgtttttgt	attgaaagct	5760
tttgttatca	agattttcat	acttttacct	tccatggctc	tttttaagat	tgatactttt	5820
aagaggtggc	tgatattctg	caacactgta	cacataaaaa	atacggtaag	gatactttac	5880
atgggttaagg	taaagtaagt	ctccagttgg	ccaccattag	ctataatggc	actttgtttg	5940
tggtgttgga	aaaagtcaca	ttgccattaa	actttccttg	tctgtctagt	taatatgtgt	6000

3/23

aagaaaaata aagtacagtg tgagatactg

6030

<210> 3
 <211> 926
 <212> DNA
 <213> Homo sapiens
 <300>
 <308> z23115
 <309> 1994-07-26
 <313> (1)..(926)
 <400> 3

gaatctcttt	ctctcccttc	agaatcttat	cttggctttg	gatcttagaa	gagaatcact	60
aaccagagac	gagactcagt	gagtgcagc	gtgttttga	caatggactg	gttgagccca	120
tccctattat	aaaaatgtct	cagagcaacc	gggagctggt	ggttgacttt	ctctcctaca	180
agctttccca	gaaaggatac	agctggagtc	agtttagtga	tgtggaagag	aacaggactg	240
agggcccgga	agggactgaa	tgggagatgg	agaccccgag	tgccatcaat	ggcaacccat	300
cctggcacct	ggcagacagc	cccgcggtga	atggagccac	tgccacagc	agcagtttgg	360
atgcccgga	ggtgatcccc	atggcagcag	ttaaagcaagc	gctgaggag	gcaggcgacg	420
agtttgaact	gcggtaccgg	cgggcattca	gtgacctgac	atcccagctc	cacatcaccc	480
cagggacagc	atatcagagc	tttgaacagg	tagtgaatga	actcttcagg	gatggggtaa	540
actggggtcg	cattgtggcc	tttttctcct	tggcgggggc	actgtgcgtg	gaaagcgtag	600
acaaggagat	gcagggtattg	gtgagtcgga	tgcgagcttg	gatggccact	tacctgaatg	660
accacctaga	gccttggatc	caggagaacg	gcggctggga	tacttttgtg	gaactctatg	720
ggaacaatgc	agcagccgag	agccgaaagg	gccaggaacg	cttcaaccgc	tgggttcctga	780
cgggcatgac	tgtggccggc	gtggttctgc	tgggctcact	cttcagtcgg	aaatgaccag	840
acactgacca	tccactctac	cctcccaccc	ccttctctgc	tccaccacat	cctcgtcca	900
gccgccattg	ccaccaggag	aacccg				926

<210> 4
 <211> 3542
 <212> DNA
 <213> Homo sapiens
 <300>
 <308> NM_004050
 <309> 2001-06-28
 <313> (1)..(3542)
 <400> 4

cccacgcgtc	cgctccctct	ctccctccct	cccagctcct	gcaccaggaa	acggccccga	60
tcccggcagc	ggcctgaccc	ggctccacgc	tggccaggag	gatgaaaggc	cccagctggg	120
ggctccttgc	caccagtgtc	gtgtcttaag	agctgccatc	ccggctggcc	gcccggatgg	180
cgaccccagc	ctcgccccca	gacacacggg	ctctgggtggc	agactttgtg	ggttataagc	240
tgaggcagaa	gggttatgtc	tgtggagctg	gccccgggga	gggcccagca	gctgacccgc	300
tgcaccaagc	catgcgggca	gctggagatg	agttcgagac	ccgcttccgg	cgcaccttct	360
ctgatctggc	ggctcagctg	catgtgaccc	caggctcagc	ccaacaacgc	ttcacccagg	420
tctccgatga	actttttcaa	gggggccccca	actggggccg	ccttgtagcc	ttctttgtct	480
ttggggctgc	actgtgtgct	gagagtgtca	acaaggagat	ggaaccactg	gtgggacaag	540
tgcaggagtg	gatggtggcc	tacctggaga	cgcggctggc	tgactggatc	cacagcagtg	600
ggggctgggc	ggagttcaca	gctctatacg	gggacggggc	cctggaggag	gcgcggcgtc	660
tgcgggaggg	gaactgggca	tcagtgagga	cagtgtgac	gggggccgtg	gcactggggg	720
ccctggtaac	tgtaggggcc	ttttttgcta	gcaagtgaag	gtccagggcc	aggtggggct	780
aggtgtggct	ggggggccagg	agagcaggaa	cagaacagag	aaatgccctt	ggaagaagtg	840
gagttggtgg	atgggtgggc	atggaacagg	atgggcagag	aaagggtagt	gtgtgaggga	900
gctgagttag	ccaggtaggc	gattggaaga	gtgagcagga	cacagagggg	aggggaatgt	960
tttggaagt	ttaggggcac	aggagatgta	gtcgtttccg	ggctggggga	ggtgggaggg	1020
atcacgccta	taggtgtggg	cacatgaaac	gacctggaac	ttgcttcaca	gccctgagga	1080
aggtggactt	acataagcag	ctgtattcca	ttagatgagt	gggatttagg	gaacgcagaa	1140
ggcacatccc	tttggaatgg	aagcttaggg	gttctcaggt	gatagggaga	ggtggctgtt	1200
aacagtgggc	tgcttgga	cgcgtgtgca	tgtgcacgca	tgctggtgtg	catgctgggc	1260
tgcctggcaa	atctggtggt	ggtgggattc	ctcaaggaga	aaacattccc	tcttgcaatg	1320
gcaagaacta	ggggcagttc	tctgtccctc	ctcccaaccc	ctcctttccc	ctgcccttgt	1380
cctgatgcct	caaggcttag	agagaacat	tgtatccaga	ccgagggtc	tgctgcttct	1440
ttccagaaag	tgattggcaa	ggctttggag	agaagagcag	ttctgcagct	ggccttgttc	1500

4/23

```

cttcatcacc ccccttcctt gtgcattatg cacttgctgc tgcctcctgg gctctgatag 1560
aagggcaggg ctgttgagcc tggatgggtg gaggccttagg tagccggacc tgcctgccac 1620
cctcctctcc cactcaggca caatgggtgcc taaagtgttt ccaatctctg ggacctctgt 1680
acccaaactg aaactctaaa ttggggccct aactaatttt ccttttgagg ttgtgggcat 1740
aagtgtgat ctagaataca gtctgggtcc cacactgtgt ctcagtgaga ctgttgatgc 1800
ctttagatga ccatttcaga tctgaatccc atgggtgtga ggggtatggg tactccagga 1860
ctggcctatg ctgtgttggt ggcttttggt cggctttatc aggggccagg catatgggtt 1920
ctagagtacc taccatgacc tagaagcatt tatgatttat ttgaagccac actgtttgca 1980
tgggtgttac ttgtctgtac ctcagagtct gaggatgtta actttggaac tcgcagtcc 2040
ctagaacagc ttcagattat ggctttttct tttgaggaaag aaattattca ctccagatgc 2100
atgccttgag ccagacctca ctgctgcact ttccaagggt ctaagattgc tgcctccaa 2160
tgctaacttt ctgacacagt gctctagaac cctgcctgtg gtcctgagca ctgacagct 2220
tagctagacc atggttgact cttcttgagg attttcaact ggctctagaa tgtggcaacg 2280
tagttgtgct cgccagaacg tgggaccaa ttggcctcag gtgttgagtc cagacttctg 2340
cttttgagag agggctgcac tttttcatgg tatttctagg ggaggtggta ggctgcatgt 2400
gccacttggt cttgttggtga gtatgctgac accagaaact cagagccagc ttgtggcaag 2460
cagttggggg ggggggtctc tgacttgctc aggacaaact aggccagtg ttttcaaact 2520
gcttggcaga gcctgaagt ttcctagggg ttgctcagg agtccttggg gagatgaagg 2580
gggtggggag ctgagcaggg tgggcaattt gccctcaaac agaacagctc cccttgtagc 2640
tgtcttacat attgggggtc agggtaagat tttatttgca ttaaggggtt tgctgctgaa 2700
aaaaagttgg aaaaccactg actagaccat cggctccaaa ttggagtctg tgcttccctc 2760
cccaggtatg gagcacactc ttcaccctac cctctaccac aggacacata tccctgttag 2820
cattccccgg gaccttttagc caagaggagc tgcagggacc atggccaggt taccaaaatg 2880
ccctgctctg aagccttgac acctgggtgg aaagagaggc tgttttctga aagggtaaag 2940
ggcttggtct ggattccagc aagcatagct tagatgggac cacagtgggc aattttgacc 3000
tgtctgccc ttcttagctt gaagggaaac ccagagact cttctgtcag ggaaaactag 3060
ggactctctt ctagagccat atagtctctt gggattagct cttggccaag aaggctgagt 3120
atggttccca atttttaaat ccatttcatt ttttaaaaaa taagggaaat aaatgtaatt 3180
gccatttttc aaagattaag taggaggaga ggggtttctt gctctccaga gcccaaagg 3240
acaaataggg accttgttta ggccaaggaa ggagcggaag tagggcaact cggctctgag 3300
attattaatc ccactcccca cttattctag ggcacacaaa cactatttta cttttttaa 3360
atcataaaac ggcagaacag atttggttag tttagaagaa aagaaagctc tataaatata 3420
aatctatatt cctgtatttt tatttaataa tttataaata ccaagttcat ttgactttta 3480
ttttgtgta atatgtaatg atcgtattaa aaacaataaa taaagcccag aagtttaatt 3540
ag 3542

```

```

<210> 5
<211> 737
<212> DNA
<213> Homo sapiens
<300>
<308> U27467
<309> 1995-11-29
<313> (1)..(737)
<400> 5

```

```

ccagctcaag actttgctct ccaccaggca gaagatgaca gactgtgaat ttggatatat 60
ttacaggctg gctcaggact atctgcagtg cgtcctacag ataccacaac ctggatcagg 120
tccaagcaaa acgtccagag tgctacaaaa tgttgcgttc tcagtccaaa aagaagtggg 180
aaagaatctg aagtcatgct tggacaatgt taatgttggt tccgtagaca ctgccagaac 240
actattcaac caagtgatgg aaaaggagtt tgaagacggc atcattaact ggggaagaat 300
tgtaaccata tttgcatttg aaggtattct catcaagaaa cttctacgac agcaaattgc 360
cccggtatgt gatacctata aggagatttc atattttgtt gcggagttca taatgaataa 420
cacaggagaa tggataaggc aaaacggagg ctgggaaaat ggctttgtta agaagtttga 480
acctaaatct cctcggtatg cttttctaga agttacagga aagatctgtg aaatgctatc 540
tctcctgaag caatactgtt gaccagaaag gacactccat attgtgaaac cggcctaatt 600
tttctgactg atatggaaac gattgccaac acatacttct acttttaaat aaacaacttt 660
gatgatgtaa cttgaccttc cagagttatg gaaattttgt ccccatgtaa tgaataaatt 720
gtatgtattt ttctcta 737

```

```

<210> 6
<211> 6502
<212> DNA

```

5/23

<213> Homo sapiens

<300>

<308> AF147742

<309> 2000-06-09

<313> (1) .. (6502)

<400> 6

ccttgaggac	aggagtgtga	gaccatcctg	gataacatag	caagactttg	ttactttctt	60
tctttttttt	tttttgagac	agagtctcgt	tctgttgccc	aggctggagt	gcagtggcac	120
gatctcggct	cactgaaagc	tctgcctccc	ggattcatgc	cattctcctg	cctcagcctc	180
ctgagtagct	gggactatag	gcacccgcca	ccatgcccag	ctaatttttc	gtattttttt	240
tttagtagag	acgggggttc	accgtgttgg	ccaggatggt	cttgatctcc	tgacctcgtg	300
atctgcccgc	ctcagcctcc	caaagtgcgt	ggattacagg	cgtgagcaac	cgtgcccgcg	360
caagacttgc	ttcctaacaa	acagggccag	ttgcaatggc	tcatgcctat	aatcctagca	420
ctttgggagg	ccaaggaggg	cagatggctt	gaggccagga	gttcgagatt	ggcctggaca	480
acatggtgaa	acccccatct	ctacaaaaaa	acacaaaaat	tagccaggca	tgggtggtgct	540
ggcctgttgt	cccagctact	tgggaagctg	aggtaggagt	atcactttag	ctcaggagggt	600
caaggttgca	gtgagccgag	actgcaccac	tgcactccag	cctgagcaac	atggtgatac	660
ccgtctcaaa	aaataataat	aacaaatgca	gaataaatgc	aattttatct	aaagtgaac	720
ttgcatttcc	tttttttagcc	tctgtacaag	gaaaaatcat	tgctctcctc	atttctcctc	780
tctctttcca	ctttaccacc	tgataaaatt	ttactttata	aagcatgaga	gcaaagctac	840
ctctccata	acactttcct	ctagctctct	cagcccaaag	tgaatttccc	aacctcttaa	900
ctccaaaatg	aagttgttaa	tgccttgtgt	agagcataca	ttccatctca	cattatggtt	960
agttgtctga	caagattaga	cattccttaa	atagagaaac	tattttctat	tactataaac	1020
caaaaaatgc	tctatccttg	ccactcatat	tataaacccc	tatggttcta	ggtcctgccc	1080
aaaaacataa	tggtgtgtat	ggagccgcta	tcactttact	aaactgtgac	attttgggga	1140
ttaggaactt	ttggccaaga	gggagactca	cgcctataat	tccaacactt	tattttattta	1200
tttatttttg	agatggcggt	tcgtctctgt	tgtccaggct	ggagtgcatt	ggcgactctt	1260
cagctcaccg	caacctccgc	ctcccagggt	caagcgattc	tcccgctcca	gccttccaag	1320
tagctgggat	tacaggcacg	tgccaccacg	gcccggctaa	ttttgtatct	ttagtagaga	1380
tggagtttct	ccatggttgg	tgggtgtgtc	tcaaaactct	gacctcagat	gattcgcccc	1440
ccttggccctc	ccaaagtgc	gggattgcag	gtgtgagcca	ctgcccagg	cctcattatt	1500
attattattt	tttttgagac	caagtcttgc	tctgttgccc	aggctggagt	gcagtggcac	1560
tatcttggct	caccgcaacc	tccgtctctt	gggttcaagc	agttctcctg	tctcagcctc	1620
cagagtagct	ggtattacag	atgcgcacca	ccacacccat	ctaatttttg	tgtttttagt	1680
agagacaggg	tttcgccatg	tttcctaggc	tgggtcctaa	ctcctgggct	caagcgatcc	1740
accacctcta	gcctcccaaa	gtgctgggat	tattggcatg	aggcacagag	cccggctgtg	1800
aatcccaaca	ctttgggagg	ccaaggtagg	aggatcacct	gagtcaggga	gttcaagacc	1860
ggcctgggca	aaatagtgat	accccatctc	tacaagaaat	aaaaaaatta	gccaagtata	1920
ggggcatgca	cctgtgttcc	tcgtactctg	cgaggctgtg	gtgggaggat	cacttcagcc	1980
caggagggtt	aggcagcaat	gagcactgat	ggtgccactg	cactccagcc	tggtgtacag	2040
ggcaagacct	catctcaaaa	aaataaataa	aaagttagct	tgctcacctt	tcctatgtct	2100
ctcagcacct	tgcttttgaa	ttttagctat	tatttttaca	gatcttttaa	caaaaaggct	2160
gcttttaatta	acgttaacta	acatacatgg	catataagaa	gacccctgtt	ctcaagggtc	2220
ttacaaacct	ctagagtcaa	atgtgcctta	ttatcagtac	aaaaataaat	ggtgtcagct	2280
gggtgcagtg	actcacacct	gtaatcccag	cactttaaga	ggctgaggca	ggtggatcac	2340
ctgaggccag	gagtttgaga	ccagcctggc	caacatgggt	aaaccacatt	gtcaggcctc	2400
tgagcccaag	ccaagccatc	gcatccctcg	tgacttgcac	gtatacatcc	agatggcctg	2460
aagtaattga	agatcccaaa	aagaagtaaa	aatagcctta	actgatgaca	ttccaccatt	2520
gtgatttgtt	tctgccccac	ccgaactgat	caatgtactt	tgtaatctcc	cccaccctta	2580
agaagggttct	ttgtaattct	ccccaccctt	gagaatgtac	tttgtgagat	ccacccttgc	2640
ccacaaaaca	ttgtctctca	cttcaccacc	tatcccaaaa	cctgtaagaa	ctaataataa	2700
ttcatcaccc	tttgtctgact	ctcttttcgg	actcagcccc	cctgcaccca	ggtgaaataa	2760
acagccatgt	tgtctacaca	aagcctgttt	ggtggtgtct	tcacacagac	gcgcatgaaa	2820
cacatctcta	ctaaaaatac	aataatcagc	tgggcgagggt	ggctcacagc	tgtaactctc	2880
gcactttggg	aggccgagac	aggcaggcca	cttgaggcca	tgagttcgag	accagcctgg	2940
ccaacatcgt	gaaaacccca	tctctaccaa	aaatacaaaa	actagccaga	tgtggtggcg	3000
cacgcttgta	atcccagcta	ctcgggaggc	tgaggtaccg	aatcgtctga	acgtgggaag	3060
tggagcttgt	agttagccga	gatcgcccca	ctgcactcca	gcctgggcaa	cagagctaga	3120
ctgtctcaaa	acaaaacaaa	aatggtgtca	agactctcag	acgagattct	aatggattaa	3180
ggcctatatg	taaatagcac	caaagactat	ggaacagaga	tgggagaagc	aagcaggagg	3240
gcaggaaatg	tttagctgtg	caggttttag	cttagtcac	ttacataaat	ggttctttag	3300
ggtagcacgt	ggagcatcct	catttccaaa	cattggactg	agagtagaga	gctgtgcaaa	3360
ataaccacaa	gtccccaact	atgcctctct	aattatccct	atcatctaag	actgttgttc	3420
ccatccatca	ctgaacttcc	cctgtctctt	ccttcaaccc	ctgtgttagt	caatggttga	3480

6/23

aatthttgatt	tggtataaaaa	cctctggcga	aaaccagcaa	aaagggtca	caaatacaggt	3540
ctcagggaag	cacagaggta	gccacgagaa	ggcccgaggt	gctcatggaa	agagctcgag	3600
cccaggagct	ctgggaggac	cccaggcgct	cggagccgcc	gttacgtaac	cggcactcag	3660
agcctccgaa	gaccggaag	ccccgctcag	gccccggctc	aggccccggc	cccgcccccg	3720
gccccggccc	cgccccggcc	cgcccgggca	gctggtaggt	gccgtgcgca	accctccgga	3780
agctgcgcgc	cctttccctt	tttatgggaa	tacttttttt	aaaaaaaaag	agttcgctgg	3840
cgccaccccc	taggactggc	cgccctaaaa	ccgtgataaa	ggagctgctc	gccacttctc	3900
acttccgctt	ccttccagta	aggagtcggg	gtcttcccca	gttttctcag	ccaggcggcg	3960
gcggcgactg	gcaatgtttg	gcctcaaaa	aaacgcggta	atcggactca	acctctactg	4020
tgggggggcc	ggcttggggg	ccggcagcgg	cgccgccacc	cgcccgggag	ggcgactttt	4080
ggctacggag	aaggaggcct	cgcccgggcg	agagataggg	ggaggggagg	ccggcgcggt	4140
gattggcgga	agcgccggcg	caagccccc	gtccaccctc	acgccagact	cccgagggtg	4200
cgcgccggcg	ccgcccattg	gcgcggaggt	ccccgacgtc	accgcgaccc	cccgagggtc	4260
gcttttcttc	gcgcccaccc	gccgcgcggc	gccgcttgag	gagatggaag	ccccggccgc	4320
tgacgccatc	atgtcgcccg	aagaggagct	ggacgggtac	gagccggagc	ctctcgggaa	4380
gcggccggct	gtcctgccgc	tgctggagtt	ggtcggggaa	tctggttaata	acaccagtac	4440
ggacgggtca	ctaccctcga	cgccgcccgc	agcagaggag	gaggaggacg	agttgtaccg	4500
gcagtcgctg	gagattatct	ctcggtacct	tgggagcag	gccaccggcg	ccaaggacac	4560
aaagccaatg	ggcaggctctg	gggcccaccg	caggaaaggcg	ctggagacct	tacgacgggt	4620
tggggatggc	gtgcagcgca	accacgagac	ggccttccaa	ggtaaggggg	ttcatataatc	4680
gccaaggcct	cactcccttt	tttccatctc	tccccggact	caccgcgcaa	gggtgggttg	4740
gaaaccgaaa	cgagtcagtg	ttgaaacgtg	tctcatccta	ttcctgaagc	cagaatatctc	4800
tgcccatgag	tcattgtttc	cgcccatctt	gattcttttg	gaaatggcag	ctcttgtttca	4860
aagaccggaa	agggtgggat	gtcaatttca	agtgggtgca	acctgagttc	tgtaaatccc	4920
agtagcgatt	ttcccgccgc	gggtggggcag	gcgaatcttg	cgccggttta	gacaaaggag	4980
gccgtgagga	cctgcatgct	tttctttctc	aggcatgctt	cggaaactgg	acatcaaaaa	5040
cgaagacgat	gtgaaatcgt	tgtctcgagt	gatgatccat	gttttcagcg	acggcgtaac	5100
aaactggggc	aggattgtga	ctctcatctt	ttttggtgcc	tttgtggcta	aacacttgaa	5160
gaccataaac	caagaaagct	gcacgcaacc	attagcagaa	agtatcacag	acgttctcgt	5220
aaggacaaaa	cgggactggc	tagttaaaca	aagaggctgg	gtaagtttgc	cttaaggatg	5280
aaaggggcct	ttgagtgga	gtagaatgaa	ggattttttt	tagagagggtg	gggatattcta	5340
aaggttttta	tgacgcacgg	ctgtttgcag	gctctaacta	aaggaccatt	gtttattttga	5400
tggtgattta	agtagtggtg	ccttagagat	agtggtagtg	cggctctgaa	ttgtatcaaa	5460
aatcttggtt	ttctctaggc	aattttttgt	tccaattcag	ttgaatactc	ttcagtggtg	5520
tcaaacatg	aaaaataaag	tcaccagggg	aggatagctg	aaataattcc	taaggcggtg	5580
cctgttttta	tggaagaagat	atgggggtga	gcctgcgttt	taaacaaacc	cagatctgat	5640
gcagatgta	cttaactacg	ttgagaaaaa	ctgatctgcg	caattgaggc	gttactgaaa	5700
tattagggtg	tggaagattg	agaataaggg	ttttcgtctt	ttacctcatg	ggaactcttg	5760
aagtcctttt	gttaggataa	atcctaataa	gaccaagata	gtactgtaaa	atgaagttta	5820
attatcatgg	gtccccgctt	aagaaactga	agaacttatt	ttcttttttt	gcccccgggg	5880
tgaataataa	ttggtttact	attgctttag	ggggaaccc	tagatatattt	aattttacctt	5940
ctctctggat	agtagtggtg	ttaagagagc	agaaacccat	acttgaaaat	gtgcttttct	6000
tttttggttt	ctaggatggg	tttgtggagt	tcttccatgt	agaggacctg	gaagggtggc	6060
tcaggaatgt	gctgctggct	tttgaggtg	ttgctggagt	aggagctggg	ttggcatatc	6120
taataagata	gccttactgt	aagtgcataa	gttgactttt	aaccaaccac	caccaccacc	6180
aaaaccagtt	tatgcagttg	gactccaagc	tgtaacttcc	tagagttgca	ccctagcaac	6240
ctagccagaa	aagcaagtgg	caagaggatt	atggctaaca	agaataaata	catgggaaga	6300
gtgctcccca	ttgattgaag	agtcactgtc	tgaaagaagc	aaagttcagt	ttcagtaaca	6360
aacaaacttt	gtttgggaag	ctatggagga	ggacttttag	atttagtgaa	gatggtaggg	6420
tgaaagact	taatttcctt	gttgagaaca	ggaaagtggc	cagtagccag	gcaagtcata	6480
gaattgatta	ccccccgaat	tc				6502

<210> 7

<211> 33

<212> DNA

<213> artificial sequence

<220>

<223> DT564

<400> 7

cgtgcgccag gctagctaca acgaatttcc cag

33

7/23

<210> 8
<211> 33
<212> DNA
<213> artificial sequence
<220>
<223> DT565
<400> 8
tccccggttag gctagctaca acgacgtacc ctg 33

<210> 9
<211> 33
<212> DNA
<213> artificial sequence
<220>
<223> DT566
<400> 9
tcatacactag gctagctaca acgactcccc gtt 33

<210> 10
<211> 33
<212> DNA
<213> artificial sequence
<220>
<223> DT567
<400> 10
cgcatccccag gctagctaca acgatcgtag ccc 33

<210> 11
<211> 33
<212> DNA
<213> artificial sequence
<220>
<223> DT568
<400> 11
tctcccgag gctagctaca acgacccact cgt 33

<210> 12
<211> 33
<212> DNA
<213> artificial sequence
<220>
<223> DT569
<400> 12
gcgcccacag gctagctaca acgactcccc cat 33

<210> 13
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT570
<400> 13
cggcgcccag gctagctaca acgaatctcc cgc 33

<210> 14
<211> 33
<212> DNA
<213> Artificial Sequence

8/23

<220>
<223> DT571
<400> 14
aggagaagag gctagctaca acgagcccgc cgc 33

<210> 15
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT572
<400> 15
gcggctgtag gctagctaca acgaggggcg tgt 33

<210> 16
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT573
<400> 16
gtcccgggag gctagctaca acgagcggct gta 33

<210> 17
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT574
<400> 17
tcctggcgag gctagctaca acgacgggct ccg 33

<210> 18
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT575
<400> 18
cagggtggcag gctagctaca acgacgggct gag 33

<210> 19
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT576
<400> 19
ggtggaccag gctagctaca acgaaggtgg cac 33

<210> 20
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT577
<400> 20
gcctggacag gctagctaca acgactcggc gaa 33

9/23

<210> 21
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT578
<400> 21
ctgcctggag gctagctaca acgaatctcg gcg 33

<210> 22
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT579
<400> 22
cctccaccag gctagctaca acgacgtggc aaa 33

<210> 23
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT580
<400> 23
gctcctccag gctagctaca acgacaccgt ggc 33

<210> 24
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT581
<400> 24
cccagttcag gctagctaca acgacccgtc cct 33

<210> 25
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT582
<400> 25
aggccacaag gctagctaca acgacctccc cca 33

<210> 26
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT583
<400> 26
agaaggccag gctagctaca acgaaatcct ccc 33

<210> 27
<211> 33

10/23

<212> DNA
<213> Artificial Sequence
<220>
<223> DT584
<400> 27
cacacatgag gctagctaca acgacccacc gaa 33

<210> 28
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT585
<400> 28
ccacacacag gctagctaca acgagacccc acc 33

<210> 29
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT586
<400> 29
ctccacacag gctagctaca acgaatgacc cca 33

<210> 30
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT587
<400> 30
ctctccacag gctagctaca acgaacatga ccc 33

<210> 31
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT588
<400> 31
cccgggtgag gctagctaca acgagctctc cac 33

<210> 32
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT589
<400> 32
ggggcgacag gctagctaca acgactcccg gtt 33

<210> 33
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT590

11/23

<400> 33
caggggcgag gctagctaca acgaatctcc cgg 33

<210> 34
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT591
<400> 34
tggtgtccag gctagctaca acgacagggg cga 33

<210> 35
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT592
<400> 35
acagggcgag gctagctaca acgagttgtc cac 33

<210> 36
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT593
<400> 36
agtcattccag gctagctaca acgaagggcg atg 33

<210> 37
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT594
<400> 37
actcagtcag gctagctaca acgaccacag ggc 33

<210> 38
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT595
<400> 38
tatcctggag gctagctaca acgaccaggt gtg 33

<210> 39
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT596
<400> 39
cctccgttag gctagctaca acgacctgga tcc 33

12/23

<210> 40
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT597
<400> 40
acaaaggcag gctagctaca acgacccagc ctc 33

<210> 41
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT598
<400> 41
ggggccgtag gctagctaca acgaagttcc aca 33

<210> 42
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT599
<400> 42
gaggccgcag gctagctaca acgagctggg gcc 33

<210> 43
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT600
<400> 43
aagctcccag gctagctaca acgacagggc caa 33

<210> 44
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT601
<400> 44
ccagggtgag gctagctaca acgagcaagc tcc 33

<210> 45
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT602
<400> 45
agataggcag gctagctaca acgaccaggg tga 33

<210> 46
<211> 33
<212> DNA
<213> Artificial Sequence

13/23

<220>
<223> DT603
<400> 46
tggcccagag gctagctaca acgaaggcac cca 33

<210> 47
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT604
<400> 47
ttgacttcag gctagctaca acgattgtgg ccc 33

<210> 48
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT605
<400> 48
gggcaggcag gctagctaca acgagttgac ttc 33

<210> 49
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT606
<400> 49
ggagccacag gctagctaca acgagaagcg gtg 33

<210> 50
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT607
<400> 50
ccccaatgag gctagctaca acgacaggtc ctt 33

<210> 51
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT608
<400> 51
agggaggcag gctagctaca acgaggactt ccc 33

<210> 52
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT609
<400> 52
ttcctcccag gctagctaca acgacaggta tgc 33

14/23

<210> 53
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT610
<400> 53
tttttcccag gctagctaca acgacgctgt cct 33

<210> 54
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT611
<400> 54
gcggcctgag gctagctaca acgagctctg ggt 33

<210> 55
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT612
<400> 55
ccctgttgag gctagctaca acgacatccc tgg 33

<210> 56
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT613
<400> 56
tggctcccag gctagctaca acgagctcca cgt 33

<210> 57
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT614
<400> 57
cacagccaag gctagctaca acgagtgcca tgt 33

<210> 58
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT615
<400> 58
accccatag gctagctaca acgatccaca cct 33

<210> 59
<211> 33

15/23

<212> DNA
<213> Artificial Sequence
<220>
<223> DT616
<400> 59
cagggttag gctagctaca acgactcacc ttc 33

<210> 60
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT617
<400> 60
gcccagggag gctagctaca acgagaggaa acc 33

<210> 61
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT618
<400> 61
tgctggtcag gctagctaca acgattgccca tct 33

<210> 62
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT673
<400> 62
aagagttcag gctagctaca acgatcacta cct 33

<210> 63
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT674
<400> 63
tttaccacag gctagctaca acgaccgga aga 33

<210> 64
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT675
<400> 64
cccagtttag gctagctaca acgaccatc ccg 33

<210> 65
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT676

16/23

<400> 65
acaatgcgag gctagctaca acgacccagt tta 33

<210> 66
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT677
<400> 66
aggccacaag gctagctaca acgagcgacc cca 33

<210> 67
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT678
<400> 67
aaaaggccag gctagctaca acgaaatgcg acc 33

<210> 68
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT679
<400> 68
ttccacgcag gctagctaca acgaagtgcc ccg 33

<210> 69
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT680
<400> 69
cgctttccag gctagctaca acgagcacag tgc 33

<210> 70
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT681
<400> 70
ccttgctctag gctagctaca acgagctttc cac 33

<210> 71
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT682
<400> 71
atacctgcag gctagctaca acgactcctt gtc 33

17/23

<210> 72
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT683
<400> 72
tcaccaatag gctagctaca acgactgcat ctc 33

<210> 73
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT684
<400> 73
actcaccaag gctagctaca acgaacctgc atc 33

<210> 74
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT685
<400> 74
tccgactcag gctagctaca acgacaatac ctg 33

<210> 75
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT686
<400> 75
gcgatccgag gctagctaca acgatcacca ata 33

<210> 76
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT687
<400> 76
aagctgcgag gctagctaca acgaccgact cac 33

<210> 77
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT688
<400> 77
aagtggccag gctagctaca acgaccaagc tgc 33

<210> 78
<211> 33
<212> DNA
<213> Artificial Sequence

18/23

<220>
<223> DT689
<400> 78
agggtggtcag gctagctaca acgatcagggt aag 33

<210> 79
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT690
<400> 79
tctcctggag gctagctaca acgaccaagg ctc 33

<210> 80
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT691
<400> 80
acaaaagtag gctagctaca acgacccagc cgc 33

<210> 81
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT692
<400> 81
gtctggtcag gctagctaca acgattccga ctg 33

<210> 82
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT693
<400> 82
tttttataag gctagctaca acgaagggat ggg 33

<210> 83
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT694
<400> 83
acatttttag gctagctaca acgaaatagg gat 33

<210> 84
<211> 33
<212> DNA
<213> Artificial Sequence
<220>
<223> DT695
<400> 84
tctgagacag gctagctaca acgattttat aat 33

19/23

<210> 85
 <211> 33
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> DT696
 <400> 85
 gctctgagag gctagctaca acgaattttt ata 33

<210> 86
 <211> 33
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> DT697
 <400> 86
 agtcaaccag gctagctaca acgacagctc ccg 33

<210> 87
 <211> 33
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> DT698
 <400> 87
 gtggctccag gctagctaca acgatcaccg cgg 33

<210> 88
 <211> 885
 <212> DNA
 <213> Homo sapiens
 <300>
 <308> NM_004049
 <309> 2001-06-28
 <313> (1)..(885)
 <400> 88
 agcctacgca cgaaagtgac tagggaggaa ggatattata aagtgatgca aacagaaatt 60
 ccaccagcct ccatgtatca tcatgtgtca taactcagtc aagctcagtg agcatttctca 120
 gcacattgcc tcaacagctt caaggtgagc cagctcaaga ctttgctctc caccaggcag 180
 aagatgacag actgtgaatt tggatatatt tacaggctgg ctcaggacta tctgcagtgc 240
 gtcctacaga taccacaacc tggatcaggt ccaagcaaaa cgtccagagt gctacaaaat 300
 gttgctgttct cagtccaaaa agaagtggaa aagaatctga agtcatgctt ggacaatgtt 360
 aatgtttgtgt ccgtagacac tgccagaaca ctattcaacc aagtgatgga aaaggagtgt 420
 gaagacggca tcattaactg gggaagaatt gtaaccatat ttgcatttga aggtattctc 480
 atcaagaaac ttctacgaca gcaaattgcc ccggtgtggg atacctataa ggagatttca 540
 tattttgttg cggagtcat aatgaataac acaggagaat ggataaggca aaacggaggc 600
 tgggaaaatg gctttgtaaa gaagtttgaa cctaaatctg gctggatgac ttttctagaa 660
 gttacaggaa agatctgtga aatgctatct ctctgaagc aatactgttg accagaaagg 720
 acactccata ttgtgaaacc ggcctaattt ttctgactga tatggaaacg attgccaaaca 780
 catacttcta cttttaaata aacaactttg atgatgtaac ttgaccttcc agagttatgg 840
 aaatthttgtc cccatgtaat gaataaattg tatgtatttt tctct 885

<210> 89
 <211> 960
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature

20/23

```

<222> (279)..(279)
<223> n
<220>
<221> misc_feature
<222> (797)..(797)
<223> n
<220>
<221> misc_feature
<222> (858)..(858)
<223> n
<300>
<308> S82185
<309> 1998-05-26
<313> (1)..(960)
<400> 89
gaactggcgg tggatccctc taacgctggc accacgggac ttcttgatct aaccacccgt 60
gactggaaac ctgcattgct ggatatggct ggcctacgtg ccgatattct ttctcctgtc 120
aaagaaaccg gcacattgct gggcgtggta agttcacaag cggcgggaact ctgcggtctg 180
aaggcgggca ctccggtggt cggtggagga ggcgacgtgc agcttggttg ccttgggtta 240
ggcgttggtg gtccggcaca aaccgcggtt cttggcgga cattctggca gcaagttgta 300
aatttagccg cgccggtgac agaccagaa atgaacgtgc gcgttaatcc tcatgttatt 360
cctggcatgg tacaagctga atctataagc ttttttaccg gactcaccat gcgctgggtc 420
cgcatgctt tctgtgccga agaaaaactg attgccgaac gtttaggcat cgacacctat 480
acgctgctgg aagagatggc cagtcgggtg ccgcctgggt cgtggggcgt aatgccgatc 540
ttctccgaca gaatgcgctt taaaacctgg tatcacgctg cgccttcctt tattaacttg 600
tccattgacc cggataaatg taacaaagcg acattgttcc gtgcgctgga agaaaatgcg 660
gcgattgtat cagcgtgtaa cttgcagcaa attgctgatt tctcgaatat tcatccttca 720
tcgctagtct ttgcaggcgg aggttcaaaa gggaaattat ggagtcaaat tctcgctgat 780
gtctcgggat taccgcncaa tattccggtg gtcaaagaag cactgcatt aggatgtgcc 840
attgcagctg gcgtcgngc cggaattttt tcatcaatgg cagaaaccgg agaacgcctg 900
gttcgctggg aacggacgca cacaaccaga cccggaaaag catgaacttt atcaggattc 960

<210> 90
<211> 22
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 90
cacagcatta aacattgaac ag 22

<210> 91
<211> 21
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 91
tggaactttt tttttgtcag g 21

<210> 92
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 92
tcctcacgtt cccagccttc 20

<210> 93
<211> 20

```

21/23

<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 93
cagacattcg gagaccacac 20

<210> 94
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 94
cagtattggg agttgggggg 20

<210> 95
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 95
ccaactctttt tcctcccacc 20

<210> 96
<211> 21
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 96
cgacgttttg cctgaagact g 21

<210> 97
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 97
cagggccaaa ctgagcagag 20

<210> 98
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 98
atcctcccc agttcacccc 20

<210> 99
<211> 18
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer

22/23

<400> 99
ggatgcggct gtatgggg 18

<210> 100
<211> 21
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-2 cleavage detection primer
<400> 100
aggccacgta aagcaactct c 21

<210> 101
<211> 18
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-xL cleavage detection primer
<400> 101
cgggttctcc tggtagga 18

<210> 102
<211> 18
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-xL cleavage detection primer
<400> 102
cctttcggct ctcggctg 18

<210> 103
<211> 18
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-xL cleavage detection primer
<400> 103
ccgccgaagg agaaaaag 18

<210> 104
<211> 21
<212> DNA
<213> Artificial Sequence
<220>
<223> bcl-xL cleavage detection primer
<400> 104
gcctcagtc tgttctcttc c 21

<210> 105
<211> 18
<212> DNA
<213> Artificial Sequence
<220>
<223> antisense oligonucleotide
<400> 105
gtgccggggt cttcgggc 18

23/23

<210> 106
<211> 15
<212> DNA
<213> Artificial Sequence
<220>
<223> antisense oligonucleotide
<400> 106
gctttgcat ttctg

15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU02/00739

A. CLASSIFICATION OF SUBJECT MATTER		
Int. Cl. ⁷ : C12N 9/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
SEE ELECTRONIC DATABASES		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SEE ELECTRONIC DATABASES		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
WPIDS CA MEDLINE: DNazyme, catalytic dna, Bcl 2, Bcl xl, Bcl w, Bfl 1, Brag 1 Mcl 1, ribozyme, antisense		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00/20432 A1 (ISIS PHARMACEUTICALS INC) 13 April 2000. Particularly pp 1-6, 24 and 30.	1-3, 10-17
X	US 6 001 992 A (AKERMAN EJ et al) 14 December 1999. Particularly pp 4 and 16	1, 2, 10-17
X	WO 97/35971 A1 (AMRAD OPERATIONS PTY LTD 2 October 1997. Particularly pp 2-6, 19	1-3, 10-12, 14-16
Y		13, 17
X	WO 95/13292 A1 (LA JOLLA CANCER RESEARCH FOUNDATION) 18 May 1995 Particularly pp 1-3, 15, 22-23, 27-28	1, 2, 10-17
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>		
Date of the actual completion of the international search 27 June 2002		Date of mailing of the international search report 16 JUL 2002
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustalia.gov.au Facsimile No. (02) 6285 3929		Authorized officer Gillian Allen Telephone No : (02) 6283 2266

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU02/00739

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Jen K-Y et al. Suppression of DNA expression by targeted disruption of messenger RNA: available options and current strategies. Stem Cells. 2000. 18: 307-319.	All
Y	Khachigian LM. Catalytic DNAs as potential therapeutic agents and sequence-specific molecular tools to dissect biological function. J Clinical Investigation. 2000. 106(10): 1189-1195	All
Y	Cairns MJ et al. Target site selection for an RNA-cleaving catalytic DNA. Nature Biotech. 1999. 17:480-486. Whole document	All
X	Olie RA et al. Targeting tumor cell resistance to apoptosis induction with antisense oligonucleotides: progress and therapeutic potential. Drug Resist Updat. Feb 2001. 4(1): 9-15. Review. Whole document.	1-3, 10-17
Y		4-9
X	Miyake H et al. Novel therapeutic strategy for advanced prostate cancer using antisense oligodeoxynucleotides targeting anti-apoptotic genes upregulated after androgen withdrawal to delay androgen-independent progression and enhance chemosensitivity. Int J Urol. Jul 2001. 8(7): 337-49. Review. Whole document	1-3, 10-17
Y		4-9
X	Gibson SA et al. Induction of apoptosis in oral cancer cells by an anti-bcl-2 ribozyme delivered by an adenovirus vector. Clinical Cancer Research. 2000. 6(1): 213-22	1-3, 10-17
Y	Whole document	4-9
X	Dorai T et al. A recombinant defective adenoviral agent expressing anti-bcl-2 ribozyme promotes apoptosis of bcl-2-expressing human prostate cancer cells.; International Journal of Cancer. 1999. 82(6): 846-52. Whole Document.	1-3, 10-17
Y		4-9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/AU02/00739

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report	Patent Family Member
WO 95/13292	US 5539094 AU 11742/95 CA 2176378 EP 742793 JP 9509304T
WO 97/35971	CA 2250207 EP 0932674 JP 2000513566T
WO 00/20432	AU 62710/99 EP 1119579
WO0040595	AU 21807/00 EP 1140971

END OF ANNEX