

Physics 122

Thermal Physics

Lecture Notes

Outline

- 1 Adiabatic Processes
- 2 Adiabatic vs Isothermal
- 3 Thermal Expansion
- 4 Worked Example

What is an Adiabatic Process?

An **adiabatic process** is a thermodynamic process in which

$$Q = 0$$

(no heat exchange with the surroundings).

What is an Adiabatic Process?

An **adiabatic process** is a thermodynamic process in which

$$Q = 0$$

(no heat exchange with the surroundings).

Occurs when:

- Process is very fast
- System is thermally insulated

First Law of Thermodynamics

Physics 122 sign convention:

Work done *by the environment on the system* is positive.

$$\Delta E = Q + W$$

First Law of Thermodynamics

Physics 122 sign convention:

Work done *by the environment on the system* is positive.

$$\Delta E = Q + W$$

For an adiabatic process:

$$\Delta E = W$$

- **Compression:**

$$W > 0 \Rightarrow \Delta E > 0 \Rightarrow T \uparrow$$

Physical Interpretation

- **Compression:**

$$W > 0 \Rightarrow \Delta E > 0 \Rightarrow T \uparrow$$

- **Expansion:**

$$W < 0 \Rightarrow \Delta E < 0 \Rightarrow T \downarrow$$

Physical Interpretation

- **Compression:**

$$W > 0 \Rightarrow \Delta E > 0 \Rightarrow T \uparrow$$

- **Expansion:**

$$W < 0 \Rightarrow \Delta E < 0 \Rightarrow T \downarrow$$

Examples:

- Diesel engine compression
- Opening a soda bottle
- Spraying deodorant

Ideal Gas Processes

$$PV = nRT$$

$$PV = nRT$$

Isothermal:

$$T = \text{constant}, \quad P = \frac{nRT}{V}$$

Ideal Gas Processes

$$PV = nRT$$


Isothermal:

$$T = \text{constant}, \quad P = \frac{nRT}{V}$$


Adiabatic:

$$\boxed{PV^\gamma = \text{constant}} \quad \gamma = \frac{C_P}{C_V}$$

P-V Diagram



P-V Diagram



Adiabatic curves are steeper.

Linear Expansion

$$\Delta L = \alpha L_0 \Delta T$$

Linear Expansion

$$\Delta L = \alpha L_0 \Delta T$$

- α = coefficient of linear expansion
- Units: K^{-1}
- Depends on material

Volume Expansion

$$\Delta V = \beta V_0 \Delta T$$


Volume Expansion

$$\Delta V = \beta V_0 \Delta T$$


For isotropic solids:

$$\beta \approx 3\alpha$$

Microscopic Picture



Microscopic Picture



Higher temperature \Rightarrow larger atomic vibrations \Rightarrow expansion.

Linear Expansion Example

Given:

$$L_0 = 10 \text{ m}, \quad \alpha = 12 \times 10^{-6} \text{ K}^{-1}$$

$$\Delta T = 40^\circ\text{C}$$

Linear Expansion Example

Given:

$$L_0 = 10 \text{ m}, \quad \alpha = 12 \times 10^{-6} \text{ K}^{-1}$$

$$\Delta T = 40^\circ\text{C}$$

Solution:

$$\Delta L = \alpha L_0 \Delta T$$

$$\boxed{\Delta L = 4.8 \text{ mm}}$$

Key Takeaways

- Adiabatic process: $Q = 0$
- Physics 122 convention: $\Delta E = Q + W$
- Compression heats, expansion cools
- Thermal expansion scales with ΔT