Tutorial 7

COMP 5361: Discrete Structures and Formal Languages

Mohammad Reza Davari

Concordia University



Mohammad Reza Davari 1 / 26

Outline

Relations and Their Properties

Closures of Relations

Sequivalence Relations



Contents of the section

Relations and Their Properties

Closures of Relations

3 Equivalence Relations



Definition

• **Binary Relation**: Let A and B be sets. A binary relation from A to B is a subset of $A \times B$.



Definition

- **Binary Relation**: Let A and B be sets. A binary relation from A to B is a subset of $A \times B$.
- Relation on a Set: A relation on a set A is a relation from A to A.



Mohammad Reza Davari 4 / 26

Example

Example

- **2** $R = \{(a, b) | a \le b\}$



Example

- $R = \{(a, b) | a \leq b\}$



Example

- $R = \{(a, b) | a \le b\}$
- $R = \{(a, b)|a > b\}$



Example

Let A be the set $\{1, 2, 3, 4\}$. Which ordered pairs are in the relation,

- $R = \{(a, b) | a \le b\}$
- $R = \{(a, b)|a > b\}$



Mohammad Reza Davari 5 / 26

Definition

• **Reflexive**: A relation R on a set A is called reflexive if $(a, a) \in R$ for every element $a \in A$.



Definition

- **Reflexive**: A relation R on a set A is called reflexive if $(a, a) \in R$ for every element $a \in A$.
- **Symmetric**: A relation R on a set A is called symmetric if $(b, a) \in R$ whenever $(a, b) \in R$, for all $a, b \in A$.



Definition

- **Reflexive**: A relation R on a set A is called reflexive if $(a, a) \in R$ for every element $a \in A$.
- **Symmetric**: A relation R on a set A is called symmetric if $(b, a) \in R$ whenever $(a, b) \in R$, for all $a, b \in A$.
- Antisymmetric: A relation R on a set A such that for all $a, b \in A$, if $(a, b) \in R$ and $(b, a) \in R$, then a = b is called antisymmetric.
- **Transitive**: A relation R on a set A is called transitive if whenever $(a,b) \in R$ and $(b,c) \in R$, then $(a,c) \in R$, for all $a,b,c \in A$.



Mohammad Reza Davari 6 / 26

Example

• Which of the relations on slide 5 are reflexive?



Example

- Which of the relations on slide 5 are reflexive?
- Which of the relations on slide 5 are symmetric?



Example

- Which of the relations on slide 5 are reflexive?
- Which of the relations on slide 5 are symmetric?
- Which of the relations on slide 5 are antisymmetric?



Example

- Which of the relations on slide 5 are reflexive?
- Which of the relations on slide 5 are symmetric?
- Which of the relations on slide 5 are antisymmetric?
- Which of the relations on slide 5 are transitive?



Mohammad Reza Davari 7 / 26

Example

lacktriangle How many relations are there on a set of n elements?



Example

- How many relations are there on a set of *n* elements?
- ② How many reflexive relations are there on a set of n elements?



Example

- **1** How many relations are there on a set of *n* elements?
- ② How many reflexive relations are there on a set of n elements?
- Mow many symmetric relations are there on a set of n elements?



Example

- \bullet How many relations are there on a set of n elements?
- ② How many reflexive relations are there on a set of *n* elements?
- How many symmetric relations are there on a set of n elements?
- \bullet How many antisymmetric relations are there on a set of n elements?



Mohammad Reza Davari 8 / 26

Combining Relations

Definition

- Let R be a relation from a set A to a set B and S a relation from B to a set C. The composite of R and S is the relation consisting of ordered pairs (a, c), where $a \in A$, $c \in C$, and for which there exists an element $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$. We denote the composite of R and S by $S \circ R$.
- Let R be a relation on the set A. The powers R^n , $n=1,2,3,\ldots$, are defined recursively by:

$$R^{1} = R \tag{1}$$

$$R^{n+1} = R^{n} \circ R \tag{2}$$

$$R^{n+1} = R^n \circ R \tag{2}$$



Mohammad Reza Davari

Combining Relations

Theorem

The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \ldots$

Example

① Let R = (1,1), (2,1), (3,2), (4,3). Find the powers R^n , $n = 2, 3, 4, \ldots$



Combining Relations

Theorem

The relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Example

- **1** Let R = (1,1), (2,1), (3,2), (4,3). Find the powers R^n , $n = 2, 3, 4, \dots$
- 2 Let R be a relation that is reflexive and transitive. Prove that $R^n = R$ for all positive integers n.



Mohammad Reza Davari

Contents of the section

Relations and Their Properties

2 Closures of Relations

3 Equivalence Relations



Definition

Let R be a relation on set A, then:

• **Reflexive Closure**: The reflexive closure of R is the smallest relation that contains R and is reflexive.



Mohammad Reza Davari 12 / 26

Definition

Let R be a relation on set A, then:

- **Reflexive Closure**: The reflexive closure of *R* is the smallest relation that contains *R* and is reflexive.
- **Symmetric Closure**: The symmetric closure of *R* is the smallest relation that contains *R* and is symmetric.



Mohammad Reza Davari 12 / 26

Definition

Let R be a relation on set A, then:

- Reflexive Closure: The reflexive closure of R is the smallest relation that contains R and is reflexive.
- **Symmetric Closure**: The symmetric closure of R is the smallest relation that contains R and is symmetric.
- Transitive Closure: The transitive closure of R is the smallest relation that contains R and is transitive.



Mohammad Reza Davari 12 / 26

Example

① What is the reflexive closure of the relation $R = \{(a, b) | a < b\}$ on the set of integers?



Example

- ① What is the reflexive closure of the relation $R = \{(a, b) | a < b\}$ on the set of integers?
- ② What is the symmetric closure of the relation $R = \{(a, b) | a > b\}$ on the set of positive integers?



Mohammad Reza Davari

Example

- What is the reflexive closure of the relation $R = \{(a, b) | a < b\}$ on the set of integers?
- ② What is the symmetric closure of the relation $R = \{(a, b) | a > b\}$ on the set of positive integers?
- Find the transitive closures of the relation R on a, b, c, d, e, where R is given by:

$$\{(a,c),(b,d),(c,a),(d,b),(e,d)\}$$



Mohammad Reza Davari 13 / 26

Definition

• Let R be a relation on a set A. The connectivity relation R^* is defined as:

$$R^* = \bigcup_{n=1}^{\infty} R^n$$



Definition

• Let R be a relation on a set A. The connectivity relation R^* is defined as:

$$R^* = \bigcup_{n=1}^{\infty} R^n$$

• The transitive closure of a relation R equals the connectivity relation R^* .



Example

Example 1

Let R be the relation on the set $\{1, 2, 3, 4, 5\}$ containing the ordered pairs (1,3), (2,4), (3,1), (3,5), (4,3), (5,1), (5,2), and (5,4). Find:

- $\mathbf{0}$ R^2
- $Q R^3$
- \circ R^4
- \bullet R^5
- \circ R^6
- R*



Mohammad Reza Davari 15 / 26

Example

Example 2

Find the transitive closures of these relations on $\{1, 2, 3, 4\}$.

- (1,2),(2,1),(2,3),(3,4),(4,1)
- **4** {(1,2), (1,3), (1,4), (2,3), (2,4), (3,4)}
- (1,1),(1,4),(2,1),(2,3),(3,1),(3,2),(3,4),(4,2)



Mohammad Reza Davari 16 / 26

Contents of the section

Relations and Their Properties

Closures of Relations

3 Equivalence Relations



Definition

- **Equivalence Relation**: A relation on a set *A* is called an equivalence relation if it is reflexive, symmetric, and transitive.
- **Equivalent**: Two elements a and b that are related by an equivalence relation are called equivalent. The notation $a \sim b$ is often used to denote that a and b are equivalent elements with respect to a particular equivalence relation.



Mohammad Reza Davari 18 / 26

Example

① Let R be the relation on the set of integers such that aRb if and only if a = b or a = -b. Is R an equivalence relation?



Example

- **1** Let R be the relation on the set of integers such that aRb if and only if a = b or a = -b. Is R an equivalence relation?
- 2 Let R be the relation on the set of real numbers such that aRb if and only if a-b is an integer. Is R an equivalence relation?



Mohammad Reza Davari 19 / 26

Example

- **1** Let R be the relation on the set of integers such that aRb if and only if a = b or a = -b. Is R an equivalence relation?
- 2 Let R be the relation on the set of real numbers such that aRb if and only if a-b is an integer. Is R an equivalence relation?
- ② Let m be an integer with m > 1. Show that the relation (Congruence Modulo m):

$$R = \{(a, b) | a \equiv b(modm)\}$$

is an equivalence relation on the set of integers.



Mohammad Reza Davari 19 / 26

Example

- Let R be the relation on the set of integers such that aRb if and only if a = b or a = -b. Is R an equivalence relation?
- Let R be the relation on the set of real numbers such that aRb if and only if a - b is an integer. Is R an equivalence relation?
- \bullet Let m be an integer with m > 1. Show that the relation (Congruence Modulo m):

$$R = \{(a, b) | a \equiv b(modm)\}$$

is an equivalence relation on the set of integers.

Show that the "divides" relation is the set of positive integers in not an equivalence relation.



Mohammad Reza Davari

Definition

- Let R be an equivalence relation on a set A. The set of all elements that are related to an element a of A is called the equivalence class of a. The equivalence class of a with respect to R is denoted by [a]_R. When only one relation is under consideration, we can delete the subscript R and write [a] for this equivalence class.
- In the above definition *a* is the representative of this equivalence class.

Properties

If $a \sim b$ then the followings are equivalent:

- aRb
- ② [a] = [b]



Mohammad Reza Davari 20 / 26

Example

• Let R be the relation on the set of integers such that aRb if and only if a = b or a = -b. What is the equivalence class of an integer?



Mohammad Reza Davari 21 / 26

Example

- Let R be the relation on the set of integers such that aRb if and only if a = b or a = -b. What is the equivalence class of an integer?
- 2 What are the equivalence classes of 0 and 1 for congruence modulo 7?



Mohammad Reza Davari 21 / 26

Example

- Let R be the relation on the set of integers such that aRb if and only if a = b or a = -b. What is the equivalence class of an integer?
- ② What are the equivalence classes of 0 and 1 for congruence modulo 7?
- 3 What are the equivalence classes of 5 and 2 for congruence modulo 5?



Mohammad Reza Davari 21 / 26

Definition

• **Partition**: A partition of a set *S* is a collection of disjoint nonempty subsets of *S* that have *S* as their union.



Mohammad Reza Davari 22 / 26

Definition

- **Partition**: A partition of a set S is a collection of disjoint nonempty subsets of S that have S as their union.
- Let R be an equivalence relation on a set S. Then the equivalence classes of R form a partition of S. Conversely, given a partition $\{A_i|i\in I\}$ of the set S, there is an equivalence relation R that has the sets A_i , $i \in I$, as its equivalence classes.



Mohammad Reza Davari

Example 1

Suppose that $S = \{1, 2, 3, 4, 5, 6\}$. Give a partition of set S with 3 classes.



Mohammad Reza Davari 23 / 26

Example 2

List the ordered pairs in the equivalence relation R produced by the partition $A_1 = \{1, 2, 3\}$, $A_2 = \{4, 5\}$, and $A_3 = \{6\}$ of $S = \{1, 2, 3, 4, 5, 6\}.$



Example 3

What are the sets in the partition of the integers arising from congruence modulo 7?



Mohammad Reza Davari 25 / 26

Example 4

What is the congruence class $[4]_m$ when m is:

- **1** 2?
- 2 3?
- **3** 6?
- **4** 8?

