

Hot Pair Coulomb Density Matrix on a Sphere

Riccardo Fantoni*

Università di Trieste, Dipartimento di Fisica, strada Costiera 11, 34151 Grignano (Trieste), Italy

(Dated: May 29, 2026)

We tabulate the hot diagonal pair Coulomb density matrix on a sphere.

Keywords: Pair Coulomb Density Matrix; Sphere; Table

CONTENTS

I. Introduction	1
II. Lower temperature pair Coulomb density matrix	2
III. The code	4
A. $\rho_{i0}(\mathbf{0}, \mathbf{0}; t)$	7
Author declarations	7
Conflicts of interest	7
Data availability	7
Funding	7
References	7

I. INTRODUCTION

In a quantum many body (Monte Carlo) simulation of a Coulomb liquid it is often useful to have access to the low temperature pair Coulomb density matrix ρ . Within Path Integral Monte Carlo (PIMC) [1] at sufficiently high temperature this can be obtained from the so called *primitive approximation* $\rho_{\text{primitive}}$. In order to access the quantum regime it is important to lower the temperature. Starting from $\rho_{\text{primitive}}$ at a high temperature T_0 we can halve the temperature simply by squaring it. Already this lower temperature pair density matrix possesses different properties than the primitive approximation. In position representation, on the diagonal

$$\left| \frac{\phi(\mathbf{r}_1, \mathbf{r}_2)}{T_0} \right| = \left| \ln \left[\frac{\rho_{\text{primitive}}(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_1, \mathbf{r}_2; 1/T_0)}{\rho_0(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_1, \mathbf{r}_2; 1/T_0)} \right] \right| \xrightarrow{\mathbf{r}_1 \rightarrow \mathbf{r}_2} \infty, \quad (1.1)$$

where $\phi(\mathbf{r}_1, \mathbf{r}_2)$ is the pair Coulomb potential between particle 1 at $\mathbf{r}_1$ and particle 2 at $\mathbf{r}_2$. But for the squared matrix

$$|P(\mathbf{r}_1, \mathbf{r}_2; 2/T_0)| = \left| \ln \left[\frac{\rho_{\text{primitive}}^2(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_1, \mathbf{r}_2; 2/T_0)}{\rho_0^2(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_1, \mathbf{r}_2; 2/T_0)} \right] \right| \xrightarrow{\mathbf{r}_1 \rightarrow \mathbf{r}_2} \text{finite}. \quad (1.2)$$

The same contact value smearing property is shared by all the $\rho_{\text{primitive}}^n$ at a temperature n/T_0 for $n > 1$. This feature makes it of fundamental importance not to use the primitive approximation for example in an electron proton binary mixture where the divergence of the Coulomb potential between unlike species at contact would inevitably let the fast electrons to stick onto the slow protons. In this sense the use of a lower temperature pair Coulomb density matrix can be considered as an essential feature in a PIMC computer experiment.

In this work we will show how this can be accomplished on a Sphere. We will not give the exact solution for the pair Coulomb density matrix as was done in flat space in Refs. [2–4] but we will give an approximate numerical solution instead.

* riccardo.fantoni@scuola.istruzione.it

II. LOWER TEMPERATURE PAIR COULOMB DENSITY MATRIX

We show a method to lower the temperature of the hot Pair Coulomb Density Matrix (PCDM) on a Sphere so to extend the work in flat space of Refs. [2–4] to a curved surface.

We will denote with a Greek index a coordinate component and with a Roman index a particle label. We will adopt Einstein summation convention to tacitly assume a sum over repeated Greek indexes. We distinguish between upper contravariant Greek indexes and lower covariant Greek indexes. Passing between the two is accomplished with a contraction with the metric tensor, as usual.

The metric tensor of the Sphere of radius a is

$$||g_{\alpha\beta}|| = a^2 \begin{pmatrix} 1 & 0 \\ 0 & \cos^2 \theta \end{pmatrix}. \quad (2.1)$$

with $g(\mathbf{r}) = \det||g_{\alpha\beta}|| = a^4 \cos^2 \theta$ and $\theta \in [-\pi/2, \pi/2]$ the polar angle measured from the equatorial plane, so that $\theta = \pi/2$ is the north pole and $\theta = -\pi/2$ is the south pole.

It is important to realize that on a curved surface the two body problem is generally not separable into a center of mass and a relative one body problem. Therefore, in full generality, we will study the position representation $\rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t)$ of the density matrix $\rho = \exp(-Ht)$, where H is the Hamiltonian operator for the two bodies and t is the imaginary time that in statistical physics is identified with the inverse temperature $\beta = 1/k_B T$ with k_B Boltzmann constant. These are initially at $\mathbf{r}_1$ and $\mathbf{r}_2$ and at an imaginary time t later they are at $\mathbf{r}'_1$ and $\mathbf{r}'_2$.

Therefore, the PCDM on the Sphere will satisfy the Bloch equation

$$\frac{\partial \rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t)}{\partial t} = \left(\frac{\lambda_1}{\sqrt{g(\mathbf{r}_1)}} \nabla_{1\alpha} \sqrt{g(\mathbf{r}_1)} \nabla_1^\alpha + \frac{\lambda_2}{\sqrt{g(\mathbf{r}_2)}} \nabla_{2\alpha} \sqrt{g(\mathbf{r}_2)} \nabla_2^\alpha - \phi(\mathbf{r}_1, \mathbf{r}_2) \right) \rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t), \quad (2.2)$$

$$\rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; 0) = \delta(\mathbf{r}_1 - \mathbf{r}'_1) \delta(\mathbf{r}_2 - \mathbf{r}'_2) / \sqrt{g(\mathbf{r}_1) g(\mathbf{r}_2)}, \quad (2.3)$$

where, for particle $i = 1, 2$, $\lambda_i = \hbar^2/2m_i$ with m_i the mass of particle i , $g(\mathbf{r}_i) = a^4 \sin^2 \theta_i$, $\nabla_i = (\nabla_{i1}, \nabla_{i2}) = (\partial/\partial\theta_i, \partial/\partial\varphi_i)$ is the gradient respect to the coordinates $\mathbf{r}_i = (r_i^1, r_i^2) = (\theta_i, \varphi_i)$, δ is a two dimensional Dirac delta function, ρ is the pair density matrix, and ϕ is the Coulomb pair potential.

The Coulomb pair potential for particles moving **on** the surface can be chosen as

$$\phi_{\text{on}}(\mathbf{r}_1, \mathbf{r}_2) = Z_1 Z_2 e^2 / d(\mathbf{r}_1, \mathbf{r}_2), \quad (2.4)$$

where $Z_i e$ is the charge of particle i and for the distance between the two points $\mathbf{r}_1$ and $\mathbf{r}_2$, $d(\mathbf{r}_1, \mathbf{r}_2)$, on the Sphere we can either choose the geodesic distance

$$d_g(\mathbf{r}_1, \mathbf{r}_2) = a \arccos[\sin \theta_1 \sin \theta_2 + \cos \theta_1 \cos \theta_2 \cos(\varphi_1 - \varphi_2)], \quad (2.5)$$

or, viewing the Sphere as embedded in the three dimensional space, the Euclidean distance

$$d_e(\mathbf{r}_1, \mathbf{r}_2) = 2a \sin(d_g/2a). \quad (2.6)$$

For particles living **in** the surface [5] we may choose

$$\phi_{\text{in}}(\mathbf{r}_1, \mathbf{r}_2) = -Z_1 Z_2 e^2 \ln[d_e(\mathbf{r}_1, \mathbf{r}_2)/\ell], \quad (2.7)$$

with ℓ a constant with the dimensions of a distance. We will call ϕ_{on}^g , ϕ_{on}^e , ϕ_{in} the three cases just illustrated.

We will measure energy in units of the Hartree e^2/a_B , with e the charge of the electron and $a_B = \hbar^2/m_e e^2$ Bohr radius, with m_e the mass of the electron. And length in units of the Bohr radius a_B . We will call $\mu_i = m_i/m_e$ the reduced mass of particle i .

The free particles, $\phi \rightarrow 0$, pair density matrix can be determined from the work of Bastianelli and Corradini [6]. We will call their solution for the single free particle i , $\rho_{i0}(\mathbf{r}_i | \mathbf{r}'_i; t)$. Therefore the pair free density matrix will be

$$\rho_0(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t) = \rho_{10}(\mathbf{r}_1 | \mathbf{r}'_1; t) \rho_{20}(\mathbf{r}_2 | \mathbf{r}'_2; t). \quad (2.8)$$

Then our PCDM can be written as

$$\rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t) = \rho_0(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t) e^{-P(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t)}, \quad (2.9)$$

where P is the *inter-action*.

Aim of this work is to tabulate $P(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t)$ for the three Coulomb cases illustrated above, P_{on}^g , P_{on}^e , P_{in} . In the hot case, for small $t \ll 1$, for each of these cases, the *primitive approximation* [1]

$$P_{\text{primitive}} \approx t\phi, \quad (2.10)$$

holds.

Therefore, introducing a timestep τ , we can find the PDCM at all subsequent timesteps through the following convolution path integral

$$\rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; n\tau) = \int \rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_{1,1}, \mathbf{r}_{2,1}; \tau) \rho(\mathbf{r}_{1,1}, \mathbf{r}_{2,1} | \mathbf{r}_{1,2}, \mathbf{r}_{2,2}; \tau) \cdots \rho(\mathbf{r}_{1,n-1}, \mathbf{r}_{2,n-1} | \mathbf{r}'_1, \mathbf{r}'_2; \tau) \times \prod_{k=1}^{n-1} \sqrt{g(\mathbf{r}_{1,k})g(\mathbf{r}_{2,k})} d\mathbf{r}_{1,k} d\mathbf{r}_{2,k}, \quad (2.11)$$

where we denote with $\mathbf{r}_{i,k}$ the position of particle i at timeslice k , $d\mathbf{r} = d\theta d\varphi$, and we choose τ small enough so that

$$\rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; \tau) \approx \rho_0(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; \tau) e^{-\tau[\phi(\mathbf{r}_1, \mathbf{r}_2) + \phi(\mathbf{r}'_1, \mathbf{r}'_2)]/2} \quad (2.12)$$

$$= \rho_{10}(\mathbf{r}_1 | \mathbf{r}'_1; \tau) \rho_{20}(\mathbf{r}_2 | \mathbf{r}'_2; \tau) e^{-\tau[\phi(\mathbf{r}_1 | \mathbf{r}_2) + \phi(\mathbf{r}'_1 | \mathbf{r}'_2)]/2}, \quad (2.13)$$

is a good approximation. We will perform the $4(n-1)$ integrations in Eq. (2.11) through a simple quadrature method.

According to the analysis of Ref. [6] we can further approximate

$$g^{1/4}(\mathbf{r}_i) \rho_{i0}(\mathbf{r}_i | \mathbf{r}'_i; \tau) g^{1/4}(\mathbf{r}'_i) \approx \exp \left\{ -\frac{1}{2\mu_i \tau} \delta_{\alpha\beta} (\mathbf{r}_i - \mathbf{r}'_i)^\alpha (\mathbf{r}_i - \mathbf{r}'_i)^\beta - \tau \frac{V_R(x_i) + V_R(x'_i)}{2} \right\}, \quad (2.14)$$

where $\delta_{\alpha\beta}$ is the Kronecker delta, $x_i^2 = \delta_{\alpha\beta} \mathbf{r}_i^\alpha \mathbf{r}_i^\beta$, and V_R is the effective potential due to the curvature of the Sphere, $R = 2M^2$ with $M = 1/a$, namely

$$V_R(x) = -\frac{M^2}{6} - \frac{5(Mx)^2 - 3 + [(Mx)^2 + 3] \cos(2Mx)}{48x^2 \sin^2(Mx)}, \quad (2.15)$$

which is equation (2.32) of Ref. [6].

We may additionally introduce the following exact Feynman-Kac expression for the one free body density matrix

$$g^{1/4}(\mathbf{r}_i) \rho_{i0}(\mathbf{r}_i | \mathbf{r}'_i; t) g^{1/4}(\mathbf{r}'_i) = \frac{\langle e^{-S_R} \rangle}{2\pi\mu_i t}, \quad (2.16)$$

$$S_R = \int_0^t dt' V_R(x) \quad (2.17)$$

where $\langle \dots \rangle$ denotes an average over all single particle Gaussian random walks from $\mathbf{r}_i$ to $\mathbf{r}'_i$ in a time t . In Appendix A we give the diagonal part $\ln[2\pi\mu_i t \sqrt{g(\mathbf{0})} \rho_{i0}(\mathbf{0} | \mathbf{0}; t)]$ up to orders t^8 . Our Eq. (A1) coincides with the perturbative expansion (3.23) of Ref. [6] for $d = 2$ spatial dimensions.

Even if in general we will be interested in the full $P(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}'_1, \mathbf{r}'_2; t)$ in practice in a many body simulation we may use the *end point approximation* for the inter-action as shown in section IV.F of Ref. [1] where one only needs the diagonal components $P(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_1, \mathbf{r}_2; t) = P(d_g(\mathbf{r}_1, \mathbf{r}_2)/a; t)$. Where on a sphere the geodesic distance $d_g(\mathbf{r}_1, \mathbf{r}_2)$ between any two points $\mathbf{r}_1$ and $\mathbf{r}_2$ is bounded from above by πa . Due to the Sphere symmetry, without loss of generality, we may choose $\mathbf{r}_1 = (0, 0)$ and $\mathbf{r}_2 = (\theta, 0)$, so that $d_g = a\theta$.

On the diagonal we will have

$$\rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_1, \mathbf{r}_2; t) = \rho(d_g(\mathbf{r}_1, \mathbf{r}_2)/a; t) = \rho_0(d_g(\mathbf{r}_1, \mathbf{r}_2)/a; t) e^{-P(d_g(\mathbf{r}_1, \mathbf{r}_2)/a; t)}, \quad (2.18)$$

we will then tabulate

$$P(m\delta; n\tau) = -\ln \left[\frac{\rho(m\delta; n\tau)}{\rho_0(m\delta; n\tau)} \right], \quad n = 1, 2, 3, \dots, \quad m = 1, 2, 3, \dots, m_{\max} \quad (2.19)$$

for given small τ and δ so that $m_{\max} = \pi/\delta$. For an imaginary time $t_n = n\tau$ this will require a number of $4(n-1)$ quadratures for each m to calculate $\rho(m\delta; n\tau) = \rho(d_g(\mathbf{r}_1, \mathbf{r}_2)/a; t_n) = \rho(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_1, \mathbf{r}_2; t_n)$ and $\rho_0(m\delta; n\tau) = \rho_0(d_g(\mathbf{r}_1, \mathbf{r}_2)/a; t_n) = \rho_0(\mathbf{r}_1, \mathbf{r}_2 | \mathbf{r}_1, \mathbf{r}_2; t_n)$ for $m = 1, 2, 3, \dots, m_{\max}$, through the corresponding path integral convolution (2.11). Each convolution requires 4 quadratures over the azimuthal angle $\varphi \in [-\pi, \pi[$ and the polar angle $\theta \in [-\pi/2, \pi/2]$ of each particle. Choosing an angular discretization $\delta_\varphi = 2\delta_\theta$ we will have that each angular integral reduces to a sum over $l = \pi/\delta_\theta$ terms. Then in order to tabulate $P(d_g/a; t_n)$ we need $l^{4(n-1)}$ functions evaluations for each m . This becomes more computationally demanding at later imaginary times t but in a many body simulation the pair-product inter-action is needed at a fixed small t , which determines the discretization error [1]. So one needs to tabulate it just once, at the beginning of the simulation.

In Table I we show $P(m\delta; n\tau)$ for $a = 1, m_1 = m_2 = m_e, Z_1 = -Z_2 = -1, \tau = 0.1, \delta = 0.1$ and $n = 3$ and compare it to the primitive approximation $n\tau\phi(m\delta)$. From the comparison we note how $|P(\delta; n\tau)| < n\tau|\phi(\delta)|$ for $n > 1$, in agreement with the cusp condition [2]. In particular $P(d_g \rightarrow 0; t_n)$ remains finite for $n > 1$.


```

c      loop over timesteps
10     th1=ith1(1)*dth-pi/2
        phi1=iph1(1)*dph
        th2=ith2(1)*dth-pi/2
        ph2=iph2(1)*dph
        dg=a*acos(sin(th1)*sin(th2)+cos(th1)*cos(th2)*cos(phi-ph2))
        de=2*a*sin(dg/2/a)
        if(dg.lt.eps)goto 20
        fa=rho(potk,tau,a,mu1,mu2,z1,z2,
&         th10,ph10,th20,ph20,th1,ph1,th2,ph2,dg,de,dg0,de0)
        faid=rhoid(tau,a,mu1,mu2,
&         th1,ph1,th2,ph2,th10,ph10,th20,ph20)
        do j=2,ni
            th1=ith1(j-1)*dth-pi/2
            phi1=iph1(j-1)*dph
            th2=ith2(j-1)*dth-pi/2
            ph2=iph2(j-1)*dph
            th1p=ith1(j)*dth-pi/2
            phi1p=iph1(j)*dph
            th2p=ith2(j)*dth-pi/2
            ph2p=iph2(j)*dph
            dg=a*acos(sin(th1)*sin(th2)+cos(th1)*cos(th2)*cos(phi-ph2))
            de=2*a*sin(dg/2/a)
            dgp=a*acos(sin(th1p)*sin(th2p)+cos(th1p)*cos(th2p)*
&             cos(phi1p-ph2p))
            dep=2*a*sin(dgp/2/a)
            if(dg.lt.eps)goto 20
            if(dgp.lt.eps)goto 20
            fa=fa+dth**2.*dph**2.*rho(potk,tau,a,mu1,mu2,z1,z2,
&             th1,ph1,th2,ph2,th1p,phi1p,th2p,ph2p,dg,de,dgp,dep)
            faid=faid+dth**2.*dph**2.*rhoid(tau,a,mu1,mu2,
&             th1,ph1,th2,ph2,th1p,phi1p,th2p,ph2p)
        enddo
        th1=ith1(ni)*dth-pi/2
        phi1=iph1(ni)*dph
        th2=ith2(ni)*dth-pi/2
        ph2=iph2(ni)*dph
        dg=a*acos(sin(th1)*sin(th2)+cos(th1)*cos(th2)*
&         cos(phi1-ph2))
        de=2*a*sin(dg/2/a)
        if(dg.lt.eps)goto 20
        fa=fa+dth**2.*dph**2.*rho(potk,tau,a,mu1,mu2,z1,z2,
&         th1,ph1,th2,ph2,th10,ph10,th20,ph20,dg,de,dg0,de0)
        faid=faid+dth**2.*dph**2.*rhoid(tau,a,mu1,mu2,
&         th1,ph1,th2,ph2,th10,ph10,th20,ph20)
        pcdm = pcdm+fa
        pidm = pidm+faid
20     do k=1,ni
        ith1(k)=ith1(k)+1
        if(ith1(k).le.na) goto 10
        ith2(k)=ith2(k)+1
        ith1(k)=1
        if(ith2(k).le.na) goto 10
        iph1(k)=iph1(k)+1
        ith1(k)=1
        ith2(k)=1
        if(iph1(k).le.na) goto 10
        iph2(k)=iph2(k)+1
        ith1(k)=1
        ith2(k)=1
        iph1(k)=1
        if(iph2(k).le.na) goto 10
        do j=1,k
            ith1(j)=1
            ith2(j)=1
            iph1(j)=1
            iph2(j)=1
        enddo
    enddo
c      tabulate P=-log(rho/rho0)
        pcdm = -log(pcdm/pidm)
        write(8,100) a*dg0,pcdm,ti*pot(potk,a,z1,z2,dg0,de0)
        write(*,*) a*dg0,pcdm,ti*pot(potk,a,z1,z2,dg0,de0)
    enddo

100    format(3(d10.3,3x))
        close(unit=8)

stop
end

function fact(n)
implicit none
factorial function
real*8 fact,p
integer i,n

p=1.
do i=1,n
    p=p*dble(i)
enddo
fact=p

return
end

function vr(a,mu,th,ph)
implicit none
Eq. (2.32) Bastianelli and Corradini EPJC 77, 731 (2017)
real*8 vr,a,mu,th,ph
real*8 pi,m,x,mx,eps
eps=1.d-3
pi=acos(-1.d0)

m =1/a
x =sqrt(th**2.+ph**2.)
mx=m*x
vr=-m**2./6-(5*mx**2.-3*(mx**2.+3)*cos(2*mx))/48/x**2./sin(mx)**2.
if(abs(vr).gt.1/eps) vr=0.

```

```

return
end

function pot(potk,a,z1,z2,dg,de)
implicit none
c the Coulomb potential
real*8 pot,a,z1,z2
real*8 pi,de,dg
character potk*3
pi=acos(-1.d0)

if(potk.eq.'in') then
pot=-z1*z2*log(de)
elseif(potk.eq.'one') then
pot=z1*z2/de
elseif(potk.eq.'ong') then
pot=z1*z2/dg
else
write(*,*) 'no pair-potential available !'
stop
endif

return
end

function rho(potk,tau,a,mu1,mu2,z1,z2,
& th1,ph1,th2,ph2,thip,phip,th2p,ph2p,dg,de,dgp,dep)
implicit none
c primitive approximation for rho
real*8 rho,tau,a,mu1,mu2,z1,z2,dg,de,dgp,dep
real*8 th1,ph1,th2,ph2,thip,phip,th2p,ph2p
real*8 pot,vr,ddx1,ddx2
character potk*3

ddx1=(th1-thip)**2.+(ph1-phip)**2.
ddx2=(th2-th2p)**2.+(ph2-ph2p)**2.

rho = exp(-ddx1/2/mu1/tau-tau*
& (vr(a,mu1,th1,ph1)+vr(a,mu1,thip,phip))/2)
rho =rho*exp(-ddx2/2/mu2/tau-tau*
& (vr(a,mu2,th2,ph2)+vr(a,mu2,th2p,ph2p))/2)
rho =rho*exp(-tau*(pot(potk,a,z1,z2,dg,de)+
& pot(potk,a,z1,z2,dgp,dep))/2)

return
end

function rhoid(tau,a,mu1,mu2,
& th1,ph1,th2,ph2,thip,phip,th2p,ph2p)
implicit none
c primitive approximation for rho
real*8 rhoid,tau,a,mu1,mu2
real*8 th1,ph1,th2,ph2,thip,phip,th2p,ph2p
real*8 vr,ddx1,ddx2

ddx1=(th1-thip)**2.+(ph1-phip)**2.
ddx2=(th2-th2p)**2.+(ph2-ph2p)**2.

rhoid = exp(-ddx1/2/mu1/tau-tau*
& (vr(a,mu1,th1,ph1)+vr(a,mu1,thip,phip))/2)
rhoid =rhoid*exp(-ddx2/2/mu2/tau-tau*
& (vr(a,mu2,th2,ph2)+vr(a,mu2,th2p,ph2p))/2)

return
end

```

Appendix A: $\rho_{i0}(\mathbf{0}, \mathbf{0}; t)$

Calling $\mu_i = m_i/m_e$ and $R = 2/a^2$ the Sphere scalar curvature, we find from equation (3.23) of Ref. [6]

$$\begin{aligned} \ln[2\pi\mu_i t \sqrt{g(\mathbf{0})}\rho_{i0}(\mathbf{0}|\mathbf{0}; t)] = & \frac{tR}{12} + \\ & \frac{(tR)^2}{6!} \frac{1}{2} + \\ & \frac{(tR)^3}{9!} 16 + \\ & \frac{(tR)^4}{10!} \frac{59}{4} + \\ & \frac{(tR)^5}{11!} \frac{58}{3} + \\ & \frac{(tR)^6}{13!} \frac{550789}{1260} + \\ & \frac{(tR)^7}{14!} \frac{46838}{45} + \\ & \frac{(tR)^8}{17!} \frac{596004707}{720} + \mathcal{O}(t^9). \end{aligned} \tag{A1}$$

AUTHOR DECLARATIONS

Conflicts of interest

None declared.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Funding

None declared.

-
- [1] D. M. Ceperley, Path integrals in the theory of condensed Helium, Rev. Mod. Phys. **67**, 279 (1995).
 - [2] E. L. Pollock, Properties and computation of the Coulomb pair density matrix, Comput. Phys. Commun. **52**, 49 (1988).
 - [3] P. Vieillefosse, Coulomb Pair Density Matrix I, J. Stat. Phys. **74**, 1195 (1994).
 - [4] P. Vieillefosse, Coulomb Pair Density Matrix II, J. Stat. Phys. **80**, 461 (1995).
 - [5] R. Fantoni, Plasma living in a curved surface at some special temperature, Physica A **177**, 524 (2019).
 - [6] F. Bastianelli and O. Corradini, On the simplified path integral on spheres, Eur. Phys. J. C **77**, 731 (2017).