# Brownian Bridge for Coherent State Path Integral Monte Carlo

Riccardo Fantoni*
*Università di Trieste, Dipartimento di Fisica, strada Costiera 11, 34151 Grignano (Trieste), Italy*
(Dated: April 26, 2026)

...

Keywords: Quantum Many Body; Quantum Monte Carlo; Coherent States; Path Integral; Brownian Bridge

## I. INTRODUCTION

We recently [1] constructed a new Path Integral Monte Carlo (PIMC) algorithm using Coherent States (CS), [2–7] in place of the more usual Plane Waves (PW), as the complete set over which to expand the thermal density matrix. We called this algorithm with the acronym CSPIMC. We proposed our new algorithm as an alternative to the more common Plane Waves PIMC of Ref. [8] to which we will refer with the acronym PWPIMC.

Although the mathematics of Ref. [1] where we proposed the new algorithm is completely correct, the numerics in that paper was not complete. In particular it was lacking a justification of the Brownian bridges [8, 9] used in the CSPIMC scheme to sample the permutations sum necessary to calculate the properties of identical particles (being them bosons or fermions).

In the present work we will fill that gap proposing a correct realization of the Brownian bridge move within the CSPIMC algorithm. And we will justify our new proposal.

The CSPIMC numerical results that we obtain for Helium on a plane compare well with the results from the conventional PWPIMC. Within CSPIMC the continuum limit for the imaginary time must be taken simultaneously to the limit of an infinite elastic constant of the harmonic oscillator of unit mass subtending the coherent states.

## II. PHYSICS MODEL

In Ref. [1] we performed several Path Integral Monte Carlo (PIMC) simulations for a two dimensional, $d = 2$, $^4$He liquid [10] with either Boltzmann or Bose statistics.

The liquid has a surface number density $n = N/\Omega$ where $N$ is the number of Helium atoms in an area $\Omega$ of a flat surface, at an inverse temperature $\beta = 1/k_B T$ with $k_B$ Boltzmann constant. The Hamiltonian of the fluid is $\hat{H} = \hat{T} + \upsilon\hat{V} = \hat{P}^2/2m + \upsilon V(Q)$ with $\vec{Q} = (\boldsymbol{q}_1, \boldsymbol{q}_2, \ldots, \boldsymbol{q}_N)$ the particles positions and $\vec{P} = (\hat{\boldsymbol{p}}_1, \hat{\boldsymbol{p}}_2, \ldots, \hat{\boldsymbol{p}}_N) = -i(\boldsymbol{\nabla}_{\boldsymbol{q}_1}, \boldsymbol{\nabla}_{\boldsymbol{q}_2}, \ldots, \boldsymbol{\nabla}_{\boldsymbol{q}_N})$ the particles momenta. The potential energy $V(Q)$ enters the Hamiltonian with a coupling constant $\upsilon$.

The computer experiments [1] were done for for $N = 16$ $^4$He atoms ($m = 0.0830594 \approx 1/12$ Å$^{-2}$K$^{-1}$) in two dimensions $d = 2$, in a square periodic cell of area $\Omega = L^2$, interacting with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556$ Å, $\varepsilon = 10.22$ K and a cutoff distance $r_{\rm cut} = 2.5$ Å (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a number density $n = N/L^2 = 0.05$ Å$^{-2}$ and various temperatures $T$ or timeslices number $M$, either with Boltzmann and Bose statistics. As we can see from the phase diagram of Ref. [10], at this density the fluid undergoes a phase transition from a fluid phase at high temperature to a superfluid phase at low temperatures.

In that work [1] we compared two different PIMC simulations: the conventional Plane Waves PIMC (**PW**PIMC) [8] and our newly devised Coherent States PIMC (**CS**PIMC) algorithm. The two PIMC differ for the expression of the hot kinetic density matrix at an imaginary timestep $\tau = \beta/M$ with $M$ a large number of timeslices. The short imaginary time density matrix CSPIMC expression is reviewed in the next section. The PWPIMC requires a multidimensional integral over $dNM$ coordinates whereas our CSPIMC requires $5dMN$ integrations, the usual $dNM$ real particles coordinates, $2dNM$ ghost particles coordinates, and $2dNM$ ghost particles momenta. The coherent states are generated by a harmonic oscillator (h.o.) with mass $m_{h.o.}$ and elastic constant $k = m_{h.o.}\omega^2$. We introduced the parameter $\xi \equiv m_{h.o.}\omega/2$ and the adimensional one $\varphi \equiv \xi\tau/m = (\sigma_{p.w.}/\sigma_{c.s.})^2/2$ where $\sigma_{p.w.} \equiv \sqrt{2\lambda\tau}$ is the standard deviation in the plane wave scheme, with $\lambda = 1/2m$, and $\sigma_{c.s.} \equiv \sqrt{1/m_{h.o.}\omega}$ is the standard deviation in the coherent state scheme. As illustrated in Ref. [1] in order to compare PWPIMC and CSPIMC it is necessary to keep $\varphi$ fixed.

* riccardo.fantoni@scuola.istruzione.it

In our quantum simulations we have to take care of two limiting procedures: The continuum limit where the ultraviolet cutoff $\tau \to 0$ or $M \to \infty$ at fixed absolute temperature $T$ and the thermodynamic limit where $N \to \infty$ and $\Omega \to \infty$ at fixed density $n$. In particular in Ref. [1] we explicitly worried about the continuum limit and we proved that in the CSPIMC scheme it is necessary to keep $\varphi$ constant as $\tau \to 0$ so that $\xi \to \infty$. To mimic the thermodynamic limit we simply used a periodic square cell which permeates the whole infinite space. This procedure gives rise to spurious finite size effects unavoidable on a computer.

We worked in units where $\hbar = k_B = 1$ so that the imaginary time has dimensions of temperature.

## III. THE SHORT TIME DENSITY MATRIX FOR CSPIMC

The short imaginary time, $\tau$, Green function for CSPIMC can be written as follows [1]:

$$\rho(Q, Q'; \tau) \approx e^{-\tau \upsilon V(Q)} \frac{1}{N!} \sum_P \prod_{\alpha=1}^{N} \zeta_\alpha \Big( \boldsymbol{q}_{P\alpha} | \boldsymbol{q}'_\alpha; \tau, m, \xi \Big), \tag{3.1}$$

with $P$ a permutation of the $N$ **real** identical particles and

$$\zeta_\alpha \Big( \boldsymbol{q} | \boldsymbol{q}'; \tau, m, \xi \Big) \approx \int \frac{d\boldsymbol{\mathcal{Q}}_\alpha \, d\boldsymbol{\mathcal{P}}_\alpha}{(2\pi)^d} \frac{d\boldsymbol{\mathcal{Q}}'_\alpha \, d\boldsymbol{\mathcal{P}}'_\alpha}{(2\pi)^d} \psi_\alpha(\boldsymbol{q}_\alpha, \boldsymbol{\mathcal{Q}}_\alpha, \boldsymbol{\mathcal{P}}_\alpha) {\psi_\alpha}^*(\boldsymbol{q}'_\alpha, \boldsymbol{\mathcal{Q}}'_\alpha, \boldsymbol{\mathcal{P}}'_\alpha) G_\alpha \exp\left[ -\tau(\boldsymbol{\mathcal{P}}_\alpha^2 + {\boldsymbol{\mathcal{P}}'}_\alpha^2)/4m \right], \tag{3.2}$$

where $\psi_\alpha(\boldsymbol{q}_\alpha, \boldsymbol{\mathcal{Q}}_\alpha, \boldsymbol{\mathcal{P}}_\alpha)$ is the CS wave function with ghost canonical variables $\boldsymbol{\mathcal{Q}}_\alpha$ and $\boldsymbol{\mathcal{P}}_\alpha$, the **ghost** of the real particle $\alpha$, namely

$$\psi_\alpha(\boldsymbol{q}_\alpha, \boldsymbol{\mathcal{Q}}_\alpha, \boldsymbol{\mathcal{P}}_\alpha) \equiv \langle \boldsymbol{q}_\alpha | \boldsymbol{\mathcal{Q}}_\alpha, \boldsymbol{\mathcal{P}}_\alpha \rangle = \left( \frac{2\xi}{\pi} \right)^{d/4} \exp\left[ -\xi \left( \boldsymbol{q}_\alpha - \boldsymbol{\mathcal{Q}}_\alpha \right)^2 + i \boldsymbol{\mathcal{P}}_\alpha \cdot \left( \boldsymbol{q}_\alpha - \boldsymbol{\mathcal{Q}}_\alpha \right) \right], \tag{3.3}$$

$$G_\alpha = \exp\left[ -\frac{1}{2}(|\boldsymbol{a}|^2 + |\boldsymbol{b}|^2) + \boldsymbol{a}^* \cdot \boldsymbol{b} + \frac{i}{2}(\boldsymbol{\mathcal{Q}}_\alpha \cdot \boldsymbol{\mathcal{P}}_\alpha - \boldsymbol{\mathcal{Q}}'_\alpha \cdot \boldsymbol{\mathcal{P}}'_\alpha) \right], \tag{3.4}$$

$$\boldsymbol{a} \equiv \frac{1}{\sqrt{4\xi}} (2\xi \boldsymbol{\mathcal{Q}}_\alpha + i \boldsymbol{\mathcal{P}}_\alpha), \tag{3.5}$$

$$\boldsymbol{b} \equiv \frac{1}{\sqrt{4\xi}} (2\xi \boldsymbol{\mathcal{Q}}'_\alpha + i \boldsymbol{\mathcal{P}}'_\alpha), \tag{3.6}$$

where $G_\alpha$ is the scalar product of the two ghosts $\boldsymbol{\mathcal{Q}}$ and $\boldsymbol{\mathcal{Q}}'$.

We can then define a reduced high temperature density matrix

$$\tilde{\rho}(s, s'; \tau) \propto e^{-\tau \upsilon V(Q)} \prod_{\alpha=1}^{N} \psi_\alpha(\boldsymbol{q}_\alpha, \boldsymbol{\mathcal{Q}}_\alpha, \boldsymbol{\mathcal{P}}_\alpha) {\psi_\alpha}^*(\boldsymbol{q}'_\alpha, \boldsymbol{\mathcal{Q}}'_\alpha, \boldsymbol{\mathcal{P}}'_\alpha) G_\alpha e^{-\tau(\boldsymbol{\mathcal{P}}_\alpha^2 + {\boldsymbol{\mathcal{P}}'}_\alpha^2)/4m}, \tag{3.7}$$

with $s = (\vec{Q}, \vec{\mathscr{Q}}, \vec{\mathscr{P}}, P)$ the configurations space for a single link where $\vec{\mathscr{Q}} = (\boldsymbol{\mathcal{Q}}_1, \boldsymbol{\mathcal{Q}}_2, \ldots, \boldsymbol{\mathcal{Q}}_N)$ and $\vec{\mathscr{P}} = (\boldsymbol{\mathcal{P}}_1, \boldsymbol{\mathcal{P}}_2, \ldots, \boldsymbol{\mathcal{P}}_N)$.

Discretizing the imaginary time into $M$ timeslices with $\tau = \beta/M$ we can use Trotter [12] identity to introduce $M$ links $\tilde{\rho}(s_k, s_{k+1}; \tau)$ for $k = 0, 1, 2, \ldots, M-1$ to reconstruct the density matrix at an inverse temperature $\beta$ as the following path integral

$$\begin{aligned} \rho(s_0, s'_M; \beta) \propto \sum_P \int \tilde{\rho}(s_0, s'_1; \tau) \tilde{\rho}(s_1, s'_2; \tau) \cdots \tilde{\rho}(s_{M-1}, s'_M; \tau) \times \\ \prod_{\alpha=1}^{N} \prod_{k=1}^{M-1} d\boldsymbol{q}_{\alpha,k} \frac{d\boldsymbol{\mathcal{Q}}^k_{\alpha,k-1} d\boldsymbol{\mathcal{P}}^k_{\alpha,k-1}}{(2\pi)^d} \frac{d\boldsymbol{\mathcal{Q}}^k_{\alpha,k} d\boldsymbol{\mathcal{P}}^k_{\alpha,k}}{(2\pi)^d} \frac{d\boldsymbol{\mathcal{Q}}^M_{\alpha,M-1} d\boldsymbol{\mathcal{P}}^M_{\alpha,M-1}}{(2\pi)^d} \frac{d\boldsymbol{\mathcal{Q}}^M_{\alpha,M} d\boldsymbol{\mathcal{P}}^M_{\alpha,M}}{(2\pi)^d} \end{aligned} \tag{3.8}$$

where $\boldsymbol{q}_{\alpha,k}$ are the coordinates of the real particle $\alpha$ at timeslice $k$, $\boldsymbol{\mathcal{Q}}^l_{\alpha,k}$ and $\boldsymbol{\mathcal{P}}^l_{\alpha,k}$ are respectively the canonical coordinates and momenta of the ghost of particle $\alpha$ at the timeslice $k$ in the link $l-1 \to l$. In our notation $s_k = (\vec{Q}_k, \vec{\mathscr{Q}}^{k+1}_k, \vec{\mathscr{P}}^{k+1}_k, P)$ and $s'_k = (\vec{Q}_k, \vec{\mathscr{Q}}^k_k, \vec{\mathscr{P}}^k_k, P)$ with $\vec{Q}_k = (\boldsymbol{q}_{1,k}, \boldsymbol{q}_{2,k}, \ldots, \boldsymbol{q}_{N,k})$, $\vec{\mathscr{Q}}^l_k = (\boldsymbol{\mathcal{Q}}^l_{1,k}, \boldsymbol{\mathcal{Q}}^l_{2,k}, \ldots, \boldsymbol{\mathcal{Q}}^l_{N,k})$, and $\vec{\mathscr{P}}^l_k = (\boldsymbol{\mathcal{P}}^l_{1,k}, \boldsymbol{\mathcal{P}}^l_{2,k}, \ldots, \boldsymbol{\mathcal{P}}^l_{N,k})$.

The Monte Carlo used is the standard Metropolis algorithm [9, 13]. In our CSPIMC simulations we have two kinds of moves: a singleslice *displace move* and a multislice *Brownian bridge*.

In the displace move we choose as transition move a uniform displacement of each of the $dMN$ real path coordinate $\boldsymbol{q}_{\alpha,k} \to \boldsymbol{q}_{\alpha,k} + (1/2 - \eta)\boldsymbol{\Delta}$ for $\alpha = 1, \ldots, N$ and $k = 1, \ldots, M$, where $\eta$ is a uniform pseudorandom number in $[0, 1)$ and $\boldsymbol{\Delta}$ a fixed $d$-dimensional vector whose magnitude is chosen so to have acceptance ratios close to $1/2$. And of each of the $4dMN$ ghost path canonical variables $\boldsymbol{\mathcal{Q}}^k_{\alpha,k-1} \to \boldsymbol{\mathcal{Q}}^k_{\alpha,k-1} + (1/2 - \eta)\boldsymbol{\Delta}$, $\boldsymbol{\mathcal{P}}^k_{\alpha,k-1} \to \boldsymbol{\mathcal{P}}^k_{\alpha,k-1} + (1/2 - \eta)\boldsymbol{\Delta}$, and $\boldsymbol{\mathcal{Q}}^k_{\alpha,k} \to \boldsymbol{\mathcal{Q}}^k_{\alpha,k} + (1/2 - \eta)\boldsymbol{\Delta}$, $\boldsymbol{\mathcal{P}}^k_{\alpha,k} \to \boldsymbol{\mathcal{P}}^k_{\alpha,k} + (1/2 - \eta)\boldsymbol{\Delta}$. So that the transition probability density is just a constant and drops out of the acceptance probability.

The sum over permutations in Eq. (3.8) can be applied just to the real particles coordinates in the first (or last) timeslice. In order to sample the permutation sum we need to construct two Brownian bridges [8, 9] between the initial positions taken from the paths of two particles $\alpha$ and $\gamma$ at the same timeslice and the final positions taken from the same two paths at a subsequent timeslice but exchaged, $\gamma$ and $\alpha$, so to connect particle $\alpha$ at the initial timeslice to particle $\gamma$ at the final timeslice with one bridge and particle $\gamma$ at the initial timeslice with particle $\alpha$ at the final timeslice. We accept or reject the move according to Metropolis algorithm [9, 13]. If the move is accepted one creates an exchange of two particles. Since any permutation of $N$ particles can be obtained by composing a finite number of particles exchanges this is sufficient to simulate Bose-Einstein statistics in an exact numerical way. The Brownian bridge move is described in the next section.

In our simulations we measured the total potential energy $\mathscr{E}_P = \upsilon\langle V\rangle$ and the total kinetic energy

$$\mathscr{E}_K = \begin{cases} \dfrac{dN}{2\tau} - \dfrac{\langle(\vec{Q}_k - \vec{Q}_{k-1})^2\rangle}{4\lambda\tau^2} & \text{PWPIMC} \\ \dfrac{dN}{2\tau}\dfrac{\varphi}{1+\varphi} - \mu\dfrac{\langle\vec{\mathscr{P}}_k^2 + \vec{\mathscr{P}}_k'^2\rangle}{4m} & \text{CSPIMC} \end{cases} \tag{3.9}$$

where, given an observable $\mathscr{O}$, we denote with $\langle\mathscr{O}\rangle = \mathrm{tr}(\rho\mathscr{O})/\mathrm{tr}(\rho)$ the thermal average with 'tr' the trace operation. These expressions for $\mathscr{E}_K$ and $\mathscr{E}_P$ corresponds to the *thermodynamic* estimators of Ref. [8]. $\mathscr{E} = \mathscr{E}_K + \mathscr{E}_P$ is the total internal energy. The parameter $\mu$, in the CSPIMC case, is necessary in order to find agreement with the kinetic energy of the PWPIMC, as explained in Ref. [1]. The kinetic energy from Eq. (3.9) is the small difference between two large quantities, infinite in the continuum $\tau \to 0$ limit. Note that in the $\xi \to \infty$ limit $\varphi/(1+\varphi) \to 1$ in agreement with the PWPIMC case.

## IV. THE BRIDGE MOVE

In order to take into account the particles permutations it is necessary to construct two Brownian bridges between two different [1] randomly chosen particles in two randomly chosen many body beads to generate an exchange between the two particles. With one bridge we connect a real particle 1 and its ghosts on timeslice $i$ to particle 2 and its ghosts on timeslice $j$ and with the other we connect particle 2 and its ghosts on timeslice $i$ to particle 1 and its ghosts on timeslice $j$ with $i < j$. This will produce an *exchange* of particles 1 and 2 and their respective ghosts.

The Brownian bridge between particle 1 at $\boldsymbol{q}_{1,i}$ and particle 2 at $\boldsymbol{q}_{2,j}$ is built through the following multislice transition move [8],

$$\boldsymbol{q}_{\mathrm{new},i} = \boldsymbol{q}_{1,i} \tag{4.1}$$

$$\boldsymbol{q}_{\mathrm{new},k} = \boldsymbol{q}_{\mathrm{new},k-1} + \frac{\boldsymbol{q}_{2,j} - \boldsymbol{q}_{\mathrm{new},k-1}}{j-k+1} + \xi \qquad k = i+1, \ldots, j \tag{4.2}$$

where $\xi$ is a random number with a Gaussian probability distribution [2] with zero mean and variance $\sigma^2(j-k)/(j-k+1)$ where $\sigma^2 = 2\lambda\tau$ is the diagonal free particle variance. The same move (4.1)-(4.2) is also carried out on the ghosts coordinates $\boldsymbol{\mathcal{Q}}_{\alpha,k}^k$ and $\boldsymbol{\mathcal{Q}}_{\alpha,k}^{k+1}$ and we leave untouched the momenta of the two ghosts.

The Metropolis (rejection) method [13] can sample any probability distribution provided that the transition rule satisfies detailed balance and ergodicity. The Metropolis algorithm is a particular way of ensuring that the transition rule satisfies detailed balance. It does this by splitting the *transition probability* into an "a priori" *sampling distribution* $T(S_{\mathrm{old}} \to S_{\mathrm{new}})$ (which is a probability distribution that we already know) and an *acceptance probability* $A(S_{\mathrm{old}} \to S_{\mathrm{new}})$.

$$P(S_{\mathrm{old}} \to S_{\mathrm{new}}) = T(S_{\mathrm{old}} \to S_{\mathrm{new}})A(S_{\mathrm{old}} \to S_{\mathrm{new}}), \tag{4.3}$$

where we denote with $S \equiv (\{s_k\}, \{s_k'\})$, $S_{\mathrm{old}}$ is the state configuration before the Markov move and $S_{\mathrm{new}}$ the proposed configuration state after the move produced by the transition with probability $P$.

In the generalized Metropolis procedure [9], trial moves are accepted according to:

$$A(S_{\mathrm{old}} \to S_{\mathrm{new}}) = \min[1, q(S_{\mathrm{old}} \to S_{\mathrm{new}})], \tag{4.4}$$

---

[1] A bridge between the same particle can still be used to sample the density matrix of distinguishable particles as can be done with the displacement move.

[2] This can be generated with the Box-Muller algorithm [9] for example.

where

$$q(S_{\rm old} \to S_{\rm new}) = \frac{\pi(S_{\rm new})T(S_{\rm new} \to S_{\rm old})}{\pi(S_{\rm old})T(S_{\rm old} \to S_{\rm new})}. \tag{4.5}$$

where $\pi \propto e^{-\mathscr{S}} \propto \rho(s_0, s_1'; \tau)\rho(s_1, s_2'; \tau) \cdots \rho(s_{M-1}, s_M'; \tau)$ is the action probability distribution in the $s = (\{s_k\})$ configurations space. The transition probability corresponding to the move of Eqs. (4.1)-(4.2) is then given by

$$\frac{T(S_{\rm new} \to S_{\rm old})}{T(S_{\rm old} \to S_{\rm new})} \propto \exp\left\{ \frac{1}{4\lambda\tau} \sum_{k=i+1}^{j} \Big[ (\boldsymbol{q}_{{\rm new},k} - \boldsymbol{q}_{{\rm new},k-1})^2 + (\boldsymbol{\mathcal{Q}}^k_{{\rm new},k} - \boldsymbol{\mathcal{Q}}^k_{{\rm new},k-1})^2 + (\boldsymbol{\mathcal{Q}}^{k+1}_{{\rm new},k} - \boldsymbol{\mathcal{Q}}^{k+1}_{{\rm new},k-1})^2 \right.$$
$$\left. - (\boldsymbol{q}_{{\rm old},k} - \boldsymbol{q}_{{\rm old},k-1})^2 - (\boldsymbol{\mathcal{Q}}^k_{{\rm old},k} - \boldsymbol{\mathcal{Q}}^k_{{\rm old},k-1})^2 - (\boldsymbol{\mathcal{Q}}^{k+1}_{{\rm old},k} - \boldsymbol{\mathcal{Q}}^{k+1}_{{\rm old},k-1})^2 \Big] \right\}. \tag{4.6}$$

In order to produce an exchange of two particles 1 and 2 one needs a combination of two bridge transitions as described above. Together with the real particles we propose also the exchange of the relative ghosts. Any permutation can be reached through a two particles exchange so the bridge transition move allows to sample the sum in Eq. (3.8).

In order to increase the acceptance ratios of this bridge and swap moves it is necessary to diminish $\xi$ and the number $j - i$ of its timeslices. Of course our kinetic energy estimator is left unchanged by this bridge move since the ghosts momenta are not moved at all. But the potential energy obtained with only the bridge move converges to the same value obtained with the displace and the bridge moves together, as it should. Moreover the potential energy becomes more negative as $\xi$ increases at constant $\varphi$, as expected.

## V. RESULTS

We here compare the conventional PWPIMC with our CSPIMC algorithm on specific simulations.

Our PWPIMC simulations (see Tables I, II and III, IV) confirm that at high temperature (classical regime) the nature of the statistics is not important. Whereas it becomes important at low temperatures (quantum regime). The zero temperature (ground state) limit can only be reached through an extrapolation of the PIMC results. In Tables I and II we use PWPIMC at constant $\tau = 0.025$ K$^{-1}$ (same as Ref. [10]) and various temperatures. In Tables III and V we compare the PWPIMC with the CSPIMC for Boltzmann statistics at fixed $M = 250$ and various temperatures. In Tables IV and VI we do the same for Bose statistics. We found favorable match between PWPIMC and CSPIMC results for $\varphi = 0.7 \approx 2^{-1/2}$. Each run has no less than $2 \times 10^6$ MC steps where one step is made of a displace move of all the timeslices of a single particle path and their associated ghost paths and a bridge move.

TABLE I. Results from **PW**PIMC for $N = 16$ $^4$He atoms ($m = 0.0830594$) in two dimensions, with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556, \varepsilon = 10.22$ and a cutoff distance $r_{\rm cut} = 2.5$ (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a density $N/L^2 = 0.05$ and various temperatures $T$ with $\tau = 0.025$, with **Boltzmann statistics**. In the Table $\mathscr{E}_K$ and $\mathscr{E}_P$ are the total kinetic and potential energies respectively. $\mathscr{E} = \mathscr{E}_K + \mathscr{E}_P$ is the total internal energy.

| $T$ (K) | $\mathscr{E}_K$ (K) | $-\mathscr{E}_P$ (K) | $-\mathscr{E}$ (K) |
|---|---|---|---|
| 1.0 | 66.6(2) | 94.4(1) | 27.8 |
| 0.5 | 60.8(2) | 92.9(1) | 32.1 |
| 0.2 | 59.0(2) | 93.00(8) | 34.0 |
| 0.1 | 58.2(2) | 93.23(8) | 35.0 |

In Tables VII and VIII we performed simulations at fixed $\varphi = 0.7$, and fixed temperatures $T = 1$ (K) and $T = 0.2$ (K) respectively, at increasing numbers $M$ of timeslices. We see that it is necessary to keep $\varphi$ constant upon taking the continuum limit in order to find agreement between the results for the CSPIMC and the ones for the PWPIMC for both the kinetic and potential energies. We find that, in the continuum $\tau \to 0$ limit, $\mu$ tends to $\approx 0.4$.

## VI. CONCLUSIONS

In this short work we proposed a possible realization of the Brownian bridge move in the CSPIMC algorithm presented in our recent work of Ref. [1]. As discussed in that work in the CSPIMC algorithm is necessary to take

TABLE II. Results from **PW**PIMC for $N = 16$ $^4$He atoms ($m = 0.0830594$) in two dimensions, with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556, \varepsilon = 10.22$ and a cutoff distance $r_{\rm cut} = 2.5$ (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a density $N/L^2 = 0.05$ and various temperatures $T$ with $\tau = 0.025$, with **Bose statistics**. In the Table $\mathscr{E}_K$ and $\mathscr{E}_P$ are the total kinetic and potential energies respectively. $\mathscr{E} = \mathscr{E}_K + \mathscr{E}_P$ is the total internal energy.

| $T$ (K) | $\mathscr{E}_K$ (K) | $-\mathscr{E}_P$ (K) | $-\mathscr{E}$ (K) |
|---|---|---|---|
| 1.0 | 57.1(1) | 94.8(3) | 37.7 |
| 0.5 | 55.0(5) | 93.4(2) | 38.4 |
| 0.2 | 55.8(1) | 93.55(5) | 37.7 |
| 0.1 | 55.9(1) | 93.46(5) | 37.6 |

TABLE III. Results from **PW**PIMC for $N = 16$ $^4$He atoms ($m = 0.0830594$) in two dimensions, with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556, \varepsilon = 10.22$ and a cutoff distance $r_{\rm cut} = 2.5$ (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a density $N/L^2 = 0.05$ and various temperatures $T$ with $M = 250$, with **Boltzmann statistics**. The runs are $2 \times 10^6$ steps long. In the Table $\mathscr{E}_K$ and $\mathscr{E}_P$ are the total kinetic and potential energies respectively. $\mathscr{E} = \mathscr{E}_K + \mathscr{E}_P$ is the total internal energy (note that unlike Ref. [10] we fix $M$ and not $\tau$).

| $T$ (K) | $\mathscr{E}_K$ (K) | $-\mathscr{E}_P$ (K) | $-\mathscr{E}$ (K) |
|---|---|---|---|
| 1.0 | 82.3(6) | 83.6(1) | 1.3(6) |
| 0.5 | 74.6(4) | 84.8(1) | 10.2(4) |
| 0.2 | 62.5(4) | 90.5(1) | 28.0(4) |
| 0.1 | 49.0(2) | 101.1(1) | 52.1(2) |

the continuum limit at fixed $\varphi = \xi\tau/m$ by letting $\tau \to 0$ and $\xi \to \infty$. We found favorable agreement between our CSPIMC and the conventional PWPIMC for $\varphi \approx 2^{-1/2}$.

## ACKNOWLEDGMENTS

I would like to thank prof. Saverio Moroni for his support in the development of the Brownian bridge move and the consequent particles permutation sampling.

TABLE IV. Results from **PW**PIMC for $N = 16$ $^4$He atoms ($m = 0.0830594$) in two dimensions, with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556, \varepsilon = 10.22$ and a cutoff distance $r_{\rm cut} = 2.5$ (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a density $N/L^2 = 0.05$ and various temperatures $T$ with $M = 250$, with **Bose statistics**. The runs are $2 \times 10^6$ steps long. In the Table $\mathscr{E}_K$ and $\mathscr{E}_P$ are the total kinetic and potential energies respectively. $\mathscr{E} = \mathscr{E}_K + \mathscr{E}_P$ is the total internal energy (note that unlike Ref. [10] we fix $M$ and not $\tau$).

| $T$ (K) | $\mathscr{E}_K$ (K) | $-\mathscr{E}_P$ (K) | $-\mathscr{E}$ (K) |
|---|---|---|---|
| 1.0 | 77(1) | 84.1(4) | 7.1(4) |
| 0.5 | 69.7(7) | 84.6(4) | 14.9(7) |
| 0.2 | 60.0(5) | 90.6(1) | 30.6(5) |
| 0.1 | 49.3(3) | 101.1(2) | 51.8(3) |

TABLE V. Results from **CS**PIMC with $\varphi = 0.7$ for $N = 16$ $^4$He atoms ($m = 0.0830594$) in two dimensions, with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556, \varepsilon = 10.22$ and a cutoff distance $r_{\rm cut} = 2.5$ (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a density $N/L^2 = 0.05$ and various temperatures $T$ with $M = 250$, with **Boltzmann statistics**. The runs are $2 \times 10^6$ steps long. In the Table $\mathscr{E}_K$ is taken from Table III and used to determine $\mu$. $\mathscr{E}_P$ is the total potential energy.

| $T$ (K) | $\langle \vec{\mathscr{P}}_k^2 + \vec{\mathscr{P}}_k'^2 \rangle / 4m$ (K) | $\mathscr{E}_K$ (K) | $-\mathscr{E}_P$ (K) | $\mu$ |
|---|---|---|---|---|
| 1.0 | 3550(6) | 82.3 | 84.4(4) | 0.44 |
| 0.5 | 1608(3) | 74.6 | 85.4(2) | 0.47 |
| 0.2 | 607(1) | 62.5 | 90.3(2) | 0.44 |
| 0.1 | 293.3(1) | 49.0 | 99.1(1) | 0.40 |

TABLE VI. Results from **CS**PIMC with $\varphi = 0.7$ for $N = 16$ $^4$He atoms ($m = 0.0830594$) in two dimensions, with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556, \varepsilon = 10.22$ and a cutoff distance $r_{\rm cut} = 2.5$ (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a density $N/L^2 = 0.05$ and various temperatures $T$ with $M = 250$, with **Bose statistics**. The runs are $2 \times 10^6$ steps long. In the Table $\mathscr{E}_K$ is taken from Table IV and used to determine $\mu$. $\mathscr{E}_P$ is the total potential energy.

| $T$ (K) | $\langle \vec{\mathscr{P}}_k^2 + \vec{\mathscr{P}}_k'^2 \rangle / 4m$ (K) | $\mathscr{E}_K$ (K) | $-\mathscr{E}_P$ (K) | $\mu$ |
|---|---|---|---|---|
| 1.0 | 3721(5) | 77 | 82.6(2) | 0.42 |
| 0.5 | 1611(3) | 69.7 | 86.4(2) | 0.48 |
| 0.2 | 606.8(8) | 60.0 | 89.8(1) | 0.44 |
| 0.1 | 293.0(1) | 49.3 | 100.42(6) | 0.40 |

## AUTHOR DECLARATIONS

### Conflicts of interest

None declared.

### Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

TABLE VII. Results from **CS**PIMC with $\varphi = 0.7$ for $N = 16$ $^4$He atoms ($m = 0.0830594$) in two dimensions, with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556, \varepsilon = 10.22$ and a cutoff distance $r_{\rm cut} = 2.5$ (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a density $N/L^2 = 0.05$ and a temperature $T = 0.2$ with various values of $M$, with **Boltzmann statistics**. In the Table $\mathscr{E}_K \approx 62.5$ is used to determine $\mu$. $\mathscr{E}_P$ is the total potential energy.

| $M$ | $\langle \vec{\mathscr{P}}_k^2 + \vec{\mathscr{P}}_k'^2 \rangle / 4m$ (K) | $\mu$ | $-\mathscr{E}_P$ (K) |
|---|---|---|---|
| 250 | 607(1) | 0.44 | 90.3(2) |
| 500 | 1268(1) | 0.47 | 86.7(1) |
| 750 | 2208(3) | 0.42 | 88.0(1) |
| 1000 | 4156(4) | 0.30 | 86.0(3) |

TABLE VIII. Results from **CS**PIMC with $\varphi = 0.7$ for $N = 16$ $^4$He atoms ($m = 0.0830594$) in two dimensions, with a Lennard-Jones pair potential with parameters [11] $\sigma = 2.556, \varepsilon = 10.22$ and a cutoff distance $r_{\rm cut} = 2.5$ (so that $v(r) = 0$ for $r > r_{\rm cut}\sigma$ without long range corrections), at a density $N/L^2 = 0.05$ and a temperature $T = 0.2$ with various values of $M$, with **Bose statistics**. In the Table $\mathscr{E}_K \approx 60.0$ is used to determine $\mu$. $\mathscr{E}_P$ is the total potential energy.

| $M$ | $\langle \vec{\mathscr{P}}_k^2 + \vec{\mathscr{P}}_k'^2 \rangle / 4m$ (K) | $\mu$ | $-\mathscr{E}_P$ (K) |
|---|---|---|---|
| 250 | 606.8(8) | 0.44 | 89.8(1) |
| 500 | 1272(1) | 0.47 | 86.2(1) |
| 750 | 2241(5) | 0.41 | 87.8(2) |
| 1000 | 4193(3) | 0.30 | 85.2(2) |

**Funding**

None declared.

[1] R. Fantoni, Coherent State Path Integral Monte Carlo, Eur. Phys. J. D **79**, 146 (2025).
[2] J. R. Klauder, Continuous-Representation Theory. I. Postulates of Continuous-Representation Theory, J. Math. Phys. **4**, 1055 (1963).
[3] J. R. Klauder, Continuous-Representation Theory. II. Generalized Relation between Quantum and Classical Dynamics, J. Math. Phys. **4**, 1058 (1963).
[4] J. R. Klauder, ContinuousRepresentation Theory. III. On Functional Quantization of Classical Systems, J. Math. Phys. **5**, 177 (1964).
[5] J. McKenna and J. R. Klauder, ContinuousRepresentation Theory. IV. Structure of a Class of Function Spaces Arising from Quantum Mechanics, J. Math. Phys. **5**, 878 (1964).
[6] J. R. Klauder and J. McKenna, ContinuousRepresentation Theory. V. Construction of a Class of Scalar Boson Field Continuous Representations, J. Math. Phys. **6**, 68 (1965).
[7] J. R. Klauder and B. Skagerstam, *Coherent States* (World Scientific, Singapore, 1985).
[8] D. M. Ceperley, Path integrals in the theory of condensed helium, Rev. Mod. Phys. **67**, 279 (1995).
[9] M. H. Kalos and P. A. Whitlock, *Monte Carlo Methods* (John Wiley & Sons Inc., New York, 1986).
[10] M. C. Gordillo and D. M. Ceperley, Path-integral calculation of the two-dimensional $^4$He phase diagram, Phys. Rev. B **58**, 6447 (1998).
[11] W. L. McMillan, Ground State of Liquid He$^4$, Phys. Rev. **138**, 442 (1964).
[12] H. F. Trotter, On the Product of Semi-Groups of Operators, Proc. Am. Math. Soc. **10**, 545 (1959).
[13] N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. M. Teller, and E. Teller, Equation of state calculations by fast computing machines, J. Chem. Phys. **1087**, 21 (1953).