

A Fourier-space four-field resonant response model for tokamak plasmas with applications to pedestal physics

R. Fitzpatrick* and Y. Lin

*Institute for Fusion Studies, Department of Physics,
University of Texas at Austin, Austin TX 78712, USA*

1 Introduction

1.1 Resonant magnetic perturbations

Externally generated, static (i.e., non-rotating in the laboratory frame), *resonant magnetic perturbations* (RMPs) can drive magnetic reconnection in tokamak plasmas that are intrinsically stable to tearing modes [1, 2, 3, 4, 5]. Static RMPs arise naturally as a consequence of magnetic field-coil misalignments [6], but are sometimes deliberately applied to tokamak plasmas in order to suppress edge-localized modes (ELMs) [7]. Driven reconnection is significant because it leads to the formation of magnetic island chains at so-called *rational* magnetic flux-surfaces within the plasma [8]. Such chains locally flatten the pressure profile, and, thereby, degrade the plasma energy confinement [9].

The analysis of the response of a tokamak plasma to an RMP is most efficiently formulated as an asymptotic matching problem in which the plasma is divided into two distinct regions [10]. In the so-called *outer region*, which comprises most of the plasma, the plasma perturbation is governed by the equations of linearized, marginally-stable, ideal-magnetohydrodynamics (MHD). However, these equations become singular on rational magnetic flux-surfaces at which the perturbed magnetic field resonates with the equilibrium field. In the *inner region*, which consists of a set of narrow layers centered on the various rational surfaces, non-ideal-MHD effects become important.

1.2 Four-field resonant response model

It is well known that single-fluid resistive-MHD offers a very poor description of the response of the inner region to the magnetic perturbation in the outer region. For instance, the strong diamagnetic flows present in tokamak plasmas imply that the electron and ion fluid velocities are significantly different from one another, necessitating a two-fluid treatment [11]. Moreover, resistive-MHD does not take into account the important ion sound-radius lengthscale below which electron and ion dynamics become decoupled from one another [12, 13]. Previously, Cole & Fitzpatrick [14] used the four-field model of Fitzpatrick & Waelbroeck [15], which consists of four coupled nonlinear partial differential equations, and which is based on the original four-field model of Hazeltine, Kotschenreuther & Morrison [16], to determine the linear two-fluid response of a resonant layer to the perturbation in the outer region. This treatment was extended in Ref. [17] in order to take into account the anomalously large perpendicular energy diffusivity present in tokamak plasmas.

In configuration space, the four-field models of Refs. [14] and [17] yield a set of resonant layer equations that can be expressed as *ten* coupled first-order linear ordinary differential equations (o.d.e.s) [18]. However, in Fourier space, the resonant layer equations can be written as *four* coupled first-order linear o.d.e.s [14]. It is clearly advantageous to solve the equations in Fourier space. In Refs. [14] and [17], an approximation is made

*rfitzp@utexas.edu

by which one of the terms in the Fourier-transformed layer equations is neglected. This approximation, which is valid in low- β plasmas, is such that the layer equation that governs the parallel ion dynamics decouples from the other three equations, effectively converting a four-field resonant response model into a three-field model. Furthermore, the three remaining layer equations can be combined to give a single second-order linear o.d.e. This second-order equation is most conveniently solved by means of a Riccati transformation that converts it into a first-order nonlinear o.d.e. [19, 20, 21]. The advantage of the Riccati approach is that it can deal with numerically problematic solutions that blow up as $\exp(p^2)$, or faster, at large p , where p is the Fourier-space variable.

Lee, Park & Na [18] recently demonstrated how to solve the full tenth-order four-field resonant layer equations in configuration space using a Riccati transformation. In the process, they discovered that, in a high- β plasma, the RMP rotation frequency (as seen in the local $\mathbf{E} \times \mathbf{B}$ frame at the rational surface) at which the electromagnetic torque exerted on the plasma in the immediate vicinity of the resonant layer passes through zero is shifted in the ion diamagnetic direction from the electron diamagnetic frequency. This result is significant because there is some experimental evidence for such a shift [22].

The first aim of this paper is to demonstrate how the calculation of Lee et alia can be reimplemented in Fourier space. The Fourier version of the calculation is more convenient, from a numerical point of view, because it involves the solution of a fourth-order, rather than a tenth-order, system of equations.

1.3 Magnetically diverted plasmas

The safety-factor profile of a magnetically diverted tokamak plasma diverges logarithmically as the magnetic separatrix (a.k.a. the last closed magnetic flux-surface) is approached. At first sight, this suggests that, when determining the response of such a plasma to an RMP, it is necessary to include an infinite number of rational magnetic flux-surfaces (at which the RMP resonates with the equilibrium magnetic field) in the calculation, the majority of which lie very close to the separatrix. In fact, as was demonstrated in Ref. [23], when finite plasma resistivity is taken into account, this turns out not to be the case. Instead, it is only necessary to include rational surfaces that lie in the region $0 < \Psi < 1 - \epsilon_c$, where Ψ is the normalized poloidal magnetic flux, and $0 < \epsilon_c \ll 1$. Moreover, the value of ϵ_c can be calculated from the plasma parameters at the edge of the discharge.

The analysis of Ref. [23] was based on the three-field resonant response model of Refs. [14] and [17]. The second aim of this paper is to generalize this analysis by employing the new Fourier-space four-field resonant response model.

2 Asymptotic matching

2.1 Plasma equilibrium

Consider a large aspect-ratio tokamak plasma equilibrium whose magnetic flux-surfaces map out (almost) concentric circles in the poloidal plane. Such an equilibrium can be approximated as a periodic cylinder [4]. Let r, θ, z be right-handed cylindrical coordinates. The magnetic axis corresponds to $r = 0$, and the plasma boundary to $r = a$, where a is the simulated minor radius of the plasma. The system is assumed to be periodic in the z -direction with periodicity length $2\pi R_0$, where R_0 is the simulated major radius of the plasma. The safety-factor profile takes the form $q(r) = r B_z / [R_0 B_\theta(r)]$, where B_z is the constant “toroidal” magnetic field-strength, and $B_\theta(r)$ is the poloidal magnetic field-strength. The equilibrium poloidal and toroidal magnetic fluxes (divided by 2π) are written $\psi_p(r) = B_z \int_0^r r' / q(r') dr'$ and $\psi_t(r) = B_z r^2 / 2$, respectively. The standard large aspect-ratio tokamak orderings, $r/R_0 \ll 1$ and $B_\theta/B_z \ll 1$, are adopted [21].

2.2 Outer region

Consider an externally generated, static (in the laboratory frame) magnetic perturbation that has m periods in the poloidal direction, and n periods in the toroidal direction. The response of the plasma to the perturbation is governed by the linearized equations of marginally-stable ideal-MHD everywhere in the plasma,

Regime	Q_e	Q_i	c_β	D	P_E	P_φ
HRi	1.0	-1.0	1.0	10.0	1.0	$1.0 \times 10^{+0}$
HRii	1.0	-1.0	1.0	10.0	1.0	1.0×10^{-1}
RIi	0.15	-0.15	1.0	1.0	1.0	1.0×10^{-4}
VRii	0.15	-0.15	1.0	0.1	1.0	$1.0 \times 10^{+0}$

Table 1: The normalized layer parameters used in the benchmark comparisons shown in Figs. 1–4.

apart from a (radially) narrow layer centered on the rational surface whose minor radius, r_s , is such that $q(r_s) = m/n$ [10].

The perturbed magnetic field is written $\delta \mathbf{B} \simeq \nabla \delta \psi \times \mathbf{e}_z$, where $\delta \psi(r, \theta, \varphi) = \delta \psi(r) \exp[i(m\theta - n\varphi)]$, and $\varphi = z/R_0$ is a simulated toroidal angle. In the outer region (i.e., everywhere in the plasma apart from the resonant layer), the perturbed helical magnetic flux, $\delta \psi(r)$, satisfies the *cylindrical tearing mode equation* [21, 24, 25],

$$\frac{d^2 \delta \psi}{dr^2} + \frac{1}{r} \frac{d \delta \psi}{dr} - \frac{m^2}{r^2} \delta \psi - \frac{J'_z \delta \psi}{r(1/q - n/m)} = 0, \quad (1)$$

where $J_z(r) = R_0 \mu_0 j_z(r)/B_z$, and $j_z(r)$ is the equilibrium “toroidal” current density. Here, $' \equiv d/dr$.

In general, the solution of Eq. (1) that satisfies physical boundary conditions at the magnetic axis and the plasma boundary is such that $\delta \psi(r)$ is continuous across the rational surface, whereas $d\delta \psi/dr$ is discontinuous. The discontinuity of $d\delta \psi/dr$ across the rational surface is indicative of the presence of a helical current sheet at the surface. The complex quantity $\Psi_s = \delta \psi(r_s)$ determines the amplitude and phase of the reconnected helical magnetic flux at the rational surface, whereas the complex quantity

$$\Delta \Psi_s = \left[r \frac{d \delta \psi}{dr} \right]_{r_{s-}}^{r_{s+}} \quad (2)$$

parameterizes the amplitude and phase of the helical current sheet [4].

The solution of the cylindrical tearing mode equation in the outer region, in the presence the RMP, leads to a *tearing mode dispersion relation* of the form [4, 10]

$$\Delta \Psi_s - E_{ss} \Psi_s = -E_{ss} \Psi_v, \quad (3)$$

where E_{ss} is a real dimensionless quantity known as the *tearing stability index* [10]. Moreover, Ψ_v is the so-called *vacuum flux*, and is defined as the reconnected magnetic flux that would be driven at the rational surface by the RMP were the plasma intrinsically tearing-stable (i.e., $E_{ss} < 0$), and were there no current sheet at the rational surface (i.e., $\Delta \Psi_s = 0$) [4].

In the following, it is assumed that $E_{ss} < 0$, which implies that the tearing mode that reconnects magnetic flux at the rational surface is intrinsically stable. In this case, any magnetic reconnection that takes place at the surface is entirely due to the RMP.

2.3 Inner region

The current sheet at the rational surface is resolved by employing a two-fluid resistive-MHD resonant response model in the inner region (i.e., the region of the plasma in the immediate vicinity of the rational surface) to determine the complex *layer response index*, Δ_s . Asymptotic matching between the solutions in the inner and the outer regions yields

$$\Delta_s = \frac{\Delta \Psi_s}{\Psi_s}. \quad (4)$$

The previous two equations lead to the *plasma response equation*,

$$\Psi_s = \frac{\Psi_v}{1 + \Delta_s/(-E_{ss})}. \quad (5)$$




Figure 1: The scaled layer response index, $\hat{\Delta}$, plotted as a function of the normalized $\mathbf{E} \times \mathbf{B}$ frequency, Q_E , in the first Hall-resistive (HRi) response regime. (See Table 1.) $\hat{\Delta}_3$ is the response index predicted by the three-field Fourier-space response model of Ref. [17]. $\hat{\Delta}_4$ is the response index predicted by the four-field Fourier-space response model specified in Appendix A. SLAYER is the response index predicted by the four-field configuration-space response model of Ref. [18]. The three vertical lines indicate the electron fluid resonance, $Q_E + Q_e = 0$, the MHD fluid resonance, $Q_E = 0$, and the ion fluid resonance, $Q_E + Q_i = 0$, respectively, in order from the left to the right.

Note that if $|\Delta_s|/(-E_{ss}) \ll 1$ then $\Psi_s \simeq \Psi_v$ and $\Delta\Psi_s \simeq 0$, which is termed a *vacuum response*. On the other hand, if $|\Delta_s|/(-E_{ss}) \gg 1$ then $|\Psi_s| \ll |\Psi_v|$, which is termed an *ideal response*. In the vacuum response regime, the current sheet that is excited in the resonant layer is too feeble to prevent driven magnetic reconnection, and the reconnected flux at the rational surface is the same as that which would be driven if there were no plasma present at the surface. On the other hand, in the ideal response regime, the current sheet excited in the layer is large enough to almost completely suppress driven magnetic reconnection, which implies that the response of the plasma is equivalent to that which would occur if the ideal-MHD flux freezing constraint, $\Psi_s = 0$, were imposed at the rational surface [23].

3 Fourier-space four-field resonant response model

3.1 Introduction

The Fourier-space four-field resonant response model employed in this paper is described in detail in Appendix A.




Figure 2: The scaled layer response index, $\hat{\Delta}$, plotted as a function of the normalized $\mathbf{E} \times \mathbf{B}$ frequency, Q_E , in the second Hall-resistive (HRii) response regime. See Table 1 and the caption to Fig. 1.

3.2 Dimensionless parameters

All four-field resonant response models are defined by the following seven dimensionless parameters: $Q_E = -S^{1/3} n \omega_E \tau_H$, $Q_e = -S^{1/3} n \omega_{*e} \tau_H$, $Q_i = -S^{1/3} n \omega_{*i} \tau_H$, $c_\beta = \sqrt{\beta/(1+\beta)}$, $D = S^{1/3} \iota_e^{1/2} \hat{d}_\beta$, $P_E = \tau_R/\tau_E$, and $P_\varphi = \tau_R/\tau_\varphi$. Here, τ_R is the resistive diffusion timescale, τ_H the hydrodynamic timescale, τ_E the energy confinement timescale, τ_φ the momentum confinement timescale, $S = \tau_R/\tau_H$ the Lundquist number, ω_E the $\mathbf{E} \times \mathbf{B}$ frequency, ω_{*e} the electron diamagnetic frequency, ω_{*i} the ion diamagnetic frequency, $\sqrt{\beta} = \sqrt{(5/3) \mu_0 p/B_z}$ the ratio of the sound speed to the Alfvén speed, p the plasma pressure, $d_\beta = c_\beta d_i$ the ion sound-radius, and d_i the collisionless ion skin-depth. Moreover, P_E and P_φ are magnetic Prandtl numbers, $\iota_e = -\omega_{*e}/(\omega_{*i} - \omega_{*e})$, and $\hat{d}_\beta = d_\beta/r_s$. All quantities are evaluated at the rational surface. (See Appendix A for more detailed definitions.)

Note that the previous definitions are slightly different to those adopted in Ref. [18]. We can convert our dimensionless parameter into those used by Lee et alia by letting $Q_E \rightarrow Q$, $Q_e \rightarrow -Q_{*e}$, $Q_i \rightarrow -Q_{*i}$, $\iota_e \rightarrow 1/(1+\tau)$, $P_E \rightarrow c_\beta^2$, and $P_\varphi \rightarrow P$.

3.3 Method of solution

After linearization and Fourier transformation (with respect to $x = r - r_s$), our four-field equations can be converted into the Riccati matrix differential equation

$$p \frac{d\mathbb{W}}{dp} = \mathbb{W} - \mathbb{W}\mathbb{W} - \mathbb{E}\mathbb{W} + \mathbb{F}. \quad (6)$$

Here, p is the Fourier transform variable, and $\mathbb{W}(p)$ the 2×2 solution matrix. The elements of the 2×2 matrices $\mathbb{E}(p)$ and $\mathbb{F}(p)$ are specified in Eqs. (98)–(101) and (102)–(105), respectively. The method of




Figure 3: The scaled layer response index, $\hat{\Delta}$, plotted as a function of the normalized $\mathbf{E} \times \mathbf{B}$ frequency, Q_E , in the second resistive-inertial (RIi) response regime. See Table 1 and the caption to Fig. 1.

solution is as follows. A solution of the Ricatti equation is launched from a large (with respect to unity) value of p , with the initial solution matrix specified in Eq. (162), and is then integrated to a small (with respect to unity) value of p . The *scaled layer response index*, $\hat{\Delta}$, is then deduced from the small- p elements of the solution matrix by means of Eq. (126). The true layer response index is $\Delta_s = S^{1/3} \hat{\Delta}$.

3.4 Benchmarking against SLAYER

Our first task is to benchmark the results of our Fourier-space calculation of $\hat{\Delta}$ against the results of the configuration-space calculation performed by Lee et alia. The latter calculation has been implemented in the SLAYER code [18, 20].

Figures 1–4 show the scaled layer response index, $\hat{\Delta}$, calculated as a function of the normalized $\mathbf{E} \times \mathbf{B}$ frequency, Q_E , by three different solution methods, in four of the analytic layer response regimes identified in Ref. [14]. The dimensionless layer parameters used in the four calculations are specified in Table 1. The first solution method involves using the Fourier-space analysis of Ref. [17]. In this analysis, the term involving c_β^2 is neglected in Eq. (53). Consequently, Eq. (55) decouples from Eqs. (52)–(54), which effectively converts a four-field response model (the four fields being $\bar{\psi}$, \bar{N} , $\bar{\phi}$, and \bar{V}) into a three-field response model (the three fields being $\bar{\psi}$, \bar{N} , and $\bar{\phi}$). The second solution method involves using the Fourier-space analysis set out in Appendix A. This method solves Eqs. (52)–(55), with no neglected terms, and thus constitutes a four-field response model. The first two methods have been implemented in the TJ code [26, 27]. The third solution method involves using the configuration-space analysis of Ref. [18], which is implemented in the SLAYER code [20]. Solution methods two and three constitute equivalent four-field response models.

It is clear from Figs. 1–4 that the second and third solution methods yield exactly the same results. This is remarkable because the second method involves solving four coupled first-order o.d.e.s in Fourier




Figure 4: The scaled layer response index, $\hat{\Delta}$, plotted as a function of the normalized $\mathbf{E} \times \mathbf{B}$ frequency, Q_E , in the second viscous-resistive (VRii) response regime. See Table 1 and the caption to Fig. 1.

space, whereas the third involves solving ten coupled first-order o.d.e.s in configuration space. Moreover, as is clear from a comparison of Appendix A and Ref. [18], the analysis that leads to the four coupled equations in the former method is radically different from, and completely independent of, that which leads to the ten coupled equations in the latter method. It follows that the two solution methods constitute strong independent checks on one another. Hence, we can be confident that the two methods are valid, and also that they have been implemented correctly in the TJ and SLAYER codes.

3.5 Effect of c_β

In Figs. 1–4, the *electron fluid resonance*, $Q_E + Q_e = 0$, the *MHD-fluid resonance*, $Q_E = 0$, and the *ion fluid resonance*, $Q_E + Q_i = 0$, are highlighted. The electron fluid resonance corresponds to the situation in which the rotation frequency of the electron fluid at the rational surface passes through zero, which we would expect to maximize the interaction between the electron fluid and the static RMP. Likewise for the other fluid resonances.

Figures 1–4 allow us to assess the consequences of neglecting the term involving c_β^2 in Eq. (53) by looking at the difference between $\hat{\Delta}_4$ and $\hat{\Delta}_3$. Obviously, this neglect is unlikely to make a significant difference if the ion sound parameter, c_β , is much less than unity, which explains why the rather large value, $c_\beta = 1$, is used in the calculations shown in the figures. (See Table 1.) It can be seen from the figures that the neglect of the c_β^2 term has a larger relative effect on the real part of the scaled layer response index, $\hat{\Delta}$, than on the imaginary part. Furthermore, Figs. 2 and 3 indicate that if the c_β^2 term is neglected then the plasma response model fails to capture resonant behavior that occurs, in some response regimes, close to the MHD-fluid resonance ($Q_E = 0$). All four figures indicate that if the c_β^2 term is neglected then the real and the imaginary parts of the response index pass simultaneously through zero at the electron fluid resonance ($Q_E + Q_e = 0$). From




Figure 5: The difference between the scaled layer response indices predicted by the three-field and the four-field Fourier-space resonant response models plotted as a function of the normalized $\mathbf{E} \times \mathbf{B}$ frequency, Q_E . The calculations are performed in the VRii regime of Table 1, but for various different values of c_β . See the caption to Fig. 1.

Eq. (5), this implies a complete loss of shielding at the electron fluid resonance, leading to a vacuum-like response in which the reconnected magnetic flux driven at the rational surface is maximized. The figures also indicate that if the c_β^2 term is retained then the value of Q_E at which the imaginary part of the scaled response index passes through zero shifts from the electron fluid resonance in the direction of the ion fluid resonance. The value of Q_E at which the real part of the scaled response index passes through zero also shifts from the electron fluid resonance. However, the direction of the shift is different in different response regimes.

Figure 5 shows the difference between the scaled layer response index predicted by the three-field (i.e., $c_\beta^2 = 0$) Fourier-space resonant response model of Ref. [17] and the four-field Fourier-space resonant response model of Appendix A. The calculations are performed using the Q_e , Q_i , D , P_E , and P_φ values associated with the VRii regime of Table 1, and for various different values of c_β . It can be seen that the difference between the predictions of the three-field and four-field models is an increasing function of increasing c_β , and that the difference is essentially negligible for c_β values less than 0.01.

Figure 6 shows the critical value of the normalized electron fluid rotation frequency at the rational surface, $Q_E + Q_e$, at which the real and imaginary parts of the scaled layer response index, $\hat{\Delta}$, pass through zero, plotted as a function of c_β . The calculations are performed using the Q_e , Q_i , D , P_E , and P_φ values associated with the HRi, HRii, RIi, and VRii regimes of Table 1. As we have seen, if $c_\beta = 0$ then the real and the imaginary components of $\hat{\Delta}$ pass through zero simultaneously as the electron fluid rotation frequency passes through zero. However, as is clear from the figure, as c_β increases from zero, the critical value of the electron fluid rotation frequency at which the real part of the scaled layer response index passes through zero shifts in the electron diamagnetic direction (i.e., to more negative values of $Q_E + Q_e$), whereas that at which the




Figure 6: The critical value of the normalized electron fluid rotation frequency at the rational surface, $Q_E + Q_e$, at which the real (dashed curves) and imaginary (solid curves) components of the layer response index pass through zero, plotted as a function of c_β for the four response regimes specified in Table 1.

imaginary part passes through zero shifts in opposite direction. The only exception to this rule occurs in the second resistive-inertial (RII) regime, where the critical values of the electron fluid rotation frequency at which the real and imaginary parts of the scaled layer response index pass through zero both shift in the ion diamagnetic direction by almost equal amounts. According to Eq. (5), a vacuum response (i.e., $\Psi \simeq \Psi_v$) requires the real and the imaginary parts of the scaled plasma response index to be simultaneously much smaller than unity (given that the true layer response index is $\Delta_s = S^{1/3} \hat{\Delta}$, whereas, under normal circumstances, $-E_{ss} \sim \mathcal{O}(1)$, and $S \gg 1$). As c_β increases, this state of affairs seems to become progressively harder to achieve, except in the RII regime.

4 Response of the pedestal region of tokamak plasma to an RMP

4.1 Edge-localized modes

Magnetically diverted tokamak plasmas generally operate in so-called *H-mode* [28]. In this operation mode, a transport barrier forms in the outer part of the plasma, leading to a region of enhanced density and temperature gradients known as the *pedestal*. As the name suggests, the “pedestal” is characterized by negative steps in the density and temperature profiles that are located just inside the plasma boundary. (See Fig. ??.) Unfortunately, the large pressure gradient present in the pedestal can drive ideal-MHD instabilities known as *peeling-ballooning modes* [29]. Consequently, most H-mode plasmas exhibit intermittent bursts of heat and particle transport, emanating from the pedestal, that are known as (type-I) *edge-localized modes* (ELMs) [30]. It is estimated that the heat load that ELMs will deliver to the tungsten plasma-facing components in a reactor-scale tokamak will be large enough to cause massive tungsten ion influx into the

ν_q	3.5	ν_{n_e}	1.1	ν_{T_e}	2.0	ν_{T_i}	2.0
q_0	1.01	n_{e0}	$1.2 \times 10^{20} \text{ m}^{-3}$	T_{e0}	$1.6 \times 10^4 \text{ eV}$	T_{i0}	$1.4 \times 10^4 \text{ eV}$
α_q	0.4	n_{e1}	$2.0 \times 10^{19} \text{ m}^{-3}$	T_{e1}	$0.0 \times 10^0 \text{ eV}$	T_{i1}	$2.5 \times 10^1 \text{ eV}$
Δ_q	0.6	Δ_{n_e}	$9.0 \times 10^{19} \text{ m}^{-3}$	Δ_{T_e}	$4.0 \times 10^3 \text{ eV}$	Δ_{T_i}	$4.0 \times 10^3 \text{ eV}$
\hat{r}_q	0.940	\hat{r}_{n_e}	0.950	\hat{r}_{T_e}	0.955	\hat{r}_{T_i}	0.950
δ_q	0.025	δ_{n_e}	0.025	δ_{T_e}	0.025	δ_{T_i}	0.025

Table 2: Parameters used to model the ITER pedestal.

plasma core, and that the erosion associated with this process will unacceptably limit the lifetimes of these components [31]. Consequently, the development of robust and effective methods for ELM control is a high priority for the international magnetic fusion program.

4.2 RMP-induced ELM suppression

The most promising method for the control of ELMs is via the application of externally generated, static, RMPs to the plasma. Complete RMP-induced ELM suppression was first demonstrated on the DIII-D tokamak [7]. Subsequently, either mitigation or complete suppression of ELMs via RMPs has been demonstrated on the JET [32], ASDEX-U [33], KSTAR, [34] MAST [35], EAST [36], and MAST-U [37] tokamaks.

If an RMP opens up a magnetic island chain at the foot of the pedestal then it gives rise to an undesirable phenomenon, known as a *density pump-out*, by which the density gradient in the pedestal is greatly reduced [38, 39, 40]. On the other hand, if an RMP opens up a magnetic island chain at the top of the pedestal then it generally suppresses ELMs [38, 39, 40, 41, 42], because the island chain limits the width and height of the pedestal, and, thereby, prevents the peeling-ballooning instability threshold from being crossed [29].

In the following, we shall construct a simple model of a pedestal in the ITER tokamak, and then use our Fourier-space, four-field, resonant response model to gauge the response of this model pedestal to a externally generated, static, RMP.

4.3 Model ITER pedestal

We shall again employ the cylindrical analysis of Sect. 2. Let the normalized toroidal plasma current take the form

$$J_z(\hat{r}) = \frac{2}{q_0} (1 - \hat{r}^2)^{\nu_q - 1}, \quad (7)$$

where $\hat{r} = r/a$. The model safety-factor profile is written

$$q(\hat{r}) = \frac{\nu_q q_0 \hat{r}^2}{1 - (1 - \hat{r}^2)^{\nu_q}} - \alpha_q \ln(1 - \hat{r}^2) - \frac{\Delta_q}{2} \left[1 + \tanh \left(\frac{\hat{r} - \hat{r}_q}{\delta_q} \right) \right]. \quad (8)$$

The first term on the right-hand side of the previous equation is the safety-factor profile that cylindrical theory would associate with the current profile specified in Eq. (7) [25]. Here, q_0 is the cylindrical safety-factor on the magnetic axis ($\hat{r} = 0$), whereas $\nu_q q_0$ is the cylindrical safety-factor at the plasma boundary ($\hat{r} = 1$). The second term is intended to mimic the logarithmic singularity in the q -profile that occurs at the edge of a magnetically diverted plasma (assuming that the magnetic separatrix corresponds to $\hat{r} = 1$) [23]. The final term is meant to represent the reduction in the magnetic shear generated by the strong, pressure-gradient-driven, bootstrap current [43] present in the pedestal of an H-mode plasma. (See, for example, Fig. 6 of Ref. [44] and Fig. 1 of Ref. [45].) The approach taken here is similar to that taken in the TM1 code, which employs cylindrical geometry in association with experimental safety-factor, density, and temperature profiles [46, 47]. This approach has been very successful at explaining RMP-induced density pump-out and ELM suppression in the DIII-D tokamak [38, 39, 40].




Figure 7: Safety-factor, magnetic shear, electron number density, and electron and ion temperature profiles used to model ITER pedestal.

The model electron number density, electron temperature, and ion temperature profiles take the respective forms

$$n_e(\hat{r}) = (n_{e0} - n_{e1})(1 - \hat{r}^2)^{\nu_{n_e}} + \frac{\Delta n_e}{2} \left[1 + \tanh \left(\frac{\hat{r} - \hat{r}_{n_e}}{\delta_{n_e}} \right) \right] + n_{e1}, \quad (9)$$

$$T_e(\hat{r}) = (T_{e0} - T_{e1})(1 - \hat{r}^2)^{\nu_{T_e}} + \frac{\Delta T_e}{2} \left[1 + \tanh \left(\frac{\hat{r} - \hat{r}_{T_e}}{\delta_{T_e}} \right) \right] + T_{e1}, \quad (10)$$

$$T_i(\hat{r}) = (T_{i0} - T_{i1})(1 - \hat{r}^2)^{\nu_{T_i}} + \frac{\Delta T_i}{2} \left[1 + \tanh \left(\frac{\hat{r} - \hat{r}_{T_i}}{\delta_{T_i}} \right) \right] + T_{i1}. \quad (11)$$

Here, the second terms on the right-hand sides of the previous three equations represent the negative steps in the three profiles in the pedestal region.

The parameters used in our model ITER pedestal are specified in Table 2. The electron number density and electron and ion temperature profiles approximately reproduce the model profiles shown in Fig. 8 of Ref. [48]. Our model pedestal is characterized by $q_{95} = 3.70$, $n_e(1) = 2.2 \times 10^{19} \text{ m}^{-3}$, $T_e(1) = 106 \text{ eV}$, and $T_i(1) = 97 \text{ eV}$. The magnetic shear attains its minimum value of $s = 0.21$ at $\hat{r} = 0.937$.

The parallel electrical resistivity of the plasma is [21]

$$\eta_{\parallel} = \frac{0.51 \ln A Z m_e^{1/2} e^{1/2}}{6\sqrt{2} \pi^{3/2} f_p \epsilon_0^2 T_e^{3/2}} \quad (12)$$




Figure 8: Rotation frequency profiles in the model ITER pedestal. Here, ω_{*e} is the electron diamagnetic frequency, ω_{*i} the ion diamagnetic frequency, ω_E the $\mathbf{E} \times \mathbf{B}$ frequency, and $\omega_{\perp e} = \omega_E + \omega_{*e}$ the electron fluid rotation frequency. All frequencies are in radians per second.

where

$$\ln \Lambda = 24 - \ln \left[\left(\frac{n_e}{10^6} \right)^{1/2} \frac{1}{T_e} \right], \quad (13)$$

$$f_p = 1 - 1.46 \left(\frac{\hat{r} a}{R_0} \right)^{1/2} + 0.46 \left(\frac{\hat{r} a}{R_0} \right)^{3/2}. \quad (14)$$

Here, n_e is measured in SI units, whereas T_e is measured in units of eV. Moreover, m_e is the electron mass, e the magnitude of the electron charge, $Z = 3.0$ the effective ion charge number, $R_0 = 5.3$ m the plasma major radius, and $a = 2.0$ m the plasma minor radius [49]. The factor f_p accounts for the neoclassical enhancement of resistivity [50].

All of the remaining plasma parameter, apart from the $\mathbf{E} \times \mathbf{B}$ rotation profile, can be calculated from their definitions given in Appendix A, the profile data specified in Eqs. (8)–(14) and Table 2, and the following information. The vacuum toroidal magnetic field-strength is $B_0 = 5.3$ T [49], the ion mass number is $M = 2$, and the perpendicular energy and momentum diffusivities take the (spatially) constant values $\chi_E = 1$ m²/s and $\chi_\varphi = 1$ m²/s, respectively [51].

Figure 8 shows the rotation frequencies in our model ITER pedestal. The adopted $\mathbf{E} \times \mathbf{B}$ frequency profile is written

$$\omega_E(\hat{r}) = \omega_{E0} (1 - \hat{r}^2)^{\nu_E} + \alpha_E \omega_{*e}, \quad (15)$$

where $\omega_{E0} = 3 \times 10^4$ rad./s, $\nu_E = 1.1$, and $\alpha_E = 0.3$. The adopted $\mathbf{E} \times \mathbf{B}$ frequency profile has the appropriate sign in the pedestal and the core [45, 48], and is designed to ensure that the electron fluid rotation frequency, $\omega_{\perp e} = \omega_E + \omega_{*e}$, passes through zero close to the top of the pedestal. This feature of the electron fluid

rotation frequency is almost always observed in H-mode tokamak plasmas with co-current neutral beam injection, and has been identified as a key feature in RMP-induced ELM suppression [38, 39, 40, 41, 42].

Acknowledgements

The authors would like to thank Y. Lee, J.-K. Park, J. Waybright, and D. Burgess for help benchmarking the results of this paper against those of Ref. [18].

Funding

This research was supported by the U.S. Department of Energy, Office of Science, Office of Fusion Energy Sciences, under contract DE-FG02-04ER54742.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

A Fourier-space four-field resonant plasma response model

A.1 Fundamental definitions

The plasma is assumed to consist of two species. First, electrons of mass m_e , electrical charge $-e$, number density n_e , and temperature T_e . Second, ions of mass m_i , electrical charge $+e$, number density n_i , and temperature T_i . Let $p = n_e (T_e + T_i)$ be the total plasma pressure.

Let r_s be the minor radius of the rational surface. It is helpful to define $n_0 = n_e(r_s)$, $p_0 = p(r_s)$, [21]

$$\eta_e = \left. \frac{d \ln T_e}{d \ln n_e} \right|_{r=r_s}, \quad (16)$$

$$\eta_i = \left. \frac{d \ln T_i}{d \ln n_e} \right|_{r=r_s}, \quad (17)$$

$$\iota = \left(\frac{T_e}{T_i} \right)_{r=r_s} \left(\frac{1 + \eta_e}{1 + \eta_i} \right) = \left(\frac{dp_e}{dp_i} \right)_{r_s}, \quad (18)$$

where $n_e(r)$, $p(r)$, $p_e(r)$, $p_i(r)$, $T_e(r)$, and $T_i(r)$ refer to electron number density, total pressure, electron pressure, ion pressure, electron temperature, and ion temperature profiles, respectively, in the absence of the magnetic perturbation.

For the sake of simplicity, the perturbed electron and ion temperature profiles are assumed to be functions of the perturbed electron number density profile in the immediate vicinity of the rational surface. In other words, $T_e = T_e(n_e)$ and $T_i = T_i(n_e)$. This implies that $p = p(n_e)$. The “MHD velocity”, which is the velocity of a fictional MHD fluid [52], is defined $\mathbf{V} = \mathbf{V}_E + V_{\parallel i} \mathbf{b}$, where \mathbf{V}_E is the $\mathbf{E} \times \mathbf{B}$ drift velocity, $V_{\parallel i}$ is the parallel component of the ion fluid velocity, $\mathbf{b} = \mathbf{B}/|\mathbf{B}|$, and \mathbf{B} is the magnetic field-strength.

A.2 Fundamental fields

The four fundamental fields in our four-field model—namely, ψ , N , ϕ , and V —have the following definitions [21]:

$$\nabla\psi = \frac{\mathbf{e}_{\parallel} \times \mathbf{B}}{r_s B_z}, \quad (19)$$

$$N = -\hat{d}_i \left(\frac{p - p_0}{B_z^2 / \mu_0} \right), \quad (20)$$

$$\nabla\phi = \frac{\mathbf{e}_{\parallel} \times \mathbf{V}}{r_s V_A}, \quad (21)$$

$$V = \hat{d}_i \left(\frac{\mathbf{e}_{\parallel} \cdot \mathbf{V}}{V_A} \right). \quad (22)$$

Here, $\mathbf{e}_{\parallel} = (0, \epsilon/q_s, 1)$, $\epsilon = r/R_0$, $q_s = m/n$, $V_A = (B_z / \sqrt{\mu_0 n_0 m_i})_{r_s}$, $d_i = \sqrt{m_i / (n_0 e^2 \mu_0)}_{r_s}$, and $\hat{d}_i = d_i / r_s$. Our model also employs the auxiliary field

$$J = -\frac{2\epsilon_s}{q_s} + \hat{\nabla}^2 \psi, \quad (23)$$

where $\epsilon_s = r_s / R_0$, and $\hat{\nabla} = r_s \nabla$. Note that V_A is the *Alfvén speed* at the rational surface, whereas d_i is the *collisionless ion skin-depth*.

A.3 Fundamental equations

For the case of a static (in the laboratory frame) magnetic perturbation, the four-field model takes the form [14, 15, 17, 21]:

$$0 = [\phi, \psi] - \iota_e [N, \psi] + \hat{\eta}_{\parallel} J + \hat{E}_{\parallel}, \quad (24)$$

$$0 = [\phi, N] + \hat{d}_{\beta}^2 [J, \psi] + c_{\beta}^2 [V, \psi] + \hat{\chi}_E \hat{\nabla}^2 N, \quad (25)$$

$$0 = [\phi, \hat{\nabla}^2 \phi] - \frac{\iota_i}{2} \left(\hat{\nabla}^2 [\phi, N] + [\hat{\nabla}^2 \phi, N] + [\hat{\nabla}^2 N, \phi] \right) + [J, \psi] + \hat{\chi}_{\varphi} \hat{\nabla}^4 (\phi + \iota_i N), \quad (26)$$

$$0 = [\phi, V] + [N, \psi] + \hat{\chi}_{\varphi} \hat{\nabla}^2 V. \quad (27)$$

Here, $[A, B] \equiv \hat{\nabla} A \times \hat{\nabla} B \cdot \mathbf{e}_{\parallel}$, $\iota_e = \iota / (1 + \iota)$, $\iota_i = 1 / (1 + \iota)$, $\hat{t} = t / (r_s / V_A)$, $\hat{\eta}_{\parallel} = \eta_{\parallel} / (\mu_0 r_s V_A)$, $\hat{E}_{\parallel} = E_{\parallel} / (B_z V_A)$, $\hat{\chi}_E = \chi_E / (r_s V_A)$, $\hat{\chi}_{\varphi} = \chi_{\varphi} / (r_s V_A)$, where η_{\parallel} is the parallel plasma electrical resistivity at the rational surface, E_{\parallel} the parallel inductive electric field that maintains the equilibrium toroidal plasma current in the vicinity of the rational surface, χ_E the anomalous perpendicular heat diffusivity at the rational surface, and χ_{φ} the anomalous perpendicular ion momentum diffusivity at the rational surface. Moreover, $d_{\beta} = c_{\beta} d_i$, and $\hat{d}_{\beta} = d_{\beta} / r_s$, where $c_{\beta} = [\beta / (1 + \beta)]^{1/2}_{r_s}$, and $\beta = (5/3) \mu_0 p_0 / B_z^2$. Here, d_{β} is usually referred to as the *ion sound-radius*.

A.4 Matching to plasma equilibrium

The unperturbed plasma equilibrium is such that $\mathbf{B} = (0, B_{\theta}(r), B_z)$, $p = p(r)$, $\mathbf{V} = (0, V_E(r), V_z(r))$, where $V_E(r) \simeq -E_r / B_z$ is the (dominant θ -component of the) $\mathbf{E} \times \mathbf{B}$ velocity. Now, the resonant layer is assumed to have a radial thickness that is much smaller than r_s . Hence, we only need to evaluate plasma equilibrium quantities in the immediate vicinity of the rational surface. Equations (19)–(22) suggest that

[21]

$$\psi(\hat{x}) = \frac{\hat{x}^2}{2\hat{L}_s}, \quad (28)$$

$$N(\hat{x}) = -\hat{V}_* \hat{x}, \quad (29)$$

$$\phi(\hat{x}) = -\hat{V}_E \hat{x}, \quad (30)$$

$$V(\hat{x}) = \hat{V}_\parallel, \quad (31)$$

where $\hat{x} = (r-r_s)/r_s$, $\hat{L}_s = L_s/r_s$, $L_s = R_0 q_s/s_s$, $\hat{V}_E = V_E(r_s)/V_A$, $\hat{V}_* = V_*(r_s)/V_A$, $V_*(r) = (dp/dr)/(e n_0 B_z)$ is the (dominant θ -component of the) diamagnetic velocity, and $\hat{V}_\parallel = \hat{d}_i V_z(r_s)/V_A$. Here, $s_s = s(r_s)$ and $s(r) = d \ln q / d \ln r$. We also have

$$J(\hat{x}) = -\left(\frac{2}{s_s} - 1\right) \frac{1}{\hat{L}_s}, \quad (32)$$

and $\hat{E}_\parallel(\hat{x}) = (2/s_s - 1)(\hat{\eta}_\parallel/\hat{L}_s)$.

A.5 Derivation of linear layer equations

In accordance with Eqs. (28)–(32), let us write [21]

$$\psi(\hat{x}, \zeta) = \frac{\hat{x}^2}{2\hat{L}_s} + \tilde{\psi}(\hat{x}) e^{i\zeta}, \quad (33)$$

$$\phi(\hat{x}, \zeta) = -\hat{V}_E \hat{x} + \tilde{\phi}(\hat{x}) e^{i\zeta}, \quad (34)$$

$$N(\hat{x}, \zeta) = -\hat{V}_* \hat{x} + \iota_e \tilde{N}(\hat{x}) e^{i\zeta}, \quad (35)$$

$$V(\hat{x}, \zeta) = \hat{V}_\parallel + \iota_e \tilde{V}(\hat{x}) e^{i\zeta}, \quad (36)$$

$$J(\hat{x}, \zeta) = -\left(\frac{2}{s_s} - 1\right) \frac{1}{\hat{L}_s} + \hat{\nabla}^2 \tilde{\psi}(\hat{x}) e^{i\zeta}, \quad (37)$$

where $\zeta = m\theta - n\varphi$. Substituting Eqs. (33)–(37) into Eqs. (23)–(27), and only retaining terms that are first order in perturbed quantities, we obtain the following set of linear equations:

$$-i n (\omega_E + \omega_{*e}) \tau_H \tilde{\psi} = -i \hat{x} (\tilde{\phi} - \tilde{N}) + S^{-1} \hat{\nabla}^2 \tilde{\psi}, \quad (38)$$

$$-i n \omega_E \tau_H \tilde{N} = i n \omega_{*e} \tau_H \tilde{\phi} - i \iota_e \hat{d}_\beta^2 \hat{x} \hat{\nabla}^2 \tilde{\psi} - i c_\beta^2 \hat{x} \tilde{V} + S^{-1} P_E \hat{\nabla}^2 \tilde{N}, \quad (39)$$

$$-i n (\omega_E + \omega_{*i}) \tau_H \hat{\nabla}^2 \tilde{\phi} = -i \hat{x} \hat{\nabla}^2 \tilde{\psi} + S^{-1} P_\varphi \hat{\nabla}^4 \left(\tilde{\phi} + \frac{\tilde{N}}{\iota} \right), \quad (40)$$

$$-i n \omega_E \tau_H \tilde{V} = -i n \omega_{*e} \tau_H \tilde{\psi} - i \hat{x} \tilde{N} + S^{-1} P_\varphi \hat{\nabla}^2 \tilde{V}. \quad (41)$$

Here, $\tau_H = L_s/(m V_A)$ is the hydromagnetic time, $\omega_E = -(q_s/r_s) V_E(r_s) = -(d\Phi/d\psi_p)_{r_s}$ the $\mathbf{E} \times \mathbf{B}$ frequency, $\omega_{*e} = \iota_e (q_s/r_s) V_*(r_s) = [(dp_e/d\psi_p)/(e n_e)]_{r_s}$ the electron diamagnetic frequency, $\omega_{*i} = -\iota_i (m/r_s) V_*(r_s) = -[(dp_i/d\psi_p)/(e n_e)]_{r_s}$ the ion diamagnetic frequency, $\Phi(r)$ the equilibrium electrostatic potential, $S = \tau_R/\tau_H$ the Lundquist number, $\tau_R = \mu_0 r_s^2/\eta_\parallel$ the resistive diffusion time, $\tau_E = r_s^2/\chi_E$ the energy confinement time, and $\tau_\varphi = r_s^2/\chi_\varphi$ the toroidal momentum confinement time. Furthermore, $P_E = \tau_R/\tau_E$ and $P_\varphi = \tau_R/\tau_\varphi$ are magnetic Prandtl numbers.

Let us define the stretched radial variable $X = S^{1/3} \hat{x}$. Assuming that $X \sim \mathcal{O}(1)$ in the layer (i.e., assuming that the layer thickness is roughly of order $S^{-1/3} r_s$), and making use of the fact that $S \gg 1$ in

conventional tokamak plasmas, Eqs. (38)–(41) reduce to the following set of linear layer equations [14, 17]:

$$\mathrm{i}(Q_E + Q_e) \tilde{\psi} = -\mathrm{i}X \left(\tilde{\phi} - \tilde{N} \right) + \frac{d^2 \tilde{\psi}}{dX^2}, \quad (42)$$

$$\mathrm{i}Q_E \tilde{N} = -\mathrm{i}Q_e \tilde{\phi} - \mathrm{i}D^2 X \frac{d^2 \tilde{\psi}}{dX^2} - \mathrm{i}c_\beta^2 X \tilde{V} + P_E \frac{d^2 \tilde{N}}{dX^2}, \quad (43)$$

$$\mathrm{i}(Q_E + Q_i) \frac{d^2 \tilde{\phi}}{dX^2} = -\mathrm{i}X \frac{d^2 \tilde{\psi}}{dX^2} + P_\varphi \frac{d^4}{dX^4} \left(\tilde{\phi} + \frac{\tilde{N}}{\iota} \right), \quad (44)$$

$$\mathrm{i}Q_E \tilde{V} = \mathrm{i}Q_e \tilde{\psi} - \mathrm{i}X \tilde{N} + P_\varphi \frac{d^2 \tilde{V}}{dX^2}. \quad (45)$$

Here, $Q_E = -S^{1/3} n \omega_E \tau_H$, $Q_{e,i} = -S^{1/3} n \omega_{*e,i} \tau_H$, and $D = S^{1/3} \iota_e^{1/2} \hat{d}_\beta$.

If we write $P_E = c_\beta^2$ then Eqs. (42)–(45) become equivalent to the set of layer equations solved by Lee et alia. (To be slightly more exact, we get from our equations to those of Lee et alia by making the following transformation: $Q_E \rightarrow Q$, $Q_e \rightarrow -Q_{*e}$, $Q_i \rightarrow -Q_{*i}$, $\iota \rightarrow 1/\tau$, $P_E \rightarrow c_\beta^2$, $P_\varphi \rightarrow P$, $\tilde{\psi} \rightarrow -\tilde{\psi}$, $\tilde{N} \rightarrow \tilde{Z}$, $\tilde{\phi} \rightarrow \tilde{\phi}$, and $\tilde{V} \rightarrow -\tilde{V}_z$.)

The low- β approximation used in Refs. [14] and [17] involves neglecting the term containing c_β^2 in Eq. (43). This approximation decouples Eq. (45) from the three preceding equations, and effectively converts a four-field resonant response model into a three-field model. In the following, we shall not make use of this approximation.

A.6 Asymptotic matching

The linear layer equations, (42)–(45), possess tearing parity solutions characterized by the symmetry $\tilde{\psi}(-X) = \tilde{\psi}(X)$, $\tilde{N}(-X) = -\tilde{N}(X)$, $\tilde{\phi}(-X) = -\tilde{\phi}(X)$, $\tilde{V}(-X) = \tilde{V}(X)$. As is easily demonstrated, the asymptotic behavior of the tearing parity solutions to Eqs. (42)–(45) are such that

$$\tilde{\psi}(X) \rightarrow \psi_0 \left[\frac{\hat{\Delta}}{2} |X| + 1 + \mathcal{O}\left(\frac{1}{X^2}\right) \right], \quad (46)$$

$$\tilde{\phi}(X) \rightarrow -\psi_0 Q_E \left[\frac{\hat{\Delta}}{2} \operatorname{sgn}(X) + \frac{1}{X} + \mathcal{O}\left(\frac{1}{X^2}\right) \right], \quad (47)$$

$$\tilde{N}(X) \rightarrow \psi_0 Q_e \left[\frac{\hat{\Delta}}{2} \operatorname{sgn}(X) + \frac{1}{X} + \mathcal{O}\left(\frac{1}{X^2}\right) \right], \quad (48)$$

$$\tilde{V}(X) \rightarrow \mathcal{O}\left(\frac{1}{X^3}\right) \quad (49)$$

as $|X| \rightarrow \infty$, where ψ_0 is an arbitrary constant. It follows from Eq. (4) that layer response index is

$$\Delta_s = S^{1/3} \hat{\Delta}. \quad (50)$$

A.7 Fourier transformation

Equations (42)–(45) are most conveniently solved in Fourier transform space [14]. Let

$$\bar{\phi}(p) = \int_{-\infty}^{\infty} \tilde{\phi}(X) e^{-\mathrm{i}pX} dX, \quad (51)$$

et cetera. The Fourier transformed linear layer equations become

$$\mathrm{i}(Q_E + Q_e) \bar{\psi} = \frac{d}{dp} (\bar{\phi} - \bar{N}) - p^2 \bar{\psi}, \quad (52)$$

$$\mathrm{i} Q_E \bar{N} = -\mathrm{i} Q_e \bar{\phi} - D^2 \frac{d(p^2 \bar{\psi})}{dp} + c_\beta^2 \frac{d\bar{V}}{dp} - P_E p^2 \bar{N}, \quad (53)$$

$$\mathrm{i}(Q_E + Q_i) p^2 \bar{\phi} = \frac{d(p^2 \bar{\psi})}{dp} - P_\varphi p^4 \left(\bar{\phi} + \frac{\bar{N}}{\iota} \right), \quad (54)$$

$$\mathrm{i} Q_E \bar{V} = \mathrm{i} Q_e \bar{\psi} + \frac{d\bar{N}}{dp} - P_\varphi p^2 \bar{V}, \quad (55)$$

where, for a tearing parity solution,

$$\bar{\phi}(p) - \bar{N}(p) \equiv \bar{Y}(p) \rightarrow \bar{Y}_0 \left[\frac{\hat{\Delta}}{\pi p} + 1 + \mathcal{O}(p) \right] \quad (56)$$

as $p \rightarrow 0$. Here, \bar{Y}_0 is an arbitrary constant.

Finally, if we define

$$\bar{J}(p) = p^2 \bar{\psi}, \quad (57)$$

then Eqs. (52)–(55) can be converted into the following equivalent set of four coupled first-order o.d.e.s:

$$\frac{d\bar{Y}}{dp} = \left[\frac{\mathrm{i}(Q_E + Q_e) + p^2}{p^2} \right] \bar{J}, \quad (58)$$

$$\frac{d\bar{N}}{dp} = - \left(\frac{\mathrm{i} Q_e}{p^2} \right) \bar{J} + (\mathrm{i} Q_E + P_\varphi p^2) \bar{V}, \quad (59)$$

$$\frac{d\bar{J}}{dp} = [\mathrm{i}(Q_E + Q_i) p^2 + P_\varphi p^4] \bar{Y} + [\mathrm{i}(Q_E + Q_i) p^2 + \iota_e^{-1} P_\varphi p^4] \bar{N}, \quad (60)$$

$$\begin{aligned} c_\beta^2 \frac{d\bar{V}}{dp} &= [\mathrm{i} Q_e + \mathrm{i}(Q_E + Q_i) D^2 p^2 + D^2 P_\varphi p^4] \bar{Y} \\ &\quad + \{ \mathrm{i}(Q_E + Q_e) + [P_E + \mathrm{i}(Q_E + Q_i) D^2] p^2 + \iota_e^{-1} D^2 P_\varphi p^4 \} \bar{N}, \end{aligned} \quad (61)$$

Note that $\iota_e = -Q_e/(Q_i - Q_e)$.

A.8 Small argument expansion

Let us search for power-law solutions of Eqs. (58)–(61) at small values of p . Given that we have four coupled first-order o.d.e.s, we expect to find four independent power-law solutions. The first solution is such that

$$\bar{Y}(p) = \mathrm{i}(Q_E + Q_e) a_{-1} p^{-1} - \left[\frac{\mathrm{i}}{2} Q_E (Q_E + Q_e) (Q_E + Q_i) + 1 \right] a_{-1} p + \mathcal{O}(p^3), \quad (62)$$

$$\bar{N}(p) = -\mathrm{i} Q_e a_{-1} p^{-1} + \frac{\mathrm{i}}{2} Q_E Q_e (Q_E + Q_i) a_{-1} p + \mathcal{O}(p^3), \quad (63)$$

$$\bar{J}(p) = -a_{-1} - \frac{1}{2} Q_E (Q_E + Q_i) a_{-1} p^2 + \mathcal{O}(p^4), \quad (64)$$

$$\bar{V}(p) = - \frac{[\mathrm{i} Q_e (1 + P_E) + Q_E (Q_E + Q_i) D^2]}{2 c_\beta^2} a_{-1} p^2 + \mathcal{O}(p^4), \quad (65)$$

where a_{-1} is an arbitrary constant. The second solution is such that

$$\bar{Y}(p) = i(Q_E + Q_e) a_0 - \frac{i}{6} Q_E (Q_E + Q_e) (Q_E + Q_i) a_0 p^2 + \mathcal{O}(p^4), \quad (66)$$

$$\bar{N}(p) = -i Q_e a_0 + \frac{i}{6} Q_E Q_e (Q_E + Q_i) a_0 p^2 + \mathcal{O}(p^4), \quad (67)$$

$$\bar{J}(p) = -\frac{1}{3} Q_E (Q_E + Q_i) a_0 p^3 + \mathcal{O}(p^5), \quad (68)$$

$$\bar{V}(p) = -\frac{1}{3} \frac{[i Q_e P_E + Q_E (Q_E + Q_i) D^2]}{c_\beta^2} a_0 p^3 + \mathcal{O}(p^5), \quad (69)$$

where a_0 is an arbitrary constant. The third solution is such that

$$\bar{Y}(p) = -\frac{1}{6} (Q_E + Q_e) (Q_E + Q_i) a_2 p^2 + \mathcal{O}(p^4), \quad (70)$$

$$\bar{N}(p) = a_2 + \frac{i}{2} (Q_E + Q_i) \left(-\frac{i}{3} Q_e + \frac{g}{c_\beta^2} \right) a_2 p^2 + \mathcal{O}(p^4), \quad (71)$$

$$\bar{J}(p) = \frac{i}{3} (Q_E + Q_i) a_2 p^3 + \mathcal{O}(p^5), \quad (72)$$

$$\bar{V}(p) = i \frac{(Q_E + Q_e)}{c_\beta^2} a_2 p + \mathcal{O}(p^3), \quad (73)$$

where a_2 is an arbitrary constant. The final solution is such that

$$\bar{Y}(p) = -\frac{i}{12} Q_E (Q_E + Q_e) (Q_E + Q_i) a_3 p^3 + \mathcal{O}(p^5), \quad (74)$$

$$\bar{N}(p) = -i Q_E a_3 p + \mathcal{O}(p^3), \quad (75)$$

$$\bar{J}(p) = -\frac{1}{4} Q_E (Q_E + Q_i) a_3 p^4 + \mathcal{O}(p^6), \quad (76)$$

$$\bar{V}(p) = -\frac{Q_E (Q_E + Q_e)}{2 c_\beta^2} a_3 p^2 + \mathcal{O}(p^4), \quad (77)$$

where a_3 is an arbitrary constant. Note, however, that the third solution is not consistent with Eqs. (46)–(49), which mandate that $\bar{N}(p) \rightarrow -[Q_e/(Q_E + Q_e)] \bar{Y}(p) + \mathcal{O}(p)$ and $\bar{V}(p) \rightarrow \mathcal{O}(p^2)$ as $p \rightarrow 0$. Hence, we deduce that $a_2 = 0$.

We conclude that, at small values of p , the most general solution for $\bar{Y}(p)$ and $\bar{N}(p)$ takes the form

$$\bar{Y}(p) = i(Q_E + Q_e) (a_{-1} p^{-1} + a_0) + \mathcal{O}(p), \quad (78)$$

$$\bar{N}(p) = -i Q_e (a_{-1} p^{-1} + a_0) + \mathcal{O}(p), \quad (79)$$

which is consistent with Eqs. (46)–(49).

A.9 Ricatti matrix differential equation

Let

$$\underline{u} = \begin{pmatrix} \bar{Y} \\ \bar{N} \end{pmatrix}, \quad (80)$$

$$\underline{v} = \begin{pmatrix} \bar{J} \\ c_\beta^2 \bar{V} \end{pmatrix}. \quad (81)$$

Equations (58)–(61) can be written in the form

$$\frac{du}{dp} = \underline{A} \underline{v}, \quad (82)$$

$$\frac{dv}{dp} = \underline{B} \underline{u}, \quad (83)$$

where

$$A_{11} = \frac{i(Q_E + Q_e) + p^2}{p^2}, \quad (84)$$

$$A_{12} = 0, \quad (85)$$

$$A_{21} = -\frac{iQ_e}{p^2}, \quad (86)$$

$$A_{22} = \frac{iQ_E + P_\varphi p^2}{c_\beta^2}, \quad (87)$$

$$B_{11} = i(Q_E + Q_i) p^2 + P_\varphi p^4, \quad (88)$$

$$B_{12} = i(Q_E + Q_i) p^2 + \iota_e^{-1} P_\varphi p^4, \quad (89)$$

$$B_{21} = iQ_e + i(Q_E + Q_i) D^2 p^2 + D^2 P_\varphi p^4, \quad (90)$$

$$B_{22} = i(Q_E + Q_e) + [P_E + i(Q_E + Q_i) D^2] p^2 + \iota_e^{-1} D^2 P_\varphi p^4. \quad (91)$$

Thus, we obtain the following matrix differential equation:

$$\frac{d}{dp} \left(\underline{A}^{-1} \frac{du}{dp} \right) = \underline{B} \underline{u}. \quad (92)$$

Let

$$p \frac{du}{dp} = \underline{W} \underline{u}. \quad (93)$$

The previous two equations can be combined to give

$$\left(p \frac{d\underline{W}}{dp} - \underline{W} + \underline{W} \underline{W} + \underline{A} p \frac{d\underline{A}^{-1}}{dp} \underline{W} - p^2 \underline{A} \underline{B} \right) \underline{u} = \underline{0}, \quad (94)$$

which yields the Riccati matrix differential equation,

$$p \frac{d\underline{W}}{dp} = \underline{W} - \underline{W} \underline{W} - \underline{E} \underline{W} + \underline{F}, \quad (95)$$

where

$$\underline{E}(p) = \underline{A} p \frac{d\underline{A}^{-1}}{dp}, \quad (96)$$

$$\underline{F}(p) = p^2 \underline{A} \underline{B}. \quad (97)$$

In fact, it is easily demonstrated that

$$E_{11} = \frac{2i(Q_E + Q_e)}{i(Q_E + Q_e) + p^2}, \quad (98)$$

$$E_{12} = 0, \quad (99)$$

$$E_{21} = -\frac{2iQ_e(iQ_E + 2P_\varphi p^2)}{[i(Q_E + Q_e) + p^2](iQ_E + P_\varphi p^2)}, \quad (100)$$

$$E_{22} = -\frac{2P_\varphi p^2}{iQ_E + P_\varphi p^2}, \quad (101)$$

and

$$F_{11} = p^2 [i(Q_E + Q_e) + p^2] [i(Q_E + Q_i) + P_\varphi p^2], \quad (102)$$

$$F_{12} = p^2 [i(Q_E + Q_e) + p^2] [i(Q_E + Q_i) + \iota_e^{-1} P_\varphi p^2], \quad (103)$$

$$F_{21} = -iQ_e p^2 [i(Q_E + Q_i) + P_\varphi p^2] + c_\beta^{-2} p^2 (iQ_E + P_\varphi p^2) [iQ_e + i(Q_E + Q_i) D^2 p^2 + D^2 P_\varphi p^4], \quad (104)$$

$$F_{22} = -iQ_e p^2 [i(Q_E + Q_i) + \iota_e^{-1} P_\varphi p^2] + c_\beta^{-2} p^2 (iQ_E + P_\varphi p^2) \{i(Q_E + Q_e) + [P_E + i(Q_E + Q_i) D^2] p^2 + \iota_e^{-1} D^2 P_\varphi p^4\}. \quad (105)$$

Finally, if

$$\underline{\underline{W}}(p) = \begin{pmatrix} W_{11}, & W_{12} \\ W_{21}, & W_{22} \end{pmatrix} \quad (106)$$

then Eq. (95) yields

$$p \frac{dW_{11}}{dp} = W_{11} - W_{11} W_{11} - W_{12} W_{21} - E_{11} W_{11} + F_{11}, \quad (107)$$

$$p \frac{dW_{12}}{dp} = W_{12} - W_{11} W_{12} - W_{12} W_{22} - E_{11} W_{12} + F_{12}, \quad (108)$$

$$p \frac{dW_{21}}{dp} = W_{21} - W_{21} W_{11} - W_{22} W_{21} - E_{21} W_{11} - E_{22} W_{21} + F_{21}, \quad (109)$$

$$p \frac{dW_{22}}{dp} = W_{22} - W_{21} W_{12} - W_{22} W_{22} - E_{21} W_{12} - E_{22} W_{22} + F_{22}. \quad (110)$$

Thus, our final system of equations consists of a set of four coupled nonlinear differential equations.

A.10 Small-argument behavior of Riccati matrix differential equation

It follows from Eqs. (98)–(101) that $\underline{\underline{E}}(p) = \underline{\underline{E}}^{(0)} + \mathcal{O}(p^2)$ at small values of p , where

$$E_{11}^{(0)} = 2, \quad (111)$$

$$E_{12}^{(0)} = 0, \quad (112)$$

$$E_{21}^{(0)} = -\frac{2Q_e}{Q_E + Q_e}, \quad (113)$$

$$E_{22}^{(0)} = 0. \quad (114)$$

Likewise, Eqs. (102)–(105) imply that $\underline{\underline{F}}(p) = \mathcal{O}(p^2)$.

Suppose that $\underline{W}(p) = \underline{W}^{(0)} + \underline{W}^{(1)} p$ at small values of p , where the elements of $\underline{W}^{(0)}$ and $\underline{W}^{(1)}$ are independent of p . Equation (95) gives

$$\underline{0} = \underline{W}^{(0)} - \underline{W}^{(0)} \underline{W}^{(0)} - \underline{E}^{(0)} \underline{W}^{(0)}, \quad (115)$$

$$\underline{0} = -\underline{W}^{(1)} \underline{W}^{(0)} - \underline{W}^{(0)} \underline{W}^{(1)} - \underline{E}^{(0)} \underline{W}^{(1)}. \quad (116)$$

Suitable solutions are

$$\underline{W}^{(0)} = \begin{pmatrix} -1, & 0 \\ -E_{21}^{(0)}/2, & 0 \end{pmatrix}, \quad (117)$$

$$W_{12}^{(1)} = 0, \quad (118)$$

$$W_{21}^{(1)} = \frac{E_{21}^{(0)}}{2} [W_{11}^{(1)} - W_{22}^{(1)}]. \quad (119)$$

At small values of p , let

$$\underline{u}(p) = \underline{u}_{-1} p^{-1} + \underline{u}_0, \quad (120)$$

where the elements of \underline{u}_{-1} (which are y_{-1} and n_{-1} , respectively) and the elements of \underline{u}_0 (which are y_0 and n_0 , respectively) are all constants. Equation (93) gives

$$\underline{W}^{(0)} \underline{u}_{-1} = -\underline{u}_{-1}, \quad (121)$$

$$\underline{W}^{(0)} \underline{u}_0 + \underline{W}^{(1)} \underline{u}_{-1} = \underline{0}. \quad (122)$$

Thus, making use of Eq. (117), we get

$$\begin{pmatrix} -1, & 0 \\ -E_{21}^{(0)}/2, & 0 \end{pmatrix} \begin{pmatrix} y_{-1} \\ n_{-1} \end{pmatrix} = - \begin{pmatrix} y_{-1} \\ n_{-1} \end{pmatrix}, \quad (123)$$

which implies that

$$\frac{E_{21}^{(0)}}{2} y_{-1} = -\frac{Q_e}{Q_E + Q_e} y_{-1} = n_{-1}, \quad (124)$$

in accordance with Eqs. (78) and (79), where use has been made of Eq. (113). Equations (117)–(119) and (122) yield

$$\frac{y_0}{y_{-1}} = W_{11}^{(1)}, \quad (125)$$

with n_0 undetermined. It follows from Equation (56) that

$$\frac{\pi}{\hat{\Delta}} \equiv \frac{y_0}{y_{-1}} = W_{11}^{(1)} = \frac{dW_{11}(0)}{dp}. \quad (126)$$

Note, incidentally, that the result $\pi/\hat{\Delta} = y_0/y_{-1}$ follows directly from Eq. (42), and would hold even if a_2 were non-zero.

A.11 Large-argument behavior of Riccati matrix differential equation

At large values of p , it is clear from Eqs. (102)–(105) that $\underline{F}(p) = \underline{F}^{(6)} p^6 + \underline{F}^{(8)} p^8$, where the elements of $\underline{F}^{(6)}$ and $\underline{F}^{(8)}$ are independent of p . On the other hand, Eqs. (98)–(101) imply that $\underline{E}(p) = \underline{E}^{(0)}$, where the

elements of $\underline{E}^{(0)}$ are independent of p . Thus, if we write $\underline{W}(p) = \underline{W}^{(2)} p^2 + \underline{W}^{(4)} p^4$, where the elements of $\underline{W}^{(2)}$ and $\underline{W}^{(4)}$ are independent of p , then Eq. (95) gives

$$\underline{W}^{(4)} \underline{W}^{(4)} = \underline{F}^{(8)}, \quad (127)$$

$$\underline{W}^{(2)} \underline{W}^{(4)} + \underline{W}^{(4)} \underline{W}^{(2)} = \underline{F}^{(6)}. \quad (128)$$

Now, according to Eqs. (102)–(105),

$$F_{11}^{(8)} = 0, \quad (129)$$

$$F_{12}^{(8)} = 0, \quad (130)$$

$$F_{21}^{(8)} = c_\beta^{-2} D^2 P_\varphi^2, \quad (131)$$

$$F_{22}^{(8)} = c_\beta^{-2} \iota_e^{-1} D^2 P_\varphi^2, \quad (132)$$

so Eq. (127) yields

$$W_{11}^{(4)} = 0, \quad (133)$$

$$W_{12}^{(4)} = 0, \quad (134)$$

$$W_{21}^{(4)} = -c_\beta^{-1} \iota_e^{1/2} D P_\varphi, \quad (135)$$

$$W_{22}^{(4)} = -c_\beta^{-1} \iota_e^{-1/2} D P_\varphi, \quad (136)$$

where we have chosen the sign of the square root that is associated with well-behaved solutions at large values of p . Here, we are assuming that $\iota_e > 0$. Equations (102)–(105) also give

$$F_{11}^{(6)} = P_\varphi, \quad (137)$$

$$F_{12}^{(6)} = \iota_e^{-1} P_\varphi, \quad (138)$$

$$F_{21}^{(6)} = i c_\beta^{-2} Q_E D^2 P_\varphi + i c_\beta^{-2} (Q_E + Q_i) D^2 P_\varphi, \quad (139)$$

$$F_{22}^{(6)} = i c_\beta^{-2} \iota_e^{-1} Q_E D^2 P_\varphi + c_\beta^{-2} [P_E + i (Q_E + Q_i) D^2] P_\varphi. \quad (140)$$

Thus, Eq. (128) yields

$$W_{12}^{(2)} W_{21}^{(4)} = F_{11}^{(6)}, \quad (141)$$

$$W_{12}^{(2)} W_{22}^{(4)} = F_{12}^{(6)}, \quad (142)$$

which gives

$$W_{12}^{(2)} = -c_\beta \iota_e^{-1/2} D^{-1}. \quad (143)$$

Now, if

$$\underline{W} \underline{u} = \lambda(p) \underline{u} \quad (144)$$

then Eq. (93) yields

$$p \frac{d\underline{u}}{dp} = \lambda \underline{u}, \quad (145)$$

which implies that

$$\underline{u}(p) = \underline{u}(p_0) \exp \left[\int_{p_0}^p \frac{\lambda_r(p')}{p'} dp' \right] \exp \left[i \int_{p_0}^p \frac{\lambda_i(p')}{p'} dp' \right], \quad (146)$$

where λ_r and λ_i are the real and imaginary parts of λ , respectively. Of course, a solution that is well behaved at large values of p is such that λ_r is negative. As we have seen, the large- p limit of Eq. (95) is

$$\underline{\underline{W}} \underline{\underline{W}} = \underline{\underline{F}}. \quad (147)$$

Hence, if

$$\underline{\underline{F}} u = \Lambda u \quad (148)$$

then Eqs. (144) and (148) imply that

$$\lambda^2 = \Lambda. \quad (149)$$

The eigenvalue problem for the F -matrix reduces to

$$\Lambda^2 - (F_{11} + F_{22}) \Lambda + F_{11} F_{22} - F_{12} F_{21} = 0. \quad (150)$$

Now,

$$F_{11} + F_{22} \simeq F_{22}^{(8)} p^8 = c_\beta^{-2} \iota_e^{-1} D^2 P_\varphi^2 p^8, \quad (151)$$

$$\begin{aligned} F_{11} F_{22} - F_{12} F_{21} &\simeq \left[F_{11}^{(6)} F_{22}^{(8)} - F_{12}^{(6)} F_{21}^{(8)} \right] p^{14} \\ &+ \left[F_{11}^{(6)} F_{22}^{(6)} - F_{12}^{(6)} F_{21}^{(6)} \right] p^{12} = c_\beta^{-2} R P_\varphi^2 p^{12}, \end{aligned} \quad (152)$$

where

$$R = P_E + i(1 - \iota_e^{-1})(Q_E + Q_i) D^2, \quad (153)$$

Hence, the two eigenvalues of the F -matrix are

$$\Lambda_1 \simeq F_{22}^{(8)} p^8 = c_\beta^{-2} \iota_e^{-1} D^2 P_\varphi^2 p^8, \quad (154)$$

$$\Lambda_2 \simeq \frac{[F_{11}^{(6)} F_{22}^{(6)} - F_{12}^{(6)} F_{21}^{(6)}]}{F_{22}^{(8)}} p^4 = \iota_e D^{-2} R p^4. \quad (155)$$

Thus, we deduce that the two eigenvalues of the W -matrix are

$$\lambda_1 = -\Lambda_1^{1/2} = -c_\beta^{-1} \iota_e^{-1/2} D P_\varphi p^4, \quad (156)$$

$$\lambda_2 = -\Lambda_2^{1/2} = -\iota_e^{1/2} D^{-1} R^{1/2} p^2, \quad (157)$$

Here, the square root of R is taken such that the real part of λ_2 is negative. Now, the eigenvalue problem for the W -matrix reduces to

$$\lambda^2 - W_{22}^{(4)} p^4 \lambda + \left[W_{11}^{(2)} W_{22}^{(4)} - W_{12}^{(2)} W_{21}^{(4)} \right] p^6 = 0. \quad (158)$$

which yields

$$\lambda_1 \simeq W_{22}^{(4)} p^4, \quad (159)$$

which is in agreement with Eq. (156), and

$$\lambda_2 \simeq \left[W_{11}^{(2)} - \frac{W_{12}^{(2)} W_{21}^{(4)}}{W_{22}^{(4)}} \right] p^2, \quad (160)$$

which implies that

$$W_{11}^{(2)} = -\iota_e^{1/2} D^{-1} R^{1/2} - c_\beta \iota_e^{1/2} D^{-1}. \quad (161)$$

Hence, the large- p boundary condition for the W -matrix is

$$\underline{\underline{W}}(p) = \begin{pmatrix} -\iota_e^{1/2} D^{-1} R^{1/2} p^2 - c_\beta \iota_e^{1/2} D^{-1} p^2, & -c_\beta \iota_e^{-1/2} D^{-1} p^2 \\ -c_\beta^{-1} \iota_e^{1/2} D P_\varphi p^4, & -c_\beta^{-1} \iota_e^{-1/2} D P_\varphi p^4 \end{pmatrix}. \quad (162)$$

A.12 Method of solution

The method of solution is to launch the well-behaved asymptotic solution (162) of Eqs. (107)–(110) from large p , and then integrate the equations backward to small p . The scaled layer response index, $\hat{\Delta}$, is then determined from Eq. (126). Note that, because there are no free parameters in expression (162), the scaled layer response index is uniquely determined by this procedure. Finally, the true layer response index is given by $\Delta_s = S^{1/3} \hat{\Delta}$. [See Eq. (50).]

References

- [1] J.T. Scoville, R.J. La Haye, A.G. Kellman, T.H. Osborne, R.D. Stambaugh, E.J. Strait and T.S. Taylor, *Locked modes in DIII-D and a method for prevention of the low density mode*, Nucl. Fusion **31**, 875 (1991).
- [2] T.C. Hender, R. Fitzpatrick, A.W. Morris, P.G. Carolan, R.D. Durst, T. Edlington, J. Ferreira, S.J. Fielding, P.S. Haynes, J. Hugill, et al., *Effect of resonant magnetic perturbations on COMPASS-C tokamak discharges*, Nucl. Fusion **32**, 2091 (1992).
- [3] G.M. Fishpool and P.S. Haynes, *Field error instabilities in JET*, Nucl. Fusion **34**, 109 (1994).
- [4] R. Fitzpatrick, *Interaction of tearing modes with external structures in cylindrical geometry (plasma)*, Nucl. Fusion **33**, 1049 (1993).
- [5] R. Fitzpatrick, *Bifurcated states of a rotating tokamak plasma in the presence of a static error-field*, Phys. Plasmas **5**, 3325 (1998).
- [6] M. Pharr, N.C. Logan, C. Paz-Soldan, J.-K. Park and C. Hansen, *Error field predictability and consequences for ITER*, Nucl. Fusion **64**, 126025 (2024).
- [7] T.E. Evans, R.A. Moyer, J.G. Watkins, P.R. Thomas, T.H. Osborne, J.A. Boedo, M.E. Fenstermacher, K.H. Finken, R.J. Groebner, M. Groth, et al., *Suppression of large edge-localized modes in high-confinement DIII-D plasmas with a stochastic magnetic boundary*, Phys. Rev. Lett. **92**, 235003 (2004).
- [8] A.H. Boozer, *Physics of magnetically confined plasmas*, Rev. Mod. Phys. **76**, 1071 (2004).
- [9] Z. Chang and J.D. Callen, *Global energy confinement degradation due to macroscopic phenomena in tokamaks*, Nucl. Fusion **30**, 219 (1990).

- [10] H.P. Furth, J. Killeen and M.N. Rosenbluth, *Finite-resistivity instabilities of a sheet pinch*, Phys. Fluids **6**, 459 (1963).
- [11] G. Ara, B. Basu, B. Coppi, G. Laval, M.N. Rosenbluth and B.V. Waddell, *Magnetic reconnection and $m = 1$ oscillations in current carrying plasmas*, Ann. Phys. (NY) **112**, 443 (1978).
- [12] J.F. Drake and Y.C. Lee, *Kinetic theory of tearing instabilities*, Phys. Fluids **20**, 1341 (1977).
- [13] F.L. Waelbroeck, *Shielding of resonant magnetic perturbations in the long mean-free-path regime*, Phys. Plasmas **10**, 4040 (2003).
- [14] A. Cole and R. Fitzpatrick, *Drift-magnetohydrodynamical model of error-field penetration in tokamak plasmas*, Phys. Plasmas **13**, 032503 (2006).
- [15] R. Fitzpatrick and F.L. Waelbroeck, *Two-fluid magnetic island dynamics in slab geometry. I. Isolated islands*, Phys. Plasmas **12**, 022307 (2005).
- [16] R.D. Hazeltine, M. Kotschenreuther and P.G. Morrison, *A four-field model for tokamak plasma dynamics*, Phys. Fluids **28**, 2466 (1985).
- [17] R. Fitzpatrick, *Influence of anomalous perpendicular transport on linear tearing mode dynamics in tokamak plasmas*, Phys. Plasmas **29**, 032507 (2022).
- [18] Y. Lee, J.-K. Park and Y.-S. Na, *Effect of parallel flow on resonant layer responses in high beta plasmas*, Nucl. Fusion **64**, 106058 (2024).
- [19] D.P. Brennan, A.J. Cole, C. Akcay and J.M. Finn, *Two fluid stability of rotating tokamak plasmas using machine learning*, Bul. Am. Phys. Soc. **64**, 249 (2019).
- [20] J.-K. Park, *Parametric dependencies of resonant layer responses across linear, two-fluid, drift-MHD regimes*, Phys. Plasmas **29**, 072506 (2022).
- [21] R. Fitzpatrick, *Tearing mode dynamics in tokamak plasmas*, (IOP Publishing, Bristol UK, 2023).
- [22] C. Paz-Soldan, R. Nazikian, L. Cui, B.C. Lyons, D.M. Orlov, A. Kirk, N.C. Logan, T.H. Osborne, W. Suttrop and D.B. Weisberg, *The effect of plasma shape and neutral beam mix on the rotation threshold for RMP-ELM suppression*, Nucl. Fusion **59**, 056012 (2019).

- [23] R. Fitzpatrick, *Response of a magnetically diverted tokamak plasma to a resonant magnetic perturbation*, arXiv 2511.07666 (2025).
- [24] H.P. Furth, P.H. Rutherford and H. Selberg, *Tearing mode in the cylindrical tokamak*, Phys. Fluids **16**, 1054 (1973).
- [25] J.A. Wesson, *Hydromagnetic stability of tokamaks*, Nucl. Fusion **18**, 87 (1978).
- [26] R. Fitzpatrick, *Calculation of tearing mode stability in an inverse aspect-ratio expanded tokamak plasma equilibrium*, Phys. Plasmas **31**, 102507 (2024).
- [27] R. Fitzpatrick, *Investigation of tearing mode stability near ideal stability boundaries via asymptotic matching techniques*, Phys. Plasmas **32**, 062509 (2025).
- [28] F. Wagner, G. Becker, K. Behringer, D. Campbell, A. Eberhagen, W. Engelhardt, G. Fussmann, O. Gehre, J. Gernhardt, et al., *Regime of improved confinement and high beta in neutral-beam-heated divertor discharges of the ASDEX tokamak*, Phys. Rev. Lett. **49**, 1408 (1982).
- [29] P.B. Snyder, H.R. Wilson, J.R. Ferron, L.L. Lao, A.W. Leonard, T.H. Osborne, A.D. Turnbull, D. Mossessian, M. Murakami and X.Q. Xu, *Edge localized modes and the pedestal: A model based on coupled peeling-ballooning modes*, Phys. Plasmas **9**, 2037 (2002).
- [30] H. Zohm, *Edge localized modes (ELMs)*, Plasma Phys. Controlled Fusion **38**, 105 (1996).
- [31] A. Loarte, G. Saibene, R. Sartori, M. Bécoulet, L. Horton, T. Eich, A. Herrmann, M. Laux, G. Matthews, S. Jachmich, et al., *ELM energy and particle losses and their extrapolation to burning plasma experiments*, J. Nucl. Mater. **313**, 962 (2003).
- [32] Y. Liang, H.R. Koslowski, P.R. Thomas, E. Nardon, B. Alper, P. Andrew, Y. Andrew, G. Arnoux, Y. Baranov, M. Bécoulet, et al., *Active control of type-I edge-localized modes with perturbation fields in the JET tokamak*, Phys. Rev. Lett. **98**, 265004 (2007).
- [33] W. Suttrop, T. Eich, J.C. Fuchs, S. Günter, A. Janzer, A. Herrmann, A. Kallenbach, P.T. Lang, T. Lunt, M. Maraschek, et al., *First observation of edge localized modes mitigation with resonant and nonresonant magnetic perturbations in ASDEX-Upgrade*, Phys. Rev. Lett. **106**, 225004 (2011).

- [34] Y.M. Jeon, J.-K. Park, S.W. Yoon, W.H. Ko, S.G. Lee, K.D. Lee, G.S. Yun, Y.U. Nam, W.C. Kim, J.-G. Kwak, K.S. Lee, H.K. Kim, H.L. Yang, et al., *Suppression of edge localized modes in high-confinement KSTAR plasmas by nonaxisymmetric magnetic perturbations*, Phys. Rev. Lett. **109**, 035004 (2012).
- [35] A. Kirk, I.T. Chapman, Y. Liu, P. Cahyna, P. Denner, G. Fishpool, C.J. Ham, J.R. Harrison, Y. Liang, E. Nardon, S. Saarelma, R. Scannell, A.J. Thornton, and MAST Team, *Understanding edge-localized mode mitigation by resonant magnetic perturbations on MAST*, Nucl. Fusion **53**, 043007 (2013).
- [36] T. Sun, Y. Liang, Y.Q. Liu, S. Gu, X. Yang, W. Guo, T. Shi, M. Jia, L. Wang, B. Lyu, et al., *Nonlinear transition from mitigation to suppression of the edge localized mode with resonant magnetic perturbations in the EAST tokamak*, Phys. Rev. Lett. **117**, 115001 (2016).
- [37] D.A. Ryan, C. Ham, A. Kirk, T. Markovic, S. Munaretto, L. Piron, S. Saarelma, W. Suttrop, A.J. Thornton, E. Viezzer, M. Willensdorfer, the MAST Upgrade Team and the EUROfusion Tokamak Exploitation Team, *First observation of RMP ELM mitigation on MAST Upgrade*, Plasma Phys. Control. Fusion **66**, 105003 (2024).
- [38] Q.M. Hu, R. Nazikian, B.A. Grierson, N.C. Logan, J.-K. Park, C. Paz-Soldan and Q. Yu, *The density dependence of edge-localized-mode suppression and density pump-out by resonant magnetic perturbations in the DIII-D tokamak*, Phys. Plasmas **26**, 120702 (2019).
- [39] Q.M. Hu, R. Nazikian, B.A. Grierson, N.C. Logan, C. Paz-Solden and Q. Yu, *The role of edge resonant magnetic perturbations in edge-localized-mode suppression and density pump-out in low-collisionality DIII-D plasmas*, Nucl. Fusion **60**, 076001 (2020).
- [40] Q.M. Hu, R. Nazikian, B.A. Grierson, N.C. Logan, D.M. Orlav, C. Paz-Solden and Q. Yu, *Wide operational windows of edge-localized mode suppression by resonant magnetic perturbations in the DIII-D tokamak*, Phys. Rev. Lett. **125**, 045001 (2020).
- [41] R. Fitzpatrick, *Theory of edge localized mode suppression by static resonant perturbations in the DIII-D tokamak*, Phys. Plasmas **27**, 042506 (2020).
- [42] R. Fitzpatrick and A.O. Nelson, *An improved theory of the response of DIII-D H-mode discharges to static resonant magnetic perturbations and its implications for the suppression of edge localized modes*, Phys. Plasmas **27**, 072501 (2020).

- [43] R.J. Bickerton, J.W. Connor and J.B. Taylor, *Diffusion driven plasma currents and bootstrap tokamak*, Nat. Phys. Sci. **229**, 110 (1971).
- [44] R. Fitzpatrick, Phys. Plasmas **31**, *Inverse aspect-ratio expanded tokamak equilibria*, 082505 (2024).
- [45] R.J. Groebner and S. Saarelma, *Elements of H-mode pedestal structure*, Plasma Phys. Control. Fusion **65**, 073001 (2023).
- [46] Q. Yu, S. Günter, and B.D. Scott, *Numerical modeling of linear drift-tearing mode stability*, Phys. Plasmas **10**, 797 (2003).
- [47] Q. Yu and S. Günter, *Plasma response to externally applied resonant perturbations*, Nucl. Fusion **51**, 073030 (2011).
- [48] C. Chrystal, B.A. Grierson, S.R. Haskey, A.C. Sontag, F.M. Poli, M.W. Shafer and J.S. de Grassie, *Predicting the rotation profile in ITER*, Nucl. Fusion **60**, 036003 (2020).
- [49] Z.S. Hartwig and Y.A. Podpaly, *Magnetic fusion energy formulary*, (Massachusetts Institute of Technology, Cambridge MA, 2011).
- [50] F.L. Hinton and C. Oberman, *Electrical conductivity of plasma in a spatially homogeneous magnetic field*, Nucl. Fusion **9**, 319 (1969).
- [51] J.A. Wesson, *Tokamaks*, 4th ed., (Oxford University Press Oxford UK, 2011).
- [52] R.D. Hazeltine and J.D. Meiss, *Plasma confinement*, (Dover, New York NY, 2003).