

Homework 4

Due date: someday, sometime!

Submit on NYU Brightspace.

Exercise 1. [100 pts] Fruits on trees can freeze in Winter time even when the ambient temperature is above 0 °C. This exercise tries to explain this counterintuitive phenomenon.

Consider a spherical-shape orange of radius a in an ambient with temperature T_∞ and heat transfer coefficient h . The orange also radiates, with emissivity ϵ , to the sky with an effective temperature T_s . Assume a spatially uniform temperature within and on the surface of the orange.

- (i) Write an energy balance for the orange taking into account the convection to the ambient and radiation to the sky (neglect radiation to the ambient). Simplify the equation for constant ρ and C_p of the orange, to reach to an ordinary differential equation (ODE) for the temperature of the orange, T , as a function of time t .
- (ii) Use Euler's method to solve the ODE. Assume that the physical properties of orange are close to those of water, i.e., $\rho = 1000 \text{ kg/m}^3$, $C_p = 4184 \text{ J/(kg} \cdot \text{K)}$. Let $T_\infty = 2 \text{ }^\circ\text{C}$, $h = 20 \text{ W/(m}^2 \cdot \text{K)}$, $a = 5 \text{ cm}$, $\epsilon = 1$. Consider three different effective sky temperatures $-20, -10, -5 \text{ }^\circ\text{C}$. (Careful! The radiation equation works with absolute temperature in K not $^\circ\text{C}$.) Make a plot of the orange temperature T in Celsius ($^\circ\text{C}$) versus time in minutes for each sky temperature. Also, plot horizontal black (different from the other three curves) lines at $T = 0 \text{ }^\circ\text{C}$ and $T = T_\infty$ to denote the freezing and ambient temperatures. Use an initial temperature $10 \text{ }^\circ\text{C}$ and simulate each problem for 7τ , where $\tau = mC_p/(hA)$ is the convection time scale. Also, use $\Delta t = 0.01\tau$. For each of the sky temperatures, you should see that the orange temperature reaches to a constant, equilibrium (or steady-state), value (not changing significantly with time). If this constant value is below the freezing temperature, the orange freezes! Does this happen here for any of the sky temperatures?
- (iii) The steady-state value can be obtained directly from the ODE by using the Newton's method. Find the steady-state temperature for the three cases above directly, and compare to the values you obtained by time marching in item (ii). With the latter, take the last calculated temperature as the equilibrium temperature.
- (iv) In a separate figure, plot the sky temperature required for a $0 \text{ }^\circ\text{C}$ (273.15 K) equilibrium temperature as a function of $0 \leq T_\infty \text{ (}^\circ\text{C)} \leq 10$. Next, explain, mathematically, under what conditions (for what T_∞), there is no Real valued T_s . What does it mean physically?

notes: You need to submit your **CODE** (commented) as well. Also, include your **PLOTS** in the report. As a general rule, any results obtained by the code (whether it is a plot or a computed value) should be included in the report. Solutions without interpretation of the results will not receive full credits.