

# Data Mining

## Classification: Alternative Techniques

---

### Bayesian Classifiers

### Bayes Classifier

---

- A probabilistic framework for solving classification problems

- Conditional Probability:  $P(Y | X) = \frac{P(X, Y)}{P(X)}$

$$P(X | Y) = \frac{P(X, Y)}{P(Y)}$$

- Bayes theorem:

$$P(Y | X) = \frac{P(X | Y)P(Y)}{P(X)}$$

## Example of Bayes Theorem

- Given:

- A doctor knows that meningitis causes stiff neck 50% of the time
- Prior probability of any patient having meningitis is 1/50,000
- Prior probability of any patient having stiff neck is 1/20

- If a patient has stiff neck, what's the probability he/she has meningitis?

$$P(M | S) = \frac{P(S | M)P(M)}{P(S)} = \frac{0.5 \times 1/50000}{1/20} = 0.0002$$

## Using Bayes Theorem for Classification

- Consider each attribute and class label as random variables
- Given a record with attributes  $(X_1, X_2, \dots, X_d)$ 
  - Goal is to predict class  $Y$
  - Specifically, we want to find the value of  $Y$  that maximizes  $P(Y | X_1, X_2, \dots, X_d)$
- Can we estimate  $P(Y | X_1, X_2, \dots, X_d)$  directly from data?

## Using Bayes Theorem for Classification

- Approach:
  - compute posterior probability  $P(Y | X_1, X_2, \dots, X_d)$  using the Bayes theorem

$$P(Y | X_1 X_2 \dots X_n) = \frac{P(X_1 X_2 \dots X_d | Y) P(Y)}{P(X_1 X_2 \dots X_d)}$$

- *Maximum a-posteriori*: Choose  $Y$  that maximizes  $P(Y | X_1, X_2, \dots, X_d)$
  - Equivalent to choosing value of  $Y$  that maximizes  $P(X_1, X_2, \dots, X_d | Y) P(Y)$
- How to estimate  $P(X_1, X_2, \dots, X_d | Y)$ ?

## Naïve Bayes Classifier

- Assume independence among attributes  $X_i$  when class is given:
  - $P(X_1, X_2, \dots, X_d | Y_j) = P(X_1 | Y_j) P(X_2 | Y_j) \dots P(X_d | Y_j)$
  - Can estimate  $P(X_i | Y_j)$  for all  $X_i$  and  $Y_j$  from data
  - New point is classified to  $Y_j$  if  $P(Y_j) \prod P(X_i | Y_j)$  is maximal.

## Conditional Independence

- **X** and **Y** are conditionally independent given **Z** if  
 $P(\mathbf{X}|\mathbf{YZ}) = P(\mathbf{X}|\mathbf{Z})$
- Example: Arm length and reading skills
  - Young child has shorter arm length and limited reading skills, compared to adults
  - If age is fixed, no apparent relationship between arm length and reading skills
  - Arm length and reading skills are conditionally independent given age

## Estimate Probabilities from Data

Tid	Refund	Marital Status	Taxable Income	Evade
1	Yes	Single	125K	No
2	No	Married	100K	No
3	No	Single	70K	No
4	Yes	Married	120K	No
5	No	Divorced	95K	Yes
6	No	Married	60K	No
7	Yes	Divorced	220K	No
8	No	Single	85K	Yes
9	No	Married	75K	No
10	No	Single	90K	Yes

- Class:  $P(Y) = N_c / N$

- e.g.,  $P(\text{No}) = 7/10$ ,  
 $P(\text{Yes}) = 3/10$

- For discrete attributes:

$$P(X_i | Y_k) = |X_{ik}| / N_{c_k}$$

- where  $|X_{ik}|$  is number of instances having attribute value  $X_i$  and belonging to class  $Y_k$
- Examples:

$$P(\text{Status}=\text{Married}|\text{No}) = 4/7$$
$$P(\text{Refund}=\text{Yes}|\text{Yes})=0$$

## Estimate Probabilities from Data

- For continuous attributes:
  - **Discretization:** Partition the range into bins:
    - ◆ Replace continuous value with bin value
      - Attribute changed from continuous to ordinal<sup>k</sup>
  - **Probability density estimation:**
    - ◆ Assume attribute follows a normal distribution
    - ◆ Use data to estimate parameters of distribution (e.g., mean and standard deviation)
    - ◆ Once probability distribution is known, use it to estimate the conditional probability  $P(X_i|Y)$

## Estimate Probabilities from Data

Tid	Refund	Marital Status	Taxable Income	Evade
1	Yes	Single	125K	No
2	No	Married	100K	No
3	No	Single	70K	No
4	Yes	Married	120K	No
5	No	Divorced	95K	Yes
6	No	Married	60K	No
7	Yes	Divorced	220K	No
8	No	Single	85K	Yes
9	No	Married	75K	No
10	No	Single	90K	Yes

- Normal distribution:

$$P(X_i | Y_j) = \frac{1}{\sqrt{2\pi\sigma_{ij}^2}} e^{-\frac{(X_i - \mu_{ij})^2}{2\sigma_{ij}^2}}$$

- One for each  $(X_i, Y_i)$  pair

- For (Income, Class=No):

- If Class=No

- ◆ sample mean = 110
- ◆ sample variance = 2975

$$P(\text{Income} = 120 | \text{No}) = \frac{1}{\sqrt{2\pi(54.54)}} e^{-\frac{(120-110)^2}{2(2975)}} = 0.0072$$

## Example of Naïve Bayes Classifier

**Given a Test Record:**

$X = (\text{Refund} = \text{No}, \text{Married}, \text{Income} = 120\text{K})$

naive Bayes Classifier:

$P(\text{Refund}=\text{Yes}|\text{No}) = 3/7$   
 $P(\text{Refund}=\text{No}|\text{No}) = 4/7$   
 $P(\text{Refund}=\text{Yes}|\text{Yes}) = 0$   
 $P(\text{Refund}=\text{No}|\text{Yes}) = 1$   
 $P(\text{Marital Status}=\text{Single}|\text{No}) = 2/7$   
 $P(\text{Marital Status}=\text{Divorced}|\text{No}) = 1/7$   
 $P(\text{Marital Status}=\text{Married}|\text{No}) = 4/7$   
 $P(\text{Marital Status}=\text{Single}|\text{Yes}) = 2/7$   
 $P(\text{Marital Status}=\text{Divorced}|\text{Yes}) = 1/7$   
 $P(\text{Marital Status}=\text{Married}|\text{Yes}) = 0$

For taxable income:

If class=No: sample mean=110  
                   sample variance=2975  
 If class=Yes: sample mean=90  
                   sample variance=25

- $P(X|\text{Class}=\text{No}) = P(\text{Refund}=\text{No}|\text{Class}=\text{No})$   
 $\times P(\text{Married}|\text{Class}=\text{No})$   
 $\times P(\text{Income}=120\text{K}|\text{Class}=\text{No})$   
 $= 4/7 \times 4/7 \times 0.0072 = 0.0024$
- $P(X|\text{Class}=\text{Yes}) = P(\text{Refund}=\text{No}|\text{Class}=\text{Yes})$   
 $\times P(\text{Married}|\text{Class}=\text{Yes})$   
 $\times P(\text{Income}=120\text{K}|\text{Class}=\text{Yes})$   
 $= 1 \times 0 \times 1.2 \times 10^{-9} = 0$

Since  $P(X|\text{No})P(\text{No}) > P(X|\text{Yes})P(\text{Yes})$

Therefore  $P(\text{No}|X) > P(\text{Yes}|X)$   
 $\Rightarrow \text{Class} = \text{No}$

## Naïve Bayes Classifier

- If one of the conditional probabilities is zero, then the entire expression becomes zero
- Probability estimation:

$$\text{Original : } P(A_i | C) = \frac{N_{ic}}{N_c}$$

$$\text{Laplace : } P(A_i | C) = \frac{N_{ic} + 1}{N_c + c}$$

$$\text{m - estimate : } P(A_i | C) = \frac{N_{ic} + mp}{N_c + m}$$

c: number of classes

p: prior probability

m: parameter

## Example of Naïve Bayes Classifier

Name	Give Birth	Can Fly	Live in Water	Have Legs	Class
human	yes	no	no	yes	mammals
python	no	no	no	no	non-mammals
salmon	no	no	yes	no	non-mammals
whale	yes	no	yes	no	mammals
frog	no	no	sometimes	yes	non-mammals
komodo	no	no	no	yes	non-mammals
bat	yes	yes	no	yes	mammals
pigeon	no	yes	no	yes	non-mammals
cat	yes	no	no	yes	mammals
leopard shark	yes	no	yes	no	non-mammals
turtle	no	no	sometimes	yes	non-mammals
penguin	no	no	sometimes	yes	non-mammals
porcupine	yes	no	no	yes	mammals
eel	no	no	yes	no	non-mammals
salamander	no	no	sometimes	yes	non-mammals
gila monster	no	no	no	yes	non-mammals
platypus	no	no	no	yes	mammals
owl	no	yes	no	yes	non-mammals
dolphin	yes	no	yes	no	mammals
eagle	no	yes	no	yes	non-mammals

**A: attributes**

**M: mammals**

**N: non-mammals**

$$P(A|M) = \frac{6}{7} \times \frac{6}{7} \times \frac{2}{7} \times \frac{2}{7} = 0.06$$

$$P(A|N) = \frac{1}{13} \times \frac{10}{13} \times \frac{3}{13} \times \frac{4}{13} = 0.0042$$

$$P(A|M)P(M) = 0.06 \times \frac{7}{20} = 0.021$$

$$P(A|N)P(N) = 0.004 \times \frac{13}{20} = 0.0027$$

Give Birth	Can Fly	Live in Water	Have Legs	Class
yes	no	yes	no	?

**P(A|M)P(M) > P(A|N)P(N)**


**=> Mammals**

## Naïve Bayes (Summary)


- Robust to isolated noise points
- Handle missing values by ignoring the instance during probability estimate calculations
- Robust to irrelevant attributes
- Independence assumption may not hold for some attributes
  - Use other techniques such as Bayesian Belief Networks (BBN)

## Bayesian Belief Networks

- Provides graphical representation of probabilistic relationships among a set of random variables
- Consists of:
  - A directed acyclic graph (dag)
    - ◆ Node corresponds to a variable
    - ◆ Arc corresponds to dependence relationship between a pair of variables
  - A probability table associating each node to its immediate parent



## Conditional Independence




**D is parent of C**  
**A is child of C**  
**B is descendant of D**  
**D is ancestor of A**

- A node in a Bayesian network is conditionally independent of all of its nondescendants, if its parents are known



## Conditional Independence

- Naïve Bayes assumption:



## Probability Tables


- If  $X$  does not have any parents, table contains prior probability  $P(X)$
- If  $X$  has only one parent ( $Y$ ), table contains conditional probability  $P(X|Y)$
- If  $X$  has multiple parents ( $Y_1, Y_2, \dots, Y_k$ ), table contains conditional probability  $P(X|Y_1, Y_2, \dots, Y_k)$



## Example of Bayesian Belief Network

Exercise=Yes	0.7
Exercise=No	0.3

Diet=Healthy	0.25
Diet=Unhealthy	0.75



## Example of Inferencing using BBN

- Given:  $X = (E=No, D=Yes, CP=Yes, BP=High)$ 
  - Compute  $P(HD|E,D,CP,BP)?$

- $P(HD=Yes | E=No, D=Yes) = 0.55$   
 $P(CP=Yes | HD=Yes) = 0.8$   
 $P(BP=High | HD=Yes) = 0.85$ 
  - $P(HD=Yes | E=No, D=Yes, CP=Yes, BP=High)$   
 $\propto 0.55 \times 0.8 \times 0.85 = 0.374$
- $P(HD=No | E=No, D=Yes) = 0.45$   
 $P(CP=Yes | HD=No) = 0.01$   
 $P(BP=High | HD=No) = 0.2$ 
  - $P(HD=No | E=No, D=Yes, CP=Yes, BP=High)$   
 $\propto 0.45 \times 0.01 \times 0.2 = 0.0009$

**Classify X  
as Yes**