
A Computationally Feasible Framework for Causal Probabilistic Explanation

Rafal Urbaniak^{*}

Basis Research Institute

Eli Bingham[†]

Basis Research Institute

Sam Witty

Poorva Garg[‡]

University of California, Los Angeles

Jack Feser[§]

Basis Research Institute

Daniel Waxman[¶]

Basis Research Institute

Abstract

Explaining why a specific outcome occurred, and which inputs deserve the blame or credit, is central to philosophical, scientific, and policy analysis. Existing tools split into two camps. The Halpern–Pearl theory of *actual causality* gives principled verdicts, but only for toy-sized models, because computing them requires enumerating counterfactual scenarios. Scalable attribution methods like SHAP and LIME ignore the causal structure that generated the data, and can give answers that conflict with a careful causal analysis. We close this gap with Probabilistic Causal Impact (PCI).

PCI builds on actual causality and on Pearl’s notions of probability of necessity and sufficiency, but recasts the question “which variables explain this outcome?” as an estimation problem on a probabilistic causal model that one can solve by approximate methods, such as MC. To compute a PCI score, the analyst specifies three ingredients: a distribution Γ over which subsets of variables to consider as candidate explanations, a distribution Δ over the counterfactual values to which those variables might be changed, and a function ci that measures how much such a change shifts the outcome. The result is a tractable, causally grounded explanation procedure that applies to large models. A correspondence theorem shows that PCI agrees with actual causality on the small models where AC is well-defined; against Pearl’s probability of actual causation, PCI recovers Pearl’s within-scenario ranking and additionally captures path-reliability and cross-scenario reliability, intuitions Pearl’s binary indicator cannot capture.

We evaluate PCI in four increasingly demanding settings: a synthetic structural model with redundant and preempting causes, where PCI recovers the AC literature’s verdicts and generalises them to continuous variables; a scaling benchmark where PCI remains tractable as AC enumeration becomes infeasible; a dynamical (Bayesian, ODE) epidemiological model with continuous outcomes and asymmetric causes, where AC offers no verdict; and a deployed spatio-temporal model with deep ML components trained on millions of points, where PCI and SHAP attributions diverge in the same qualitative ways as in the synthetic setting.

Keywords: Probabilistic Causal Models, Explanation, Responsibility, Actual Causality, Probability of Necessity and Sufficiency.

^{*}rafal@basis.ai

[†]eli@basis.ai

[‡]poorvagarg@cs.ucla.edu

[§]jack@basis.ai

[¶]dan@basis.ai

Contents

1	Introduction	4
2	Causal Impact: Motivations	5
2.1	Actual Causality	6
2.2	Probability of Necessity and Sufficiency	7
2.3	Path Forward	10
3	Causal Impact: Definitions	11
3.1	Continuous Outcomes and the General Causal Impact Function	20
3.2	Back to the Running Example: Verifying the Desiderata	21
4	Examples and Comparison to SHAP and Causal SHAP	22
4.1	SHAP and Causal SHAP	23
4.2	OBCB: SHAP and Causal SHAP	24
4.3	Signal with Mediation	28
4.4	Computations and Desiderata Analysis	28
4.5	Summary Across Both Examples	31
5	Synthetic Evaluation with Overdetermination and Undercutting	31
6	Comparison to Differential Causal Effect	35
7	Relation with Actual Causality	37
8	Relation with the Probability of Actual Causality	43
8.1	Pearl's Worked Example: The Desert Traveller	43
8.2	PCI on the Basic Desert Traveller	44
8.3	PCI Expectation vs the AC Machinery	45
8.4	Weak-Poison Variant: Two Intuitions Pearl Misses	45
8.5	Further Differences Between Pearl's Probability of Causation and PCI	47
9	Scaling Actual-Cause Computation: An Empirical Comparison	48
10	Dynamical SIR Benchmark: PCI on Continuous Outcomes	52
11	Scaling PCI to a Real-World Automated Valuation Model	55
12	Discussion and Conclusions	56
A	Computations for the OBCB Running Example	61
A.1	Alice's probability of necessity	61
A.2	Alice's probability of sufficiency	62
A.3	SHAP value function for Option A	62

B	Computations for the Signal-with-Mediation Example	62
B.1	Distributional facts	63
B.2	Conditional expectations	63
B.3	Plain SHAP at Instance 1	63
B.4	Causal SHAP at Instance 1	63
B.5	PCI at Instance 1	64
C	Setup for the Desert-Traveller PCI Computation	64
C.1	Structural model (basic variant)	64
C.2	PCI components	65

1 Introduction

Causal attribution is a central reasoning task in many important problems, such as gauging the efficacy of medical treatments [Shepherd et al., 2024], pinpointing drivers of economic growth [Cheshire and Magrini, 2009], root cause analysis in automated systems [Papageorgiou et al., 2022], or assessing the impact of public policy [Heckman and Pinto, 2022]. We may want to understand why an adverse outcome occurred so as to prevent recurrence, or how a predicted positive outcome could be brought about.

In the absence of scalable, formal causal attribution tools, analysts often resort to ad-hoc methods: strong correlation, historical comparison, domain expertise, or non-causal attribution like Granger causality [Granger, 1969] and SHAP [Lundberg and Lee, 2017]. Even researchers with an advanced command of statistical and experimental methods may not possess a thorough understanding of how to ascribe cause [Shepherd et al., 2024], and would benefit from scalable automation. Here we present a tool for accurate and tractable causal attribution applicable to a wide array of causal, probabilistic, machine-learned models.

What would a satisfactory causal attribution method look like? At minimum, it should be (i) *causally grounded*, tracking counterfactual dependence rather than correlation; (ii) *context-sensitive*, distinguishing causes whose path was active in the factual world from those whose path was blocked; (iii) *broadly applicable*, handling stochastic models, discrete and continuous variables, and finite (not just infinitesimal) counterfactual changes; and (iv) *tractable*, estimable from samples without combinatorial enumeration over witness sets and alternatives. Existing methods each forfeit one or more of these. Actual causality [Halpern, 2016] is grounded and context-sensitive, but neither broadly applicable nor tractable in general, with no clear approximation [Eiter and Lukasiewicz, 2002]. Methods that *do* scale either give up causal grounding or are restricted in scope. SHAP [Lundberg and Lee, 2017] grounds attribution in observational correlations, so a feature can receive high attribution merely for being correlated with a true cause; Causal SHAP [Heskes et al., 2020] addresses this to some extent, but failure modes persist (Section 4). Butler et al. [2022]’s Differential Causal Effect (DCE) is causally grounded but local: it relies on gradients, so requires differentiability and does not capture effects from non-infinitesimal changes in potential causes.⁶ Standard treatment-effect quantities, ATE and CATE, are causally grounded and tractable but lack context-sensitivity: every intervention propagates downstream through the structural equations, overriding the mediator values that mark which causal paths were active in the factual world (Section 2).

We introduce *probabilistic causal impacts* (PCI), an attribution framework that combines the context-sensitivity of actual causality with the scalability of probability-of-causation methods. PCI generalises Pearl’s probability of necessary and sufficient causation [Pearl, 2022] from binary to arbitrary variable domains, and incorporates the *witness* mechanism from actual causality [Halpern, 2016] via integration over a distribution of witness sets rather than existential quantification.

PCI also shares an intuition with the literature on *descriptive normality* [Halpern and Hitchcock, 2015, Icard et al., 2017]: attributions are computed as expectations against a distribution over counterfactual worlds conditioned on upstream factual context, so factual upstream values play the role of a default against which alternatives are graded. We do not aim to resolve the specific counterexamples that motivate the normality literature; the connection is at the level of why a non-uniform alternative-value distribution matters, not in handling explicit normality orderings. Several components have precedents — expectations over exogenous noise [Halpern, 2016], weighting by alternative-value probabilities [Beckers and Vennekens, 2015], aggregation over witness sets [Beckers and Vennekens, 2015, Halpern, 2016] — but no prior construction combines them into a single tractable estimator handling arbitrary variable domains.

PCI presumes access to a trained probabilistic causal model from which counterfactual outcomes can be sampled, a stronger requirement than methods that work from observational data and a causal graph alone (e.g., PN/PS/PNS bounds, Causal SHAP). Within that requirement, however, we place no strong restrictions on the form of the model, distributions, or types of variables. Such models can comprise standard machine learning components such as neural network parametrized distributions or Gaussian processes.

⁶DCE is, fairly, not designed for responsibility attribution: Butler et al. target rates of change of treatment.

With this breadth of application, and a measure that surfaces its design choices — the variable selection distribution, alternative value distribution, and causal impact function — as explicit user-facing components rather than hidden defaults, PCI hopes to provide causal attribution and explanation tooling for a wide array of users and applications.

Paper structure. The paper proceeds in four phases. The first establishes motivation and formalism. Section 2 works through a running example to surface what actual causality, the probability of necessity/sufficiency, and average/conditional treatment effects each get right and miss, motivating the design choices behind PCI. Section 3 then defines the method: a kernel built from a sufficiency check at the factual configuration and a necessity check against alternatives, integrated against a distribution over suspect/witness sets, with three components exposed as user choices: the variable selection distribution Γ , the alternative-value distribution Δ , and the causal impact function.

The second phase positions PCI against existing attribution methods. Section 4 compares PCI with SHAP, Causal SHAP, ATE, and CATE on two small examples (OBCB, with two binary features, and a signal-with-mediation model), tabulating which desiderata each satisfies or fails, including cases where SHAP and Causal SHAP attributions track correlation rather than counterfactual dependence. Section 5 reports a synthetic evaluation built around overdetermination and undercutting, the patterns on which pre-AC accounts most often diverge from intuition, and shows that PCI recovers the literature’s verdicts. Section 6 contrasts PCI with gradient-based attribution, instantiated by Differential Causal Effect [Butler et al., 2022], on a continuous credit-limit example, showing where the two diverge and why PCI better tracks responsibility under unit rescaling and outside the steep region of the response surface.

The third phase establishes the formal and empirical relationship between PCI and actual causality. Section 7 shows that under specific choices of Γ and Δ the PCI kernel is positive on a configuration exactly when that configuration is an actual cause in Halpern’s sense, so AC verdicts can be read off the kernel without enumerating witnesses. Section 8 positions PCI against Pearl’s probability of actual causation: on the canonical desert-traveller example, PCI recovers Pearl’s within-scenario ranking with graded magnitudes, and a weak-poison variant surfaces two additional responsibility intuitions (path-reliability and cross-scenario reliability) that Pearl’s binary indicator structurally cannot express but PCI’s expectation does. Section 9 then benchmarks PCI against explicit actual-cause computation: exact subset enumeration times out at 17 variables, while the necessity-only PCI estimator continues to roughly 73–145 variables (depending on sample budget) and sustains above a 0.85 correct-attribution rate up to about 60 variables, visiting several orders of magnitude fewer cause–witness configurations.

The fourth phase pushes PCI into settings where actual causality no longer applies. Section 10 extends the benchmarking to a continuous-outcome dynamical-SIR (Susceptible–Infected–Recovered) setting, exercising the joint necessity-sufficiency machinery on threshold events (peak overshoot) where AC has no comparable verdict. Where but-for analysis cannot separate the two policies, lockdown and masking, PCI correctly assigns lockdown roughly twice the responsibility of masking, with the gap driven entirely by the necessity term under witness pinning. Section 11 applies PCI to a deployed automated valuation model with machine-learned components trained on millions of points, a regime where actual causality is intractable; SHAP returns qualitatively different attributions, assigning virtually all responsibility to downstream variables and effectively ignoring the rest. Section 12 closes with discussion.

2 Causal Impact: Motivations

The central difficulty in causal attribution is not computing counterfactuals — it is knowing which counterfactuals to compute, and recognising when a cause is blocked by context rather than absent altogether. The closest predecessors of this approach, taken individually, fail to capture this: actual causality handles context sensitivity but is computationally intractable and not general enough; the probability of necessity and sufficiency scales well, but lacks context sensitivity. This section builds intuition for both limitations through a running example, setting up the design choices that motivate PCI in Section 3. We proceed in three steps: revisit Halpern’s actual-cause construction on a deterministic OBCB example to see what witnesses do and why a plain but-for clause misses Alice’s case; turn to Pearl’s PN/PS/PNS on a stochastic OBCB to introduce the probabilistic machinery actual causality lacks while showing that, without witnesses, PN/PS/PNS cannot register Alice’s

missing credit check; and close with a “Path forward” subsection that distills what PCI must inherit from each predecessor before Section 3 formalises it.

2.1 Actual Causality

Consider the following (deterministic, discrete) model:

Example 1 (Bob at Old Boys’ Club Bank). *Bob applied for a home mortgage loan with the local branch of Old Boys’ Club Bank (OBCB), and his application was denied. The OBCB policy requires applicants to pass a credit score check, but it only performs this check if the applicant is male; all non-male applicants are rejected outright. Thus, the procedure is:*

$$\begin{aligned} \text{check} &= (\text{gender} = \text{male}) \\ \text{check-failed} &= \text{check} \wedge (\text{credit} = \text{bad}) \\ \text{loan} &= \neg \text{check-failed} \wedge (\text{gender} = \text{male}) \end{aligned}$$

Why was Bob’s loan application denied? An intuitive explanation is that his credit was bad: had it been good, the outcome would have been different. In contrast, his gender does not appear to be responsible for the rejection: had it been different, his application still would have been denied (albeit this time due to $\text{check} = F$).

This suggests a counterfactual approach to feature responsibility attribution: perhaps we can identify the features responsible for a particular outcome by asking counterfactual questions: would the outcome have been different if a specific feature had a different value? This is called the *but-for clause*: Bob wouldn’t have been denied the loan but for his bad credit.

Concretely, consider a model M with *exogenous variables* $\mathbf{U}$, which are the source of stochasticity, and *endogenous variables* $\mathbf{V}$, which are influenced by other variables in the model. Let $\mathbf{S} \subseteq \mathbf{V}$ be the variables we suspect to be causally responsible (call them *causal suspects*), and $Y \in \mathbf{V}$ be an outcome of interest. Suppose we consider only one suspect $S \in \mathbf{S}$ and mark factual values with $*$. Under a setting $\mathbf{U} = \mathbf{u}$ of the exogenous variables, M produces $S = s^*$ and $Y = y^*$, denoted $(M, \mathbf{u}) \models S = s^*, Y = y^*$. We call this the *factual scenario*. An intervened model where a variable S is set to an alternative value s' is denoted $(M, \mathbf{u})[S \leftarrow s']$.

A first attempt at defining a counterfactual notion of causal responsibility is:

Definition 2 (But-for, single). *$S = s^*$ is causally responsible for $Y = y^*$ in $(M, \mathbf{u})$ if and only if $(M, \mathbf{u}) \models S = s^*, Y = y^*$ and $(M, \mathbf{u})[S \leftarrow s'] \not\models Y = y^*$ for some $s' \neq s^*$.*

In a given context, S is causally responsible for an outcome only when intervening on S with some alternative value would change that outcome. Bob’s credit score satisfies this condition: had his credit score been higher, the loan might have been approved.

However, this cause test is too coarse. Consider another case:

Example 3 (Alice at OBCB). *Alice is a woman with bad credit. She applies for a loan with OBCB and is rejected.*

Why was Alice denied the loan? Under the but-for clause for single cause candidates, neither her gender nor her credit score are responsible. No matter how high her credit score, her application would still have been denied because the bank does not loan to women. Had her gender been different, she would still have been denied for having bad credit.

To address this example, we could consider sets of factors as causes:

Attempt 4 (But-for, sets). *Variables $\mathbf{C} \subseteq \mathbf{S}$ with factual values $\mathbf{C} = \mathbf{c}^*$ are causally responsible for $Y = y^*$ in $(M, \mathbf{u})$ if and only if $(M, \mathbf{u}) \models \mathbf{C} = \mathbf{c}^*, Y = y^*$ and $(M, \mathbf{u})[\mathbf{C} \leftarrow \mathbf{c}'] \not\models Y = y^*$ for some $\mathbf{c}'$ everywhere different from $\mathbf{c}^*$, where no proper subset of $\mathbf{C}$ has this property.*

The minimality condition reflects our preference for explanations that include only relevant properties.

Under this generalization, the set $\mathbf{c}^* = \{\text{gender} = \text{female}, \text{credit} = \text{bad}\}$ is jointly responsible for Alice’s application being denied because there exists $\mathbf{c}' = \{\text{gender} = \text{male}, \text{credit} = \text{good}\}$ that

would change the decision and c^* is minimal. But what can we say about the individual features? We could say that the whole set bears responsibility but individual features do not, or that any feature in a cause set is partially and equally accountable. Both approaches have issues, as they imply the features are equally responsible. This conflicts with our intuition: OBCB did not evaluate Alice’s credit score, so her denial was caused by her gender.

The structural pattern is what the actual-causality literature calls *preemption* (sometimes *undercutting*): two distinct causal routes might each have been sufficient to produce the outcome, but only one is actually engaged, with the other prevented from operating. Alice’s gender preempted her credit by preventing the check from happening, so credit had no chance to act. The closely related notion of *over-determination* covers cases in which multiple sufficient causes coexist without one preempting the other (e.g., two simultaneous fatal shots). Both phenomena defeat naive but-for tests, since intervening on any one of the candidate causes leaves the outcome unchanged.⁷

Prior work suggests a way out of this difficulty: hold some of the context in which counterfactuals are evaluated fixed, calling such fixed variables *witnesses*. Let us focus on the mediator *check-failed* between *gender* and *loan*. If we fix this variable at the factual value by intervening, then counterfactually, Alice would have been granted the loan if she were male. However, there is no context fixed at the actual value such that, had her credit score been higher, she would have been granted the loan. This motivates the following formalization:

Definition 5 (Actual cause). *Let $S \subseteq V$ be a suspect pool and $W \subseteq V$ a witness pool, observed in (M, u) at factual values $S = s^*$ and $W = w^*$. A cause set $C \subseteq S$ at its factual setting $C = c^*$ is an actual cause of $Y = y^*$ in (M, u) if there is a witness set $T \subseteq W$ disjoint from C such that:*

- **Factivity.** *The cause, the witness pool, and the outcome are all observed at their factual values:*

$$(M, u) \models C = c^*, W = w^*, Y = y^*.$$

- **Witnessed necessity.** *Holding T at its factual setting t^* while intervening on the cause overturns the outcome, for some alternative c' distinct everywhere from c^* :*

$$(M, u)[C \leftarrow c', T \leftarrow t^*] \not\models Y = y^*.$$

- **Minimality.** *No proper subset of C satisfies the two conditions above.*

This definition extends but-for causality with a witness set T that is fixed to its factual value by an intervention. Under this definition, to determine the cause of an outcome, we need to find a cause set C , alternative values c' (trivial if the variables are binary and less so if they are categorical or continuous), and a witness set T . C and T are drawn from sets of size at most $2^{|V|}$, so a brute-force computation is expensive.

Putting computational difficulties aside, the deeper limitation of the construction so far is that it lives entirely in deterministic models. Real-world attribution problems are typically probabilistic: policies apply with some probability, mechanisms occasionally fail or are overridden, and the analyst usually wants to reason about the unit-level outcome in the presence of this noise. Lifting actual causality into that setting is not straightforward, and is one of the reasons we turn next to a different family of attribution tools.

2.2 Probability of Necessity and Sufficiency

To bring probability into the picture, we now consider a stochastic variant of OBCB in which the bank’s gendered policies apply only with some probability: women are checked only sometimes, men are skipped only sometimes, and even when the check happens its outcome is occasionally overridden by loan-officer discretion. This is the natural setting for Pearl’s probabilities of necessity, sufficiency, and their conjunction (PN, PS, PNS), which we introduce in this subsection. As we will see, these scale gracefully into the stochastic regime but, without an analogue of the witness mechanism, mis-attribute Alice’s denial.

⁷The canonical preemption example: Jim is both stabbed and poisoned, and dies. Stabbing works quickly, so it kills Jim before the poison takes effect, and the stabbing caused his death. But-for clauses fail to break the symmetry here as well: removing the stabbing leaves Jim dead by poison, and removing the poison leaves him dead by stab.

Example 6 (OBCB, stochastic). *The bank performs credit checks for a randomly selected 20% of women and skips checks for a randomly selected 10% of men. Unchecked applicants are rejected. Furthermore, if a man’s credit check fails, he is nevertheless accepted 5% of the time and if a woman’s credit check succeeds, she is denied 10% of the time. We assume that the population is evenly divided into male and female and good and bad credit. Formally, the model, using 1 for male and 1 for good credit, is:*

$$\begin{aligned}
 \text{loan-prob}(\text{gender}, \text{check-failed}) &= \begin{array}{c|cc} & \text{check-failed}=0 & \text{check-failed}=1 \\ \hline \text{gender}=0 & 0.9 & 0 \\ \text{gender}=1 & 1 & 0.05 \end{array} \\
 \text{credit} &\sim \text{Bern}(0.5) \\
 \text{gender} &\sim \text{Bern}(0.5) \\
 \text{check} \mid \text{gender} &\sim \text{Bern}((1 - \text{gender}) \cdot 0.2 + \text{gender} \cdot 0.9) \\
 \text{check-failed} &= \text{check} \cdot (1 - \text{credit}) \\
 \text{loan-if-checked} \mid \text{gender}, \text{check-failed} &\sim \text{Bern}(\text{loan-prob}(\text{gender}, \text{check-failed})) \\
 \text{loan} &= \text{loan-if-checked} \cdot \text{check}
 \end{aligned}$$

The witness mechanism works well for deterministic models, but actual causality in its standard form offers no straightforward way to handle probabilistic ones. Before extending the witness idea to the stochastic setting, it is instructive to examine a probabilistic approach that scales more naturally — Pearl’s probabilities of necessity and sufficiency [Pearl, 2022] — and to ask what it would miss without witnesses. This will let us identify what probabilistic machinery is needed, and what context-sensitivity is lost when witnesses are absent, motivating the synthesis that PCI provides. The following definitions allow for probabilistic models, but assume that all variables are Boolean (PCI will relax this assumption, allowing for continuous variables). Pearl considers two ways that a cause may relate to an effect: it may be necessary or sufficient for the effect. A cause is necessary if, counterfactually absent the cause, the effect is unlikely to occur; it is sufficient if the effect is counterfactually likely in the presence of the cause. We now formalize these ideas and discuss them in the context of the running example.

Definition 7 (Probability of Necessity). *Let the probability of necessity be the probability that the outcome $Y = y$ would not have occurred in the absence of $C = c^*$, given that both occurred:*

$$PN(C = c^*, Y = y^*) = P(Y_{c'} \neq y^* \mid C = c^*, Y = y^*),$$

where $Y_{c'}$ is the value of Y after intervening on C with $c' \neq c^*$.⁸

Strictly speaking, the probability of necessity is less specific to a particular situation than the actual cause test discussed above. For example, we can consider the probability that being male causes someone to fail to get a loan, but we can’t consider the probability that being male causes Bob specifically (who we know has bad credit) to fail to get a loan. We might try to patch this up by conditioning, for the time being (while also reporting population-level values that follow the definitions more closely).

We illustrate the computation for Alice’s gender. Alice is factually $\text{gender} = \text{female}$, $\text{credit} = \text{bad}$, $\text{loan} = F$; the counterfactual intervenes to set $\text{gender} = \text{male}$ while leaving $\text{credit} = \text{bad}$. Tracing the male-intervention branches through the model (Appendix A, §A.1) gives $PN(\text{gender} = \text{female}, \text{loan} = F \mid \text{Alice}) = 0.9 \cdot 0.05 = 0.045$. Computing the analogous quantities for the remaining (variable, person) pairs gives the individual-level values shown in Table 1; the table also lists population-level values for comparison.

The PN scores partially align with intuition: Bob’s bad credit is high ($PN = 0.90$), and his gender score is zero, matching the reading that his gender played no role. But Alice’s PN values do not separate gender (0.045) from credit (0.18) particularly well, and Alice’s gender PN is only marginally

⁸Read in twin-network fashion: one draws an exogenous noise $\mathbf{u} \sim P_{\mathbf{U}}$, computes the factual Y under the unintervened structural equations and the counterfactual $Y_{c'}$ under the intervened equations, and conditions on the joint event $\{C = c^*, Y = y^*\}$. Both Y and $Y_{c'}$ live on the same probability space and share the same $\mathbf{u}$.

above Bob’s, not the sharp distinction we would want, given that the bank never even evaluated Alice’s credit. PN cannot register this asymmetry: it asks counterfactually what would happen under intervention without any context about which causal paths were active. For the full picture we need a second quantity, the probability of sufficiency.

Definition 8 (Probability of Sufficiency). *Let the probability of sufficiency be the probability that the outcome $Y = y^*$ would have occurred in the presence of $C = c^*$, given that neither $C = c^*$ nor $Y = y^*$ occurred:*

$$PS(C = c^*, Y = y^*) = P(Y_{c'} = y^* \mid C = c', Y \neq y^*)$$

where $c' \neq c^*$ and $Y_{c'}$ is the value of Y after intervening on C with c' .

To query whether the factual cause was sufficient, we condition on the counterfactual: we assume the cause did *not* take its factual value and the outcome did *not* occur, then intervene to restore the factual cause value and ask whether the outcome now occurs. Intuitively, this asks: if the cause had been absent and things had gone differently, would reinstating the cause have brought the outcome about? Let’s look at the resulting values.

We illustrate PS for Alice’s gender. Following the same convention as for PN , the individual-level computation additionally conditions on Alice’s credit ($credit = bad$): we fix her credit but flip the cause and outcome to their non-factual values ($gender = male$, $loan = T$), then intervene to restore $gender = female$ and ask whether the loan is again denied. Both check branches yield $loan = F$ (Appendix A, §A.2), so $PS(gender = female, loan = F \mid Alice) = 1$. The remaining individual values, along with population-level values for comparison, appear in Table 1. Two cases warrant comment: Bob’s gender PS is undefined because the required conditioning event ($gender = female$, $credit = bad$, $loan = T$) has probability zero in the model; Bob’s credit PS is 0.955 rather than 1 because the model includes a 5% exception rate for males who fail the credit check.

To somehow combine insights provided by the two scores, we would like to find causes that are both necessary and sufficient. This notion is formalized as follows:

Definition 9 (Probability of necessity and sufficiency). *Let the probability of necessity and sufficiency be the probability that the outcome $Y = y^*$ would have occurred in the presence of $C = c^*$ and that it would not have occurred under the intervention $C = c'$:*

$$PNS(C = c^*, Y = y^*) = P(Y_{c^*} = y^*, Y_{c'} \neq y^*).$$

This quantity factorises into a weighted sum of PN and PS by the consistency rule of counterfactuals [Pearl, 2009, §9.2]:

$$PNS(C = c^*, Y = y^*) = P(C = c^*, Y = y^*) PN(C = c^*, Y = y^*) + P(C = c', Y \neq y^*) PS(C = c^*, Y = y^*).$$

Table 1: PN , PS , and PNS for Alice and Bob in Example 6, with $loan = F$ as the outcome. “Population” values condition only on the cause–outcome pair prescribed by the definition; “Individual” values additionally condition on the subject’s other factual variable (e.g. for Alice’s gender, on $credit = bad$). A dash marks an entry that is undefined because the conditioning premises are inconsistent with the model.

Variable	Level	Alice			Bob		
		PN	PS	PNS	PN	PS	PNS
<i>gender</i>	Population	0.43	0.83	0.39	0.02	0.10	0.01
<i>gender</i>	Individual	0.045	1.00	0.045	0	—	0
<i>credit</i>	Population	0.53	0.96	0.52	0.53	0.96	0.52
<i>credit</i>	Individual	0.18	1.00	0.18	0.90	0.955	0.86

The population- and individual-level PNS values appear in the rightmost column group of Table 1. These values capture the relative causal impact of *gender* and *credit* but are not entirely satisfying. Table 1 reveals the problem directly: for Alice, $PNS(gender) = 0.045 < PNS(credit) = 0.18$, so PNS ranks credit as more causally responsible than gender. Yet the bank never evaluated Alice’s credit: the *check* variable was *F* throughout. PNS has no mechanism to register this: it asks counterfactually what would happen if credit were different, without fixing *check-failed*, the mediating variable that blocked credit’s causal path.

As an attempt to help Alice better understand her particular situation, we might consider a modification of PNS with additional conditioning:

$$PNS_c(C = c^*, Y = y^* | Z = z^*) = P(Y_{c^*} = y^*, Y_{c'} \neq y^* | Z = z^*)$$

Using this definition, we can compute $PNS_c(\text{gender} = \text{female}, \text{loan} = F | \text{credit} = \text{bad}) = 0.045$ and $PNS_c(\text{credit} = \text{bad}, \text{loan} = F | \text{gender} = \text{female}) = 0.18$. Conditioning does not solve the problem described above (Alice’s credit was in reality never evaluated) because the interventions override the mediating variable that tracks whether her credit has been checked. When we compute $PNS_c(\text{credit} = \text{bad}, \text{loan} = F | \text{gender} = \text{female})$, conditioning on gender being female is consistent with the factual, but when we then intervene to set credit to good, this intervention propagates downstream and changes *check-failed*, overriding what we learned from conditioning on Alice’s factual outcome. The context we fixed is undone by the very intervention meant to probe it. We will not consider this use of conditioning further, and this observation motivates the need for intervention-based (as opposed to conditioning-based) context-sensitivity, which is what the witness mechanism provides.

The propagation problem is not specific to PNS. The two most familiar causal quantities in the ML and policy literature are the average treatment effect $ATE(T) = \mathbb{E}[Y | do(T=1)] - \mathbb{E}[Y | do(T=0)]$ and the conditional version $CATE(T | X=x) = \mathbb{E}[Y | do(T=1), X=x] - \mathbb{E}[Y | do(T=0), X=x]$. Computed on our stochastic OBCB, population ATE ranks credit (0.518) above gender (0.383), mirroring the population PN ranking. At the individual level for Alice, $CATE(\text{credit} | \text{gender}=\text{female}) = 0.18$ and $CATE(\text{gender} | \text{credit}=\text{bad}) = 0.045$. These are exactly Alice’s individual PNS values: with binary outcomes and $PS = 1$ in the relevant branches, CATE for a single treatment collapses to PNS_c , so the same *check-failed* propagation that undermined PNS_c undermines CATE. A second, sharper failure follows from CATE’s inability to condition on the factual treatment value: $CATE(\text{gender} | \text{credit}=\text{bad}) = 0.045$ is identical for Alice and Bob even though one’s gender carried her denial outright while the other’s merely opened the credit check that then rejected him. The structural limitation cuts across PNS, ATE, and CATE alike, which is why the witness mechanism in Section 3 is not just a fix for PNS but a general repair for the family of methods that average counterfactual effects without intervention-based context-sensitivity.

2.3 Path Forward

The two predecessors examined in this section have complementary strengths and blind spots. Halpern’s actual causality is context-sensitive: by holding witnesses fixed at their factual values, it correctly diagnoses Alice’s case, attributing her denial to gender rather than credit. But in its standard form it is restricted to deterministic models, it requires a brute-force search over witness sets, and it has no native treatment of continuous variables. Pearl’s PN/PS/PNS make the converse trade-off. They scale naturally to probabilistic models, because every counterfactual reduces to an expectation under the noise distribution. But they have no analogue of the witness mechanism: every intervention propagates through the structural equations and overrides whatever mediator value the factual context fixed. Table 1 shows the symptom: for Alice, $PNS(\text{credit}) > PNS(\text{gender})$, even though the credit check was never performed.

Conditioning is not a workaround: we saw with PNS_c that context fixed by observation does not survive the subsequent intervention, since the credit intervention propagates downstream and overrides the *check-failed* value that conditioning had fixed. Any replacement therefore needs intervention-based context-sensitivity, holding the relevant mediators fixed by intervention. A satisfactory replacement must combine three ingredients. First, from actual causality, the witness mechanism: a way to hold relevant context fixed by intervention so that the structural information about which causal paths were active survives the counterfactual evaluation. Second, from PN/PS/PNS, the probabilistic vocabulary: counterfactuals expressed as expectations under the noise distribution, so that the attribution lives on the same probability space as the model and naturally extends to continuous variables. Third, in place of brute-force enumeration over witness sets, an integral that can be estimated from samples even when the witness pool is large or the variables are continuous.

PCI provides this synthesis. To get some general intuition (we will revisit the example in more detail later), consider Alice’s case once more, beginning with gender. PCI considers counterfactual worlds in which gender is flipped to male while a sampled witness set is held at its factual values by intervention. For Alice, the witness sets that drive the attribution are those containing *check-failed*:

when this variable is held fixed at F — the value recording that the check was never performed — flipping gender to male leads to loan approval. Unlike under PNS_c , the gender intervention does not override the mediator, because the witness intervention keeps it pinned. The counterfactual shift is large, and gender receives high attribution.

For credit, the same mechanism yields the opposite verdict. The witness sets that contain *check-failed*= F now work against credit: holding that mediator fixed at F disconnects credit from the loan decision, because the bank’s policy is to reject unchecked applicants regardless of credit. The counterfactual shift in those worlds is zero, and credit receives low attribution. The intuition that Alice’s denial was a function of gender, not credit, is recovered.

Two features distinguish PCI from both predecessors. Unlike actual causality, it does not ask whether *some* witness set validates a counterfactual dependency; it integrates over a distribution of witness sets, weighting each by its probability. This replaces combinatorial enumeration with statistical estimation. Unlike PN/PS/PNS, it inherits the context-sensitivity of actual causality through this integration: causally inert variables are downweighted automatically, because almost every witness set leaves them disconnected from the outcome, while variables that drive the outcome are upweighted, because almost every witness set leaves them connected. The formal definitions in Section 3 make this expectation precise.

Before moving to the formal development, we flag the closest neighbour of PCI in the existing literature, so the later comparison does not come out of nowhere: Pearl’s *probability of actual causation* [Pearl, 2009, Def. 10.3.5], the posterior probability — given the observed evidence — that the noise lands in a state for which the actual-cause verdict holds. Because PCI proposes a seemingly similar bridge from structural information to a probabilistic score, and the two give the same rankings in typical simple cases, we defer a precise side-by-side comparison to Section 8, once enough machinery is on the page. (The formula and its Bayes-rule reading are stated there, in Eq. (20).)

With the limitations of both predecessors in mind, we turn in Section 3 to the formal development. The next section introduces PCI piece by piece — PSCM scaffolding, the variable selection distribution Γ , the alternative-value distribution Δ , the joint necessity-sufficiency measure, and the user-chosen causal impact function ci — and verifies on the same Alice/Bob example that the resulting attribution satisfies all seven desiderata stated at the start of that section.

3 Causal Impact: Definitions

Explanation and causal attribution seek to connect a potentially causal *event* $X = x$ to an outcome event $Y = y$.⁹ In this section, we present a formal development. To ground it, we return to Example 6 and use Alice and Bob as running illustrations throughout this section. Both are rejected for a loan; the question is which of their attributes (*gender* and *credit*) was causally responsible for each rejection, and to what degree. While the example is discrete, the machinery applies to continuous and mixed-type variables; a continuous illustration is given in Sections 4.3 and 5.

Roadmap of this section. The development proceeds in five steps: (1) write down a list of seven *desiderata* that any responsibility attribution for Alice and Bob should satisfy, so that the formal construction can be judged against an explicit target; (2) introduce the probabilistic structural causal model (PSCM) and interventions, the substrate on which everything else rests; (3) introduce the two distributions PCI exposes to the user: the variable selection distribution Γ over suspect/witness pairs and the alternative-value distribution Δ over counterfactual cause settings; (4) combine them into the joint necessity-sufficiency measure $P_k^{s,n}$ and the user-chosen causal impact function ci whose expectation is the PCI score, and show how the choice of ci generalises the framework from binary to continuous outcomes; (5) compute the score on the OCB example, verify that all seven desiderata hold, and contrast PCI’s necessity/sufficiency decomposition with Pearl’s PN/PS/PNS. The closing paragraph then points forward to Section 4, which positions PCI against SHAP and Causal SHAP.

Let $R(V \rightsquigarrow loan \mid person)$ denote the causal responsibility of variable V for the loan outcome at the individual level (R is a placeholder for any attribution score and should not be identified

⁹This is quite different from type-level causal queries, where one is interested in some group-level attribution, as when one estimates average treatment effect or conditional average treatment effect.

with a specific formal quantity; PCI interprets R as $\mathbb{E}[ci(Y^s, Y^n, y^*)]$, to be defined below, but the desiderata are method-agnostic). Attribution methods may produce signed values; the desiderata are stated in terms of absolute magnitudes $|R(\cdot)|$, allowing for directionality:

$$\begin{aligned}
\mathbf{D-A1} \quad & |R(\text{gender} \rightsquigarrow \text{loan} \mid \text{Alice})| > 0 \\
\mathbf{D-A2} \quad & |R(\text{credit} \rightsquigarrow \text{loan} \mid \text{Alice})| > 0 \\
\mathbf{D-A-rank} \quad & |R(\text{gender} \rightsquigarrow \text{loan} \mid \text{Alice})| > |R(\text{credit} \rightsquigarrow \text{loan} \mid \text{Alice})| \\
\mathbf{D-B1} \quad & |R(\text{gender} \rightsquigarrow \text{loan} \mid \text{Bob})| > 0 \\
\mathbf{D-B2} \quad & |R(\text{credit} \rightsquigarrow \text{loan} \mid \text{Bob})| > 0 \\
\mathbf{D-B-rank} \quad & |R(\text{credit} \rightsquigarrow \text{loan} \mid \text{Bob})| > |R(\text{gender} \rightsquigarrow \text{loan} \mid \text{Bob})| \\
\mathbf{D-comp} \quad & |R(\text{gender} \rightsquigarrow \text{loan} \mid \text{Alice})| > |R(\text{gender} \rightsquigarrow \text{loan} \mid \text{Bob})|
\end{aligned}$$

D-A1 and **D-A2** both require non-zero attribution for Alice because two distinct causal routes were available: gender filtered her out at the checking stage without her credit ever being evaluated, while bad credit would have mattered had she been checked. Neither is causally inert. **D-A-rank** is the substantive constraint: because Alice never underwent a credit evaluation, gender’s causal role is immediate, whereas credit’s is purely hypothetical. A method that ranks credit above gender for Alice is misattributing the cause of her rejection.

For Bob, **D-B2** reflects that the credit check occurred and directly produced the rejection: the causal chain $\text{credit} \rightarrow \text{check-failed} \rightarrow \text{loan}$ was activated. **D-B1** requires strictly positive attribution for *gender*: being male gave Bob a 90% probability of being checked, so gender opened the evaluation that rejected him. This indirect causal role is non-zero; a method returning exactly zero fails to capture it. **D-B-rank** corresponds to the intuition that credit is the proximate cause. **D-comp** captures the cross-individual asymmetry that is hardest for naive methods to express: gender is more causally responsible for Alice’s rejection than for Bob’s, because for Alice *gender*=female was sufficient on its own to produce the rejection — no evaluation took place, so credit was never a factor — while for Bob *gender*=male merely opened the evaluation that *credit*=bad then caused. We will verify at the end of this section that PCI satisfies all seven desiderata, while PN, PS, PNS fail some of them. Later, we will also revisit this example and its desiderata in the context of SHAP and Causal SHAP (Section 4).

We work within the context of probabilistic structural causal models [Pearl, 2009].

Definition 10 (Probabilistic Structural Causal Model (PSCM)). *A probabilistic structural causal model is a tuple*

$$M = \langle \mathbf{U}, \mathbf{V}, \mathbf{F}, P_{\mathbf{U}} \rangle,$$

where $\mathbf{V} = \{V_1, \dots, V_n\}$ is a finite set of endogenous variables, $\mathbf{U} = \{U_1, \dots, U_n\}$ is a set of exogenous variables, and $\mathbf{F} = \{f_1, \dots, f_n\}$ is a set of structural assignments

$$V_i := f_i(\mathbf{Pa}_i, U_i), \quad i = 1, \dots, n,$$

with $\mathbf{Pa}_i \subseteq \mathbf{V} \setminus \{V_i\}$ denoting the parents of V_i . The distribution $P_{\mathbf{U}}$ is a probability measure on $\text{dom}(\mathbf{U})$, and the assignments in $\mathbf{F}$ are assumed to be acyclic and to have a unique solution for $\mathbf{V}$ given $\mathbf{U}$.¹⁰

A PSCM encodes which variables influence which others, through what mechanisms, and subject to what randomness. The structural equations specify each variable as a deterministic function of its causal parents and local noise; the exogenous distribution captures all irreducible uncertainty in the system. Together they make every counterfactual question well-posed and computable.

The OCB model (Example 6) is a PSCM with endogenous variables

$$\mathbf{V} = \{\text{gender}, \text{credit}, \text{check}, \text{check-failed}, \text{loan}\},$$

¹⁰Working within the PSCM framework commits the practitioner to fully specified structural assignments $\mathbf{F}$ and an exogenous distribution $P_{\mathbf{U}}$, analogous to how Bayesian inference requires a likelihood and a prior. This is precisely what enables PCI to yield individual-level counterfactual attributions rather than population-level bounds, and once $\mathbf{F}$ and $P_{\mathbf{U}}$ are specified, sampling from $P_{\mathbf{U}}$ is a routine probabilistic inference operation.

and exogenous variables

$$\mathbf{U} = \{U_{gender}, U_{credit}, U_{check}, U_{loan}\}$$

which induce independent Bernoulli noise.

The structural assignments in the case at hand take the form $V_i = f_i(\mathbf{Pa}_i, U_i)$ promised by the PSCM definition above, with indicator-threshold mechanisms against uniform exogenous noise:

$$\begin{aligned} gender &= \mathbb{I}[U_g \leq \alpha_g], \\ credit &= \mathbb{I}[U_c \leq \alpha_c], \\ check &= \mathbb{I}[U_{ck} \leq \gamma_F(1 - gender) + \gamma_M gender], \\ check-failed &= check(1 - credit), \\ loan-if-checked &= \mathbb{I}[U_\ell \leq \text{loan-prob}(gender, check-failed)], \\ loan &= loan-if-checked \cdot check, \end{aligned}$$

with $U_g, U_c, U_{ck}, U_\ell \stackrel{\text{iid}}{\sim} \text{Unif}[0, 1]$ as the exogenous noise. For the running example we take $\alpha_g = \alpha_c = 0.5$, $\gamma_F = 0.2$, $\gamma_M = 0.9$, and the values of loan-prob given in Example 6.

Definition 11 (Interventions). *Given a model $M = \langle \mathbf{U}, \mathbf{V}, \mathbf{F}, P_{\mathbf{U}} \rangle$, an intervention on variables $\mathbf{X} = \{X_1, \dots, X_k\} \subseteq \mathbf{V}$ with values $\mathbf{x} = (x_1, \dots, x_k)$ replaces the structural assignments for X_i by $X_i := x_i$, where $i = 1, \dots, k$. We denote this intervention by either $[\mathbf{X} \leftarrow \mathbf{x}]$ or $\text{do}(\mathbf{X} = \mathbf{x})$.*

An intervention surgically replaces a variable's structural equation with a fixed constant, disconnecting it from its usual causes. The rest of the model propagates downstream as normal under the new assignment; the exogenous noise is unchanged.

In the OBCB model, $\text{do}(gender = 1)$ replaces Alice's gender assignment with male, leaving all other structural equations intact. The downstream effect propagates: *check* now draws from $\text{Bern}(0.9)$ rather than $\text{Bern}(0.2)$, so Alice would be checked in 90% of noise realizations instead of 20%. This is the counterfactual world in which we ask whether her loan outcome would have differed.

Because PCI supports arbitrary variable types, it will be helpful to define a measure theoretic object corresponding to interventional outcomes. Because outcomes are a deterministic function of exogenous noise, this will be a Dirac measure.

Definition 12 (Pointwise Interventional Law). *Let $M = \langle \mathbf{U}, \mathbf{V}, \mathbf{F}, P_{\mathbf{U}} \rangle$ be a probabilistic structural causal model and let $Y \in \mathbf{V}$. For an intervention $\text{do}(\mathbf{X} = \mathbf{x})$, let $Y' : \text{dom}(\mathbf{U}) \rightarrow \text{dom}(Y)$ denote the measurable map induced by the modified structural assignments. The pointwise interventional law of Y at fixed $\mathbf{u} \in \text{dom}(\mathbf{U})$ is the probability measure on $\text{dom}(Y)$ defined by*

$$P(Y \in A \mid \mathbf{u}, \text{do}(\mathbf{X} = \mathbf{x})) := \mathbb{I}\{Y'(\mathbf{u}) \in A\} = \delta_{Y'(\mathbf{u})}(A), \quad A \subseteq \text{dom}(Y).$$

The pointwise interventional law records the deterministic outcome of Y under $\text{do}(\mathbf{X} = \mathbf{x})$ for a fixed noise realization $\mathbf{u}$: since the modified structural equations make Y a function of $\mathbf{u}$ alone, its distribution collapses to a point mass. Integrating over $P_{\mathbf{U}}$ then recovers the full interventional distribution.

We define a *causal set* to be a set of assignments to the variables of a probabilistic structural causal model.

Definition 13 (Causal set). *Given a model $\langle \mathbf{U}, \mathbf{V}, \mathbf{F}, P_{\mathbf{U}} \rangle$, a causal set is a set of pairs $X = x$ where $X \in \mathbf{V}$ and $x \in \text{dom}(X)$.*

Informally, a causal set is a list of specific variable-value assignments, a precise statement of which variables are set to which values. It is the basic unit of intervention: from a causal set one can read off exactly what is being manipulated or held fixed.

In PCI, causal sets play four distinct roles, two on the cause side and two on the context side:

Suspects S. The full set of variables suspected to bear on the outcome. Subsets of **S** will be sampled and intervened on.

Active suspects $C \subseteq S$. The specific subset intervened on in a given search step, either to alternative values (necessity regime) or to factual values (sufficiency regime).

Witnesses W . The full set of context variables that may be held fixed at their factual values. Subsets of W will be sampled and held fixed.

Active witnesses $T \subseteq W$. The specific subset held fixed in a given search step. To avoid contradicting the active intervention, T is pruned so that $T \cap C = \emptyset$.

Why prune T and not C ? The conflict on $T \cap C$ has to be resolved in one direction or the other, and pruning the witnesses is the right choice for three reasons. First, C is the object of the test: it encodes the causal claim being evaluated, while T is the auxiliary device that holds context fixed. Pruning C would silently change the claim under test; pruning T leaves the claim intact and merely weakens the context held fixed — a weaker probe of the same hypothesis, not a different one. Second, the convention extends cleanly to the limit $C = V$, where every variable is a suspect: the pruned T collapses to $\emptyset$ and we recover the unconditional but-for test, the right baseline behaviour. Pruning C instead would leave this limit undefined whenever any witness is sampled. Third, the convention inherits directly from Halpern’s actual-cause definition (Definition 5), which makes $T \cap C = \emptyset$ part of the definition; pruning post-sampling is the operational implementation of that constraint and lets us specify Γ on the unconstrained product $2^S \times 2^W$ rather than baking disjointness into the prior.

To evaluate the causal impact of a variable on an outcome, we consider two interventional regimes: the *necessity regime*, in which we intervene on the variables $C \subseteq S$ to have an alternative value, and the *sufficiency regime*, in which we intervene so that C takes on factual values. Moreover, we hold some elements $T \subseteq W$ of the context fixed (i.e., intervened on to retain factual values, despite intervention on C). This is exactly what makes the necessity component *context-sensitive* in Halpern’s sense [Halpern, 2016]: the alternative cause is evaluated against a held-fixed witness configuration, not in isolation. We will reuse this terminology throughout: “context-sensitive necessity” refers to the necessity regime evaluated under an active witness set T . When looking for an explanation, we perform a search across possible cause sets and possible contexts. Complete model samples in these regimes will be called *necessity worlds* and *sufficiency worlds*, respectively.

The intuition is that the suspect set S names every variable that could plausibly have caused the outcome; the witness set W names context variables to be selectively held fixed, isolating the causal channels under investigation. Intervening on suspects tests what would have happened under an alternative cause in the necessity worlds and under the factual values in the sufficiency worlds; fixing witnesses blocks confounding paths that would otherwise absorb the effect. Responsibility attribution increases when the outcome is far from the factual value in the necessity world and decreases when it is far from the factual value in the sufficiency world.

For both Alice (female, bad credit, loan denied) and Bob (male, bad credit, loan denied) we set $S = \{gender, credit\}$ and $W = \{check-failed\}$. The witness *check-failed* records whether the applicant was checked and had bad credit; holding it at its factual value isolates the causal path from *gender* or *credit* to *loan* independently of the credit-check bottleneck. Without this witness, intervening on *credit* for Alice would leave 80% of noise realizations unaffected (those in which she is never checked), making credit’s causal role invisible. The witness makes the bottleneck explicit and manipulable.

Moving on with the general framework, we will be searching for causes by intervening on subsets of suspects and holding fixed subsets of witnesses. To make this search tractable, we need a way to prioritize which subsets to consider.

Definition 14 (Variable Selection Distribution). *Let X be a finite set of random variables. A variable selection distribution Γ is a probability distribution on 2^X . We write $Y \sim \Gamma$ to mean Y is a random subset of X drawn from Γ , with probability mass function $p^\Gamma(y) = \mathbb{P}(Y = y)$ for $y \subseteq X$.*

The variable selection distribution Γ encodes prior beliefs about which subsets of suspects are worth examining. A uniform distribution treats all suspect subsets as equally plausible; a cardinality-constrained choice focuses the search on interactions of a particular order, analogous to choosing which interaction terms to include in a regression. More complex distributions could encode domain knowledge about which variables are more likely to interact, or could be learned from data.

In particular, we will be using variable selection distributions for the powerset of causal suspects S and for the powerset of witnesses W . One simple pattern of suspect and witness selection can

be derived by setting size limits on the number of activated suspects or witnesses, and sampling uniformly otherwise.

Example 15 (Cardinality-Constrained Uniform Selection). *Let $\mathbf{X}$ be a candidate set of random variables and let $k \sim \text{Unif}\{J, \dots, K\}$ with $1 \leq J \leq K \leq |\mathbf{X}|$. A variable selection $\mathbf{Y} \sim \Gamma(2^{\mathbf{X}})$ is sampled from a cardinality-constrained uniform selection distribution $\Gamma(2^{\mathbf{X}})$ with probability mass function*

$$p^\Gamma(\mathbf{Y}) \propto \mathbb{I}[|\mathbf{Y}| = k].$$

*The parameters J and K determine the cardinality range of the search; specific choices connect PCI to existing methods and enable efficient Monte Carlo approximation.*¹¹

In PCI the variable selection distribution Γ is a probability mass function on $2^{\mathbf{S}} \times 2^{\mathbf{W}}$, governing the joint choice of an active suspect set $\mathbf{C} \subseteq \mathbf{S}$ and an active witness set $\mathbf{T} \subseteq \mathbf{W}$. The recommended construction samples suspects and witnesses separately and then enforces the constraint that no variable plays both roles in the same draw: pick a marginal Γ_s on $2^{\mathbf{S}}$ and a marginal Γ_w on $2^{\mathbf{W}}$, draw $\mathbf{C} \sim \Gamma_s$ and $\mathbf{T} \sim \Gamma_w$ independently, and *reject* any draw with $\mathbf{T} \cap \mathbf{C} \neq \emptyset$, resampling until the constraint holds. Equivalently,

$$p^\Gamma(\mathbf{C}, \mathbf{T}) \propto p_s^\Gamma(\mathbf{C}) p_w^\Gamma(\mathbf{T}) \mathbb{I}[\mathbf{T} \cap \mathbf{C} = \emptyset].$$

Throughout, Γ_s and Γ_w are abbreviations for the sampling marginals of this construction; the formal primitive in Definition 18 is the joint Γ .

For the OBCB analysis, with $|\mathbf{S}| = 2$, we use a Γ built from Γ_s uniform over the three non-empty subsets of $\mathbf{S}$ ($\{gender\}$, $\{credit\}$, $\{gender, credit\}$, each with probability $1/3$) and Γ_w uniform over $\{\emptyset, \{check-failed\}\}$, each with probability $1/2$. Since no element appears in both $\mathbf{S}$ and $\mathbf{W}$, the rejection step is vacuous and Γ is exactly the product of marginals $p_s^\Gamma \cdot p_w^\Gamma$. In larger suspect sets the choice of cardinality range becomes substantive: sampling uniformly from the full powerset yields an expected intervention size of $|\mathbf{S}|/2$, which spreads attribution across large coalitions and may make individual feature contributions harder to isolate; we return to this issue in Section 9, where the cardinality cap is treated as a tunable hyperparameter and a dynamic-set-sizes ablation measures the empirical cost of bounding it.

The choice of Γ is a design decision that shapes the search process and the resulting attributions. As a default, a uniform Γ over the full powerset is the least-informative starting point, assigning equal weight to all subset sizes, analogous to including all interaction orders in a regression. There is no universally principled criterion for restricting the cardinality range; the honest advice is to treat J and K as explicit design choices. Two practical considerations bear on this choice, however: with $K = 1$, attribution scores are based on singleton interventions and straightforward to communicate to stakeholders; higher K averages over joint interventions on X_k with other suspects simultaneously, which is harder to explain and, in large or structurally complex models, may amplify estimation noise rather than capturing genuine higher-order causal structure. For applications where attribution has serious consequences, we recommend a cardinality sensitivity analysis: if rankings are stable as K increases from 1 to $|\mathbf{S}|$, the simpler choice is supported; if they diverge, higher-order interactions

¹¹When $J = 1$ and $K = |\mathbf{X}|$, this distribution coincides with the Shapley weighting kernel, which samples coalition size uniformly and then selects a coalition uniformly at that size [Shapley, 1953]. The parameters J and K allow the practitioner to restrict the search to cause sets of a specific cardinality range: setting $J = K = 1$ confines attribution to individual variables, while $J = 2$, $K = 3$ focuses on small multi-variable interactions. Moreover, whereas exact Shapley computation requires summing over all $2^{|\mathbf{X}|}$ coalitions, PCI treats Γ as a generative model: approximation proceeds by sampling subsets from Γ and outcomes from P_U , making Monte Carlo estimation a native feature of the framework rather than an external algorithmic layer. We note that the weighting alone does not distinguish PCI from SHAP at $J = 1$, $K = |\mathbf{X}|$: the subset weights are identical. The distinction lies in the integrand, and it matters for what is being evaluated and what results follow. SHAP averages marginal contributions using the observational conditional $\mathbb{E}[f(\mathbf{X}_S = \mathbf{x}_S, \mathbf{X}_{\bar{S}})]$, marginalising over the empirical distribution of non-suspect features; this means a variable that is merely correlated with a cause, but not itself causally active, can receive positive attribution. PCI instead evaluates counterfactual outcomes $P(Y \in A \mid \mathbf{u}, \text{do}(\mathbf{C} = \mathbf{c}', \mathbf{T} = \mathbf{t}^*))$ computed via the structural model, so attribution flows only along genuine causal paths. The witness mechanism adds a further distinction: by holding intermediate variables fixed, PCI can resolve overdetermination and undercutting scenarios where SHAP assigns equal or incorrect attribution. The choice of J and K remains a design decision; practitioners can set $J = K = 1$ for single-variable attribution or allow higher-order interactions as discussed above. A detailed comparison of PCI, SHAP, and Causal SHAP on concrete examples is given in Section 4.

are materially relevant and the full powerset should be used. The key advantage of PCI is that this commitment is made transparently rather than absorbed into an implicit default.¹²

Causal attribution hinges on questions such as “if X had, contrary to fact, been fixed to $x' \neq x$ instead of x , what would the outcome have been?” For non-binary causes, this is ambiguous. To resolve this in a way that does not require committing to a single, contrastive causal event [Kawakami et al., 2024], we rely on a *distribution* over alternative values.

Definition 16 (Alternative Value Distribution). *Let $\mathbf{X}$ be a set of random variables with domain $\text{dom}(\mathbf{X})$ and let $\mathbf{X} = \mathbf{x} \in \text{dom}(\mathbf{X})$ be a factual event. An alternative value distribution $\Delta(\mathbf{x})$ is a probability measure on $\text{dom}(\mathbf{X})$ such that $\mathbf{x} \notin \text{supp}(\Delta(\mathbf{x}))$.*

Informally, $\Delta(\mathbf{x})$ specifies what counts as a genuinely different value for a suspect variable. The support condition ensures no mass falls on the factual value $\mathbf{x}$ itself, so every sampled alternative is a real counterfactual departure rather than a noisy copy of what actually happened.

Three properties guide the choice of Δ . *Plausibility under M* : the alternative should be a value the structural model could plausibly produce given the context, so the necessity test interrogates a counterfactual the model has anything to say about. *Outcome-independence*: the distribution over $\mathbf{x}'$ should not be informed by the outcome whose causes we are explaining, on pain of circularity. *Context-sensitivity*: $\mathbf{x}'$ should reflect the rest of the factual scenario, so the necessity test is individual-level rather than population-level. The construction we use throughout is the structural-model posterior of $\mathbf{X}$ conditioned on upstream factual values, optionally excised near the factual value (Example 17 below); this is the natural choice satisfying all three.

Upstream-only conditioning operationalises the intuition shared with the descriptive-normality literature [Halpern and Hitchcock, 2015, Icard et al., 2017]: factual upstream values play the role of the *default* against which counterfactual alternatives are graded. PCI does not aim to resolve the specific counterexamples motivating that literature; the link is at the level of *why* a non-uniform Δ matters at all. Since Δ is user-pluggable, other constructions are available when domain knowledge motivates them: a uniform distribution over $\text{dom}(\mathbf{X})$ trades plausibility for breadth; the empirical marginal $p(\mathbf{x}')$ trades context-sensitivity for simplicity; a fully conditional posterior on every observable (including downstream evidence) trades outcome-independence for sharper conditioning. Each reads as an informed relaxation of one of the three properties above, not an unconsidered default — exactly the user-facing knob the framework is designed to expose.

One natural alternative value distribution results from taking the one currently embodied in the model and excising it around the factual setting. Then, the semantics for “alternative value” means “a sufficiently different value, following the original distribution otherwise”. In the discrete case, of course, the analogue is simply the probability mass function conditioned on the factual value not holding.

Example 17 (ϵ -Excised Distribution). *Assume $\mathbf{X}$ is $\mathbb{R}^d$ -valued with density p . Given a factual event $\mathbf{X} = \mathbf{x}$, define the ϵ -excised alternative value distribution $\Delta_\epsilon(\mathbf{x})$ as the probability measure on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ with density*

$$p_{\Delta_\epsilon}(\mathbf{x}' | \mathbf{x}) \propto \begin{cases} 0, & \mathbf{x}' \in B_\epsilon(\mathbf{x}), \\ p(\mathbf{x}'), & \text{otherwise,} \end{cases}$$

where $B_\epsilon(\mathbf{x})$ denotes the ϵ -ball centered at $\mathbf{x}$. The support is therefore $\text{supp}(\Delta_\epsilon(\mathbf{x})) = \text{supp}(p) \setminus B_\epsilon(\mathbf{x})$ — not just the factual point $\mathbf{x}$ but its entire ϵ -neighbourhood is excluded.

Informally, the alternative value distribution Δ operationalises what “counterfactually different” means for each suspect variable: it specifies which alternative values are considered when asking whether the outcome would have changed under an alternative cause.

¹²SHAP faces an analogous decision without flagging it as one: by averaging over all $2^{|N|}$ feature coalitions, plain SHAP and Causal SHAP commit to the maximum interaction order $K = |N|$ as an unstated default, with no diagnostic offered for whether that level of joint conditioning is right for a given model. Restricting SHAP to $K < |N|$ would break the efficiency axiom, so the choice is hardcoded into the framework rather than treated as a tunable. PCI surfaces the same decision as an explicit knob.

In the OBCB model both *gender* and *credit* are binary, so Δ is trivial. For Alice, the only alternative to *gender* = 0 (female) is *gender* = 1 (male), and the only alternative to *credit* = 0 (bad) is *credit* = 1 (good); for Bob, the only alternative to *gender* = 1 is *gender* = 0. Accordingly, $\Delta(\text{gender} = 0)$ is a point mass on *gender* = 1 and $\Delta(\text{credit} = 0)$ a point mass on *credit* = 1: no distributional choice is required. The binary case is exactly where Δ is unambiguous; it is for continuous suspects such as income or loan amount that the ϵ -excised distribution becomes essential, forcing an explicit commitment to what counts as a meaningfully different value.

Why ϵ -excision? Any method that asks “what would happen if $\mathbf{X}$ had been different” must operationalise what *different* means. Three natural choices for Δ are available. The observational marginal $p(\mathbf{x}')$ — as in kernel SHAP [Lundberg and Lee, 2017] — places mass near the factual value $\mathbf{x}$, conflating the necessity question (“what if $\mathbf{x}$ had genuinely differed?”) with evaluating essentially the same setting. The fully conditional posterior $p(x'_k | \mathbf{x}_{-k})$ conditions on all other observed features, potentially including descendants of X_k , which imposes structural constraints that may rule out otherwise plausible counterfactuals, and still concentrates mass near x_k when features are correlated. The ϵ -excised distribution occupies the middle ground: it draws from the marginal informed by upstream variables only, excising the factual neighbourhood to guarantee genuine departure without downstream conditioning.

The choice parallels the Region of Practical Equivalence (ROPE) in Bayesian hypothesis testing [Kruschke, 2018]: just as ROPE forces an explicit decision about what difference is large enough to matter, ϵ -excision forces an explicit decision about what counterfactual is sufficiently far from the factual to count as genuinely alternative. The practitioner makes this commitment regardless of method; the only question is whether it is made transparently or absorbed into a default. When X is standardised to $\mathcal{N}(0, 1)$ the choice acquires a direct probabilistic interpretation: the excised mass $\Phi(x^* + \epsilon) - \Phi(x^* - \epsilon)$ is the fraction of the distribution ruled out as insufficiently different from x^* . At $x^* = 0$, $\epsilon = 0.2, 0.5, 0.8$ excise approximately 16%, 38%, 58% of the distribution, corresponding to small, medium, and large effect thresholds in the sense of Cohen [1988]. Alternatively, ϵ can reflect measurement precision or a decision-relevant threshold; sensitivity analysis across ϵ values is advisable when the appropriate scale is unclear. For concrete examples of how SHAP’s implicit choice leads to problematic attributions, see Section 4.

Where we are. The PSCM and intervention apparatus give us a notion of counterfactual outcome for any fixed noise realisation $\mathbf{u}$; Γ tells us which suspect/witness configurations to ask about; Δ tells us which non-factual values to substitute when evaluating necessity. What remains is to plug those distributions into the counterfactual mechanics and read out a number. The next two definitions do exactly that: first the joint necessity-sufficiency measure $P_k^{s,n}$, then its expectation against a user-chosen impact function *ci*, with the OBCB scores and the desiderata verification immediately after.

We are now ready to state the main definitions of PCI. The core idea is to measure two complementary things for a variable of interest X_k : (a) how much X_k ’s factual value contributes to producing y^* , the *sufficiency* question, and (b) how much replacing X_k ’s value with alternatives would shift the outcome away from y^* , the *necessity* question. We capture both by constructing two counterfactual outcome distributions conditioned on the same noise realization $\mathbf{u}$, then marginalising their product over P_U . Because PCI is not constrained to particular variable types, we work in general measure-theoretic terms supporting combinations of continuous and discrete variables.

Definition 18 (Joint Necessity and Sufficiency Measure). *Let $M = \langle \mathbf{U}, \mathbf{V}, \mathbf{F}, P_U \rangle$ be a probabilistic structural causal model and Y an outcome variable with domain $\text{dom}(Y)$. Let p^Γ be the mass function of a variable selection distribution on $2^{\mathbf{S}} \times 2^{\mathbf{W}}$, and let $\Delta(\mathbf{s}^*)$ be an alternative value distribution on $\text{dom}(\mathbf{S})$. Let $(\mathbf{S}, \mathbf{W}) = (\mathbf{s}^*, \mathbf{w}^*)$ be a factual event.*

For a variable of interest X_k , let $2_k^{\mathbf{S}} := \{\mathbf{C} \subseteq \mathbf{S} : X_k \in \mathbf{C}\}$, and for each $\mathbf{C} \subseteq \mathbf{S}$ let $\Delta_{\mathbf{C}}(\mathbf{s}^)$ denote the pushforward of $\Delta(\mathbf{s}^*)$ under the restriction map $\mathbf{s}' \mapsto \mathbf{s}'|_{\mathbf{C}}$.¹³ For any measurable $A \subseteq \text{dom}(Y)$ and exogenous realization $\mathbf{u}$, define three sub-probability measures on $\text{dom}(Y)$ (the first two) and $\text{dom}(Y) \times \text{dom}(Y)$ (the third).*

¹³For an assignment $\mathbf{x}$ to a collection of variables $\mathbf{X}$ and any subset $\mathbf{Y} \subseteq \mathbf{X}$, the restriction $\mathbf{x}|_{\mathbf{Y}}$ denotes the sub-assignment of $\mathbf{x}$ to the variables in $\mathbf{Y}$. Note that the restricted marginal $\Delta_{\mathbf{C}}(\mathbf{s}^*)$ may place positive mass on $\mathbf{s}^*|_{\mathbf{C}}$ even though $\Delta(\mathbf{s}^*)$ excludes the joint factual $\mathbf{s}^*$; the genuine-departure guarantee is on the joint alternative, not on each restricted coordinate.

(i) **Sufficiency measure.** $P_k^s(\cdot \mid \mathbf{u})$ intervenes on each active suspect set $\mathbf{C}$ to its factual values $\mathbf{s}^* \mid_{\mathbf{C}}$, holds the active witnesses $\mathbf{T}$ fixed, and takes the Γ -weighted sum of $P(Y \in A \mid \mathbf{u}, \text{do}(\cdot))$ under factual cause restoration:

$$P_k^s(A \mid \mathbf{u}) = \sum_{\substack{(\mathbf{C}, \mathbf{T}) \\ X_k \in \mathbf{C}}} P(Y \in A \mid \mathbf{u}, \text{do}(\mathbf{C} = \mathbf{s}^* \mid_{\mathbf{C}}, \mathbf{T} = \mathbf{w}^* \mid_{\mathbf{T}})) p^\Gamma(\mathbf{C}, \mathbf{T}). \quad (1)$$

(ii) **Necessity measure.** $P_k^n(\cdot \mid \mathbf{u})$ instead integrates over alternative values $\mathbf{c}'$ for $\mathbf{C}$ (with $\mathbf{T}$ still pinned at factual values), and takes the Γ -weighted average over alternative cause values:

$$P_k^n(A \mid \mathbf{u}) = \sum_{\substack{(\mathbf{C}, \mathbf{T}) \\ X_k \in \mathbf{C}}} \int P(Y \in A \mid \mathbf{u}, \text{do}(\mathbf{C} = \mathbf{c}', \mathbf{T} = \mathbf{w}^* \mid_{\mathbf{T}})) \Delta_{\mathbf{C}}(\mathbf{s}^*)(d\mathbf{c}') p^\Gamma(\mathbf{C}, \mathbf{T}). \quad (2)$$

(iii) **Joint necessity-sufficiency measure.** On product rectangles $A \times B \subseteq \text{dom}(Y) \times \text{dom}(Y)$, set

$$P_k^{s,n}(A \times B) := \int P_k^s(A \mid \mathbf{u}) P_k^n(B \mid \mathbf{u}) P_{\mathbf{U}}(d\mathbf{u}). \quad (3)$$

This rectangle-by-rectangle specification extends uniquely (by Carathéodory's extension theorem) to a sub-probability measure on the product σ -algebra of $\text{dom}(Y) \times \text{dom}(Y)$, with total mass $\Pr_{\Gamma}[X_k \in \mathbf{C}]^2 \leq 1$.

Informally: $P_k^s(\cdot \mid \mathbf{u})$ is the distribution of Y in the *sufficiency world*: restore the factual value of a candidate cause set $\mathbf{C}$ containing X_k , hold the witness set $\mathbf{T}$ at its factual values, and observe where the outcome lands. $P_k^n(\cdot \mid \mathbf{u})$ is the distribution of Y in the *necessity world*: replace $\mathbf{C}$'s values with alternatives drawn from Δ , hold $\mathbf{T}$ fixed, and ask how the outcome shifts. The sum over suspect-containing pairs $\{(\mathbf{C}, \mathbf{T}) : X_k \in \mathbf{C}\}$ weights each configuration by its Γ -mass: more focused interventions receive whatever weight the practitioner assigns them; configurations with $X_k \notin \mathbf{C}$ would contribute 0 to the attribution and are absent from the sum, so P_k^s and P_k^n are sub-probability measures with total mass $\Pr_{\Gamma}[X_k \in \mathbf{C}] \leq 1$ rather than renormalized expectations. In the OCB example, for $X_k = \text{gender}$ under a noise realization in which Alice is credit-checked: the sufficiency world restores $\text{gender} = \text{female}$ while holding check-failed fixed; the necessity world substitutes $\text{gender} = \text{male}$ under the same noise.

The product decomposition in $P_k^{s,n}$ is legitimate because, conditional on $\mathbf{u}$, the sufficiency and necessity interventions operate in separate counterfactual regimes with no causal connection: $\text{do}(\mathbf{C} = \mathbf{c}^*)$ and $\text{do}(\mathbf{C} = \mathbf{c}')$ act on the same noise but in distinct hypothetical worlds, so Y^s and Y^n are conditionally independent given $\mathbf{u}$, and $P(Y^s \in A, Y^n \in B \mid \mathbf{u}) = P_k^s(A \mid \mathbf{u}) P_k^n(B \mid \mathbf{u})$.

With the necessity-sufficiency measure fully defined, we are ready to characterize the causal impact of an event X_k . In particular, we will proceed by defining an arbitrary *causal impact function*, whose expectation over the necessity-sufficiency measure determines the causal impact. We first proceed with a formal definition of the causal impact measure, followed by several intuitive candidates for the causal impact function.

Definition 19 (Causal Impact Function and Its Expectation). *Let $y^* \in \text{dom}(Y)$ denote the factual outcome value, and let $Y^s, Y^n \sim P_k^{s,n}$. For any measurable function $ci : \text{dom}(Y)^3 \rightarrow \mathbb{R}$, the (expected) causal impact of X_k is defined as*

$$\mathbb{E}[ci(Y^s, Y^n, y^*)] := \int \int ci(y^s, y^n, y^*) P_k^{s,n}(dy^s, dy^n). \quad (4)$$

The integral is taken against the sub-probability $P_k^{s,n}$ directly, not against its renormalisation; the symbol $\mathbb{E}[\cdot]$ is shorthand for this sub-probability integral throughout. Suspect configurations with $X_k \notin \mathbf{C}$ contribute zero mass to P_k^s and P_k^n (and hence to $P_k^{s,n}$), correctly capturing that those configurations make no claim on X_k 's attribution.

Informally, ci is a user-defined scoring rule that translates the joint sufficiency–necessity measure into a single scalar. The choice is not arbitrary: the Absolute Difference example below is an instance of L_1 distance between outcomes, the same metric underlying Average Treatment Effect,

Conditional Treatment Effect, and SHAP. The binary indicator score recovers this as a special case when outcomes are binary: indicators are L_1 on $\{0, 1\}$. Practitioners with reasons to prefer L_2 or another distance measure may substitute it freely; the framework places no restriction on ci beyond measurability.

For the OCB analysis we instantiate ci as the PNS binary score defined in the following example, with $y^* = loan = 0$. This is the choice that produces the values in Table 2.

Example 20 (PNS for Binary Outcomes). *Assume the outcome Y is binary with $\text{dom}(Y) = \{0, 1\}$, and let $y^* = 1$ denote the factual outcome. Let Y^s and Y^n denote the sufficiency and necessity outcome variables. Define the causal impact score*

$$ci(y^s, y^n, y^*) = \mathbb{I}\{y^n \neq y^*\} \mathbb{I}\{y^s = y^*\}.$$

Then the expected causal impact reduces to

$$\mathbb{E}[ci(Y^s, Y^n, 1)] = \mathbb{P}(Y^n = 0, Y^s = 1),$$

which conceptually coincides with Pearl’s probability of necessity and sufficiency (PNS). The alignment is exact when $\mathbf{S} = \{X_k\}$, $\mathbf{W} = \emptyset$, and $\text{dom}(X_k) = \{0, 1\}$ where, factually, $x_k^* = 1$.

We now illustrate the full computation for Alice’s gender attribution using this ci . With $\mathbf{S} = \{gender, credit\}$, $\mathbf{W} = \{check-failed\}$, Γ_s and Γ_w uniform, Δ a point mass on the alternative binary value, and $y^* = loan = 0$, the exogenous noise U_{check} falls into three regions:

- u_1 (prob. 0.2): Alice is checked regardless of gender; $check-failed = 1$ factually.
- u_2 (prob. 0.7): Alice is not checked as female but would be as male; $check-failed = 0$ factually.
- u_3 (prob. 0.1): Alice is not checked regardless of gender.

In all regions, restoring Alice’s factual values always yields a denial, so $P_k^s(\{0\} \mid \mathbf{u}) = \frac{2}{3}$ everywhere (four suspect-containing (C, T) combinations each with $p^\Gamma(C, T) = \frac{1}{3} \cdot \frac{1}{2} = \frac{1}{6}$ — since $\mathbf{S} \cap \mathbf{W} = \emptyset$ here, rejection is vacuous and Γ is just the product of marginals — each contributing probability 1). The necessity measure varies.

In u_1 , the witness holds $check-failed = 1$. The four (C, T) contributions, each weighted $\frac{1}{6}$, are:

- $\mathbf{C} = \{gender\}$, $T = \emptyset$: male, $check-failed = 1$ propagates structurally $\Rightarrow P(loan = 1) = 0.05$.
- $\mathbf{C} = \{gender\}$, $T = \{check-failed\}$: male, witness holds failure $\Rightarrow P(loan = 1) = 0.05$.
- $\mathbf{C} = \{gender, credit\}$, $T = \emptyset$: male and good credit, check passes $\Rightarrow P(loan = 1) = 1.0$.
- $\mathbf{C} = \{gender, credit\}$, $T = \{check-failed\}$: male and good credit, but witness holds failure $\Rightarrow P(loan = 1) = 0.05$.

$$P_k^n(\{1\} \mid u_1) = \frac{1}{6}(0.05 + 0.05 + 1.0 + 0.05) \approx 0.192.$$

In u_2 , the witness holds $check-failed = 0$. The four contributions:

- $\mathbf{C} = \{gender\}$, $T = \emptyset$: male would be checked; $check-failed = 1$ propagates structurally $\Rightarrow P(loan = 1) = 0.05$.
- $\mathbf{C} = \{gender\}$, $T = \{check-failed\}$: male, witness holds $check-failed = 0$, check passes $\Rightarrow P(loan = 1) = 1.0$.
- $\mathbf{C} = \{gender, credit\}$, $T = \emptyset$: male and good credit, check passes $\Rightarrow P(loan = 1) = 1.0$.
- $\mathbf{C} = \{gender, credit\}$, $T = \{check-failed\}$: male and good credit, witness holds 0 $\Rightarrow P(loan = 1) = 1.0$.

$$P_k^n(\{1\} \mid u_2) = \frac{1}{6}(0.05 + 1.0 + 1.0 + 1.0) \approx 0.508.$$

In u_3 , male Alice is also not checked ($check = 0$ regardless of gender), so all necessity worlds yield denial: $P_k^n(\{1\} \mid u_3) = 0$.

Combining, with $ci(y^s, y^n, y^*) = \mathbb{I}\{y^n \neq y^*\} \mathbb{I}\{y^s = y^*\}$ and $y^* = loan = 0$:

$$\begin{aligned} \text{PCI}_{gender}(\text{Alice}) &= \mathbb{E}[ci(Y^s, Y^n, y^*)] = \mathbb{P}(Y^n = 1, Y^s = 0) \\ &= \sum_u p(u) P_k^s(\{0\} \mid u) P_k^n(\{1\} \mid u) \\ &= 0.2 \times \frac{2}{3} \times 0.192 + 0.7 \times \frac{2}{3} \times 0.508 + 0.1 \times 0 \\ &\approx 0.026 + 0.237 = 0.263. \end{aligned}$$

Table 2 gives the full results for Alice and Bob under these parameter choices: Pearl’s PN/PS/PNS, PCI without witnesses ($\mathbf{W} = \emptyset$), and PCI with the *check-failed* witness active, with the latter further decomposed into its necessity and sufficiency marginals.

Person, Variable	Pearl			PCI ($\mathbf{W} = \emptyset$)	PCI ($\mathbf{W} = \{\text{check-failed}\}$)		
	PN	PS	PNS		Necessity	Sufficiency	Total
Alice (<i>gender</i>)	0.045	1.000	0.045	0.210	0.394	0.667	0.263
Alice (<i>credit</i>)	0.180	1.000	0.180	0.240	0.298	0.667	0.199
Bob (<i>gender</i>)	0.000	—	0.000	0.038	0.030	0.637	0.019
Bob (<i>credit</i>)	0.900	0.955	0.860	0.228	0.188	0.637	0.119

Table 2: Per-individual responsibility scores for *gender* and *credit* on $loan = 0$ in the stochastic OCB model. Pearl’s PN/PS/PNS, PCI without witnesses ($\mathbf{W} = \emptyset$), and PCI with the *check-failed* witness shown side by side; the rightmost three columns decompose $\text{PCI}(\mathbf{W} = \{\text{check-failed}\})$ into its necessity and sufficiency marginals (their joint expectation is the bold total). Here the total happens to equal the product of the two marginals because the sufficiency measure $P_k^s(\cdot \mid \mathbf{u})$ is constant in $\mathbf{u}$ for both individuals; in general the joint $P_k^{s,n}$ does not factor into the product of its marginals. Bob’s PS for *gender* is undefined because the conditioning event (female, bad, $loan = 1$) has probability zero in the OCB model; the PCI marginals are well-defined throughout, since they integrate over P_U rather than conditioning on the factual outcome.

For Alice, PNS and PCI without witnesses both fail **D-A-rank** (p. 12): they rank *credit* above *gender* (0.180 vs. 0.045; 0.240 vs. 0.210). PCI with witnesses reverses this (0.263 vs. 0.199), satisfying **D-A-rank** as well as **D-A1** and **D-A2** (both values positive). Holding *check-failed* at its factual value blocks the credit-check bottleneck in noise realizations where Alice was checked, isolating *gender*’s structural role. The *check-failed* witness is the decisive ingredient.

For Bob, **D-B2** and **D-B-rank** hold under all three methods: *credit* is ranked above *gender* throughout. **D-B1** requires strictly positive attribution for *gender*: PNS returns exactly 0.000 and therefore fails, because $P(loan = 1 \mid gender = \text{female}, credit = \text{bad}) = 0$ makes the counterfactual necessity calculation zero, missing *gender*’s indirect role in opening the credit evaluation. PCI with witnesses returns 0.019, correctly capturing this indirect contribution. The cross-individual desideratum **D-comp** holds under both PCI (0.263 > 0.019) and PNS (0.045 > 0.000). A full verification of all seven desiderata is given at the end of this section.

The PNS and PCI (no witnesses) columns differ even though neither uses witnesses, because PCI still Γ -averages over candidate suspect sets. With $|\mathbf{S}| = 2$ PCI averages over $\{gender\}$, $\{credit\}$, and $\{gender, credit\}$ (each weighted $\frac{1}{3}$), whereas PNS evaluates a single counterfactual on one variable at a time. Property (i) below establishes that the two agree exactly when $|\mathbf{S}| = 1$ (single suspect, no witnesses, binary outcome), whereas the OCB setup uses $|\mathbf{S}| = 2$ deliberately, to demonstrate joint-subset attribution.

3.1 Continuous Outcomes and the General Causal Impact Function

The causal impact function ci is a user-defined component of PCI, and different choices encode different causal questions. In the benchmarking experiments of Section 9, we instantiate ci using only the necessity component, since Halpern’s actual causality has no sufficiency counterpart: necessity-only attribution allows a direct parallel comparison. The full framework incorporating both Y^s and Y^n is instantiated in the PNS binary example above and in the Absolute Difference example below; the OCB analysis of this section (Tables 2–3) provides a concrete validation, since the PNS binary ci jointly conditions on $Y^s = y^*$ and $Y^n \neq y^*$ and it is precisely this combination that produces

the correct **D-A-rank** ordering. Further experiments using the full ci on continuous outcomes are presented in the signal-with-mediation example (Section 4.3), the synthetic evaluation (Section 5), and the SIR benchmark (Section 10).

When the outcome variable Y is continuous rather than binary, indicators are replaced by a distance-based score that measures how far the necessity and sufficiency worlds land from the factual value y^* . The natural choice, paralleling L_1 -based scores such as ATE and SHAP, is the absolute difference.

Example 21 (Absolute Difference Impact Score). *We can develop an analogous measure for continuous outcomes where we increase attribution with distance from the factual value y^* in the necessity world and decrease impact with distance to the factual value in the sufficiency world.*

$$ci(y^s, y^n, y^*) = |y^n - y^*| - |y^s - y^*|.$$

With the full machinery in place — structural model, variable selection, alternative value distribution, joint necessity-sufficiency measure, and causal impact function — PCI delivers the following properties.

- (i) **Generalisation of PN/PS/PNS.** When $\mathbf{S} = \{X_k\}$, $\mathbf{W} = \emptyset$, and $\text{dom}(X_k) = \{0, 1\}$, the PNS binary ci recovers Pearl’s probability of necessity and sufficiency exactly; PN and PS are recovered by the corresponding one-sided ci choices.
- (ii) **Connection to actual causality.** For suitable Γ and Δ , PCI recovers Halpern’s actual causality judgements (Section 7).
- (iii) **Necessity–sufficiency decomposition.** P_k^s and P_k^n are defined and interpretable independently; practitioners may choose a ci that uses either alone or both, depending on whether the causal question concerns necessity, sufficiency, or both.
- (iv) **Witness-mediated symmetry breaking.** Holding intermediate variables fixed via $\mathbf{W}$ disambiguates overdetermination and undercutting scenarios where necessity-only methods assign equal or zero attribution, as demonstrated above for Alice and Bob, and further developed in Section 4 (SHAP and Causal SHAP comparison), Section 5 (synthetic overdetermination and undercutting archetypes), and Section 9 (witness ablation on the scaled throwing problem).
- (v) **Composability.** Γ , Δ , and ci are probability distributions and a measurable function respectively; they slot naturally into Bayesian pipelines, accept posteriors as inputs, and admit Monte Carlo estimation without modification to the core formalism.

3.2 Back to the Running Example: Verifying the Desiderata

Table 3 verifies the seven desiderata introduced at the start of this section (p. 12) against the values in Table 2. PCI with witnesses satisfies all seven; PCI without witnesses fails **D-A-rank**; PNS fails **D-A-rank** and **D-B1**. (The table uses the shorthand $|R(V \mid \text{person})|$ for $|R(V \rightsquigarrow \text{loan} \mid \text{person})|$.)

Desideratum	PNS	PCI (no wit.)	PCI (with wit.)
D-A1: $ R(\text{gender} \mid \text{Alice}) > 0$	✓ 0.045	✓ 0.210	✓ 0.263
D-A2: $ R(\text{credit} \mid \text{Alice}) > 0$	✓ 0.180	✓ 0.240	✓ 0.199
D-A-rank: $ R(\text{gender} \mid \text{Alice}) > R(\text{credit} \mid \text{Alice}) $	× 0.045 < 0.180	× 0.210 < 0.240	✓ 0.263 > 0.199
D-B1: $ R(\text{gender} \mid \text{Bob}) > 0$	× 0.000	✓ 0.038	✓ 0.019
D-B2: $ R(\text{credit} \mid \text{Bob}) > 0$	✓ 0.860	✓ 0.228	✓ 0.119
D-B-rank: $ R(\text{credit} \mid \text{Bob}) > R(\text{gender} \mid \text{Bob}) $	✓ 0.860 > 0.000	✓ 0.228 > 0.038	✓ 0.119 > 0.019
D-comp: $ R(\text{gender} \mid \text{Alice}) > R(\text{gender} \mid \text{Bob}) $	✓ 0.045 > 0.000	✓ 0.210 > 0.038	✓ 0.263 > 0.019

Table 3: Verification of the seven desiderata (p. 12) against the values in Table 2. PCI with witnesses satisfies all seven; PCI without witnesses fails **D-A-rank**; PNS fails **D-A-rank** and **D-B1**.

*PNS fails **D-A-rank** and **D-B1**. The D-A-rank failure reflects that without a check-failed witness, necessity-based attribution cannot distinguish Alice’s gender (which blocked the evaluation entirely) from her credit (which would only have mattered had she been evaluated). The D-B1 failure reflects*

that PNS assigns Bob’s gender exactly zero: because $P(\text{loan} = 1 \mid \text{female}, \text{bad}) = 0$, the counterfactual necessity calculation returns zero, missing gender’s indirect role in opening the credit evaluation. The check-failed witness resolves both.

To close this section, it is worth comparing PCI’s decomposition with the one already present in Pearl’s framework. Pearl’s framework defines three related but distinct binary scores: the probability of necessity $\text{PN} = P(Y_{x'} = y' \mid X = x^*, Y = y^*)$, the probability of sufficiency $\text{PS} = P(Y_{x^*} = y^* \mid X = x', Y = y')$, and their joint $\text{PNS} = P(Y_{x^*} = y^*, Y_{x'} = y')$. The first two are conditional probabilities; PNS is unconditional, and the three satisfy $\text{PNS} = \text{PN} \cdot P(Y_{x^*} = y^*) = \text{PS} \cdot P(Y_{x'} = y')$ rather than $\text{PNS} = \text{PN} \cdot \text{PS}$ (which would require $Y_{x^*} \perp Y_{x'}$, an assumption typically violated since the two potential outcomes share exogenous noise). PCI admits a related split: with $ci = \mathbb{I}\{y^n \neq y^*\} \mathbb{I}\{y^s = y^*\}$,

$$\mathbb{E}[ci(Y^s, Y^n, y^*)] = \int P_k^s(\{y^*\} \mid \mathbf{u}) P_k^n(\{1-y^*\} \mid \mathbf{u}) dP_{\mathbf{U}},$$

giving a *sufficiency marginal* and a *necessity marginal* that record how often, on average, restoring the factual values of a suspect set containing X_k reproduces the factual outcome, and how often substituting alternatives overturns it. The two marginals are independent given $\mathbf{u}$.

The components are not identical to Pearl’s: PN and PS condition on the factual outcome and consider a single counterfactual flip; PCI’s marginals integrate over $P_{\mathbf{U}}$, average over suspect sets weighted by Γ , and hold the witness configurations fixed when active. Table 2 (presented earlier) places both decompositions side by side for *gender* and *credit*.

For Alice’s gender, $\text{PN} = 0.045$ is the bare probability that a male with bad credit would have been approved; PCI’s necessity marginal is 0.394, nearly an order of magnitude higher, because the *check-failed* witness keeps the credit-check bottleneck active when the male counterfactual would otherwise have been checked and might still have failed, structural information that the unconditional PN cannot register. For Bob’s gender, PN is exactly 0 (a female with bad credit cannot be approved in this model) and PS is undefined; PCI’s marginals remain well-defined at 0.030 and 0.637, recovering the indirect role of gender in opening the evaluation.

The next section turns from comparing PCI with Pearl’s family of counterfactual scores to comparing it with the SHAP family, methods grounded in cooperative game theory rather than in structural counterfactual reasoning. Section 4 re-runs the OBCB analysis under plain and Causal SHAP, then introduces a continuous mediation example where the two SHAP variants diverge from each other and from PCI, with explicit desiderata recorded throughout.

4 Examples and Comparison to SHAP and Causal SHAP

Where we are. Section 3 defined PCI and verified, on Alice and Bob, that the resulting attribution satisfies all seven desiderata of that section, where PN/PS/PNS fail at least two. The natural next question is how PCI compares to the dominant practical alternative for feature attribution in the ML literature, SHAP and Causal SHAP, since these are the tools practitioners typically reach for. This section runs that comparison on two examples, recording where each method passes or fails an explicit desideratum. We first define SHAP and Causal SHAP formally (Section 4.1), then re-run OBCB (Section 4.2), where the two SHAP variants coincide because the feature set is exogenous, so the divergence between PCI and SHAP is the only contrast to draw. We then introduce a continuous mediation example (Section 4.3) where plain and Causal SHAP do diverge (one desideratum, D-YX, is fixed by the interventional correction; others, D-YM, D-MX, D-MXY, are not), and tabulate the desiderata in Section 4.4. Section 4.5 traces every failure to one of three structural gaps in the SHAP framework that PCI’s witness mechanism, suspect-set distribution Γ_s , and realised-outcome reference jointly close.

Shapley values have become one of the most widely used tools for explaining the predictions of complex machine learning models. Grounded in cooperative game theory, they provide a principled decomposition of a model’s prediction into additive contributions from individual features. Janzing et al. [2020] argued that the conditional expectation in plain SHAP is the conceptually wrong way to model feature removal: a dropped feature should be marginalised under a *do*-intervention, not under conditioning. Heskes et al. [2020] propose Causal SHAP, which implements this prescription by replacing the observational marginal used for features outside the active coalition with an

interventional distribution: when a dropped feature is causally downstream of a coalition member, its distribution under the intervention differs from its marginal, and Causal SHAP corrects for this using Pearl’s do-calculus.¹⁴ Formal definitions of both are given in Section 4.1.

Despite this repair, both methods share deeper limitations. First, both plain and Causal SHAP can assign non-zero responsibility to variables that play no causal role with respect to the target: attribution can run against the direction of the causal graph, a failure the interventional correction does not address. Second, both methods explain a model’s predicted output averaged over the population, not the individual factual outcome of a specific person: when the goal is *causal responsibility attribution*, this produces further failures that are not always made explicit in the literature. The first limitation will surface in our signal example as violations of D-YX, D-YM, and D-MX; the second will surface in OBCB as the D-A-rank overdetermination effect, and in the signal example as Instance 2’s budget collapse when the prediction equals its baseline.

We trace these failures through two examples, comparing plain SHAP, Causal SHAP, and PCI against the desiderata of Section 3 and some further desiderata related to the example we will employ. We first return to the OBCB model: it permits exact analytic computation and the desiderata are already known. In OBCB the feature set contains only exogenous roots, so plain and Causal SHAP coincide: the comparison is short. We then turn to a continuous mediation example where downstream variables are observable model inputs; there the two methods diverge, and we examine what Causal SHAP fixes and what it does not.

4.1 SHAP and Causal SHAP

In this section $\mathcal{S} \subseteq N$ denotes a SHAP coalition (a subset of features held to their factual values during evaluation), playing the same role here that $\mathbf{C} \subseteq \mathbf{S}$ played in Section 3. We use the calligraphic $\mathcal{S}$ to keep the SHAP coalition typographically distinct from PCI’s suspect set $\mathbf{S}$ on the printed page; the SHAP literature’s plain S is the same object.

Definition 22 (SHAP). A cooperative game is a pair (N, v) , where $N = \{1, \dots, n\}$ is the set of players and $v : 2^N \rightarrow \mathbb{R}$ is the characteristic function mapping each subset (coalition) $\mathcal{S} \subseteq N$ to the value that coalition can produce. The Shapley value [Shapley, 1953] assigns each player i a payoff ϕ_i , its share of the total value $v(N) - v(\emptyset)$, and is the unique attribution satisfying four axioms:

1. **Efficiency:** $\sum_i \phi_i = v(N) - v(\emptyset)$
2. **Symmetry:** If $v(\mathcal{S} \cup \{i\}) = v(\mathcal{S} \cup \{j\})$ for all $\mathcal{S}$, then $\phi_i = \phi_j$
3. **Dummy:** If $v(\mathcal{S} \cup \{i\}) = v(\mathcal{S})$ for all $\mathcal{S}$, then $\phi_i = 0$
4. **Linearity:** For two games v_1, v_2 : $\phi_i(\alpha_1 v_1 + \alpha_2 v_2) = \alpha_1 \phi_i(v_1) + \alpha_2 \phi_i(v_2)$

The unique solution is:

$$\phi_i = \sum_{\mathcal{S} \subseteq N \setminus \{i\}} \frac{|\mathcal{S}|!(|N| - |\mathcal{S}| - 1)!}{|N|!} [v(\mathcal{S} \cup \{i\}) - v(\mathcal{S})] \quad (5)$$

The weight $\frac{|\mathcal{S}|!(|N| - |\mathcal{S}| - 1)!}{|N|!}$ equals the probability that, in a uniformly random ordering of all players, exactly the members of $\mathcal{S}$ precede i . Equivalently, ϕ_i is the average marginal contribution of i across all orderings. Lundberg and Lee [2017] apply Shapley values to explain model predictions by taking players to be input features, and the characteristic function to be the expected model output given a subset of features:

$$v(\mathcal{S}) = \mathbb{E}_{X_{\mathcal{S}}} [f(X_{\mathcal{S}} = x_{\mathcal{S}}, X_{\bar{\mathcal{S}}})] \quad (6)$$

¹⁴A different counterfactual line is taken by Sharma et al. [2022], whose CF-Shapley values use a structural causal model to attribute the change in a metric from a single fixed reference x^{ref} to the observed x : $\phi_j \propto \sum_{\mathcal{S} \subseteq N \setminus \{j\}} [f(x_{\mathcal{S}}, x_j^{\text{ref}}) - f(x_{\mathcal{S}}, x_j)]$. This is the closest neighbour to PCI in spirit (single contrastive scenario, do-style intervention) and differs in three ways. (i) CF-Shapley commits to one reference x^{ref} , while PCI integrates over an alternative-value distribution $\Delta(\mathbf{x})$ and need not pick a single contrast. (ii) CF-Shapley returns a one-axis contrast $f(\cdot, x_j^{\text{ref}}) - f(\cdot, x_j)$, while PCI reports a joint (Y^s, Y^n) measure and applies a user-chosen ci , decomposing necessity and sufficiency separately. (iii) CF-Shapley has no witness mechanism, so it cannot break the overdetermination and undercutting symmetries that PCI’s witness set $\mathbf{W}$ resolves on the OBCB and signal walkthroughs of this section.

where $\bar{S} = N \setminus S$ and missing features are marginalized over their marginal distribution.

Definition 23 (Causal SHAP). *Causal SHAP replaces the characteristic function with an interventional expectation [Heskes et al., 2020]:*

$$v_{\text{causal}}(S) = \mathbb{E}[f(X) \mid \text{do}(X_S = x_S)] = \int P(X_{\bar{S}} \mid \text{do}(X_S = x_S)) f(X_{\bar{S}}, x_S) dX_{\bar{S}} \quad (7)$$

By the rules of do-calculus, intervening on X_S only affects the distribution of descendants of X_S . For non-descendants, $P(X_j \mid \text{do}(X_S = x_S)) = P(X_j)$. Writing $X_{\bar{S}, \text{nd}}$ for the features in $\bar{S}$ that are not descendants of any member of S , and $X_{\bar{S}, \text{d}}$ for those that are, equation (7) is equivalent to:

$$v_{\text{causal}}(S) = \mathbb{E}_{X_{\bar{S}, \text{nd}} \sim P(\cdot), X_{\bar{S}, \text{d}} \sim P(\cdot \mid \text{do}(X_S = x_S))} [f(X_{\bar{S}}, x_S)] \quad (8)$$

The Shapley formula (5) is otherwise unchanged. The method retains all four Shapley axioms.

At the level of subset weighting alone, PCI (with $J = 1$, $K = |S|$) and SHAP are not distinguishable: Γ 's subset weights match the Shapley kernel. The distinction lies in the integrand. SHAP averages marginal contributions using the observational conditional $\mathbb{E}[f(\mathbf{X}_S = \mathbf{x}_S, \mathbf{X}_{\bar{S}})]$, marginalising over the empirical distribution of non-coalition features; a feature merely correlated with a true cause can therefore receive positive attribution. PCI evaluates counterfactual outcomes $P(Y \in A \mid \mathbf{u}, \text{do}(\mathbf{C} = \mathbf{c}', \mathbf{T} = \mathbf{t}^*))$ computed via the structural model, so attribution flows only along genuine causal paths. The witness mechanism adds a further distinction: by holding intermediate variables fixed, PCI can resolve overdetermination and undercutting scenarios where SHAP assigns equal or incorrect attribution. The remainder of this section makes both points concrete on two examples.

4.2 OBCB: SHAP and Causal SHAP

To apply the SHAP and Causal SHAP definitions of Section 4.1 to OBCB we must specify both a feature set N and a function f on the joint feature space; the value functions and Shapley sums are derived from these. The natural choice for f is the conditional expectation of the outcome, $f(x_N) = \mathbb{E}[\text{loan} \mid X_N = x_N]$. But the OBCB PSCM has five variables — *gender*, *credit*, *check*, *check-failed*, *loan-if-checked* — so the choice of N is non-trivial. Two options are natural:

- **Option A: two upstream roots.** $N = \{\text{gender}, \text{credit}\}$. Both features are exogenous; the three internal variables (*check*, *check-failed*, *loan-if-checked*) are integrated out inside f . This is the smallest sufficient feature set and matches the way the model is normally described: predict approval from gender and credit.
- **Option B: two roots plus the mediator.** $N' = \{\text{gender}, \text{credit}, \text{check-failed}\}$. The mediator becomes an observable model input, the same status it has in the signal-with-mediation example (§4.3). This is the comparison PCI's witness mechanism naturally invites: PCI accesses *check-failed* as a witness, so allowing SHAP to access it as a feature is the fair foil.

We work through both. Option A is simpler and exposes the qualitative failures of plain SHAP. Option B then asks whether admitting the mediator rescues Causal SHAP.

Option A: two upstream roots. Here $f(g, c) = P(\text{loan} = 1 \mid g, c)$ takes values $f(\text{F}, \text{bad}) = 0.000$, $f(\text{F}, \text{good}) = 0.180$, $f(\text{M}, \text{bad}) = 0.045$, $f(\text{M}, \text{good}) = 0.900$, with the three internal variables integrated out. We apply SHAP to the rejection probability $g(x) = 1 - f(x)$ for Alice and Bob from Example 6, attributing responsibility for the denial outcome.¹⁵ By the SHAP definition (and equivalently by Definition 23 once we note that no dropped feature is a descendant of any coalition member), the value function on the 2-feature game and its closed-form specialisation under independent uniform marginals are derived in Appendix A (§A.3). Plugging in the four values of f listed at the start of Option A yields Table 4; pushing those through the closed-form Shapley gives the magnitudes in Table 5, which compares against PNS and PCI.

¹⁵Because $g = 1 - f$, the characteristic functions satisfy $v_g(S) = 1 - v_f(S)$ and $\phi_i^g = -\phi_i^f$; we compute ϕ^f and flip signs. With $|N| = 2$ and uniform marginals $P(\text{gender}) = P(\text{credit}) = \frac{1}{2}$, equation (5) reduces to $\phi_i = \frac{1}{2}[v(\{i\}) - v(\emptyset)] + \frac{1}{2}[v(N) - v(N \setminus \{i\})]$.

$\mathcal{S}$	$v(\mathcal{S})$ for Alice	$v(\mathcal{S})$ for Bob
$\emptyset$	0.281	0.281
$\{gender\}$	0.090	0.473
$\{credit\}$	0.023	0.023
$\{gender, credit\}$	0.000	0.045

Table 4: Coalition values $v(\mathcal{S}) = \mathbb{E}[f(X) \mid X_{\mathcal{S}} = x_{\mathcal{S}}^*]$ for the 2-feature game ($N = \{gender, credit\}$). $v(\emptyset) = \mathbb{E}[f(X)] = 0.281$ is the population baseline; $v(N)$ is the factual prediction $f(x^*)$. Plain SHAP and Causal SHAP coincide on this game (neither feature is downstream of the other), so a single column suffices per individual.

Person	Feature	PNS	SHAP	ATE	CATE	ITE	PCI (no wit.)	PCI (with wit.)
Alice	<i>gender</i>	0.045	0.107	0.383	0.045	0.045	0.210	0.263
Alice	<i>credit</i>	0.180	0.174	0.518	0.180	0.180	0.240	0.199
Bob	<i>gender</i>	0.000	-0.107	0.383	0.045	0.000	0.038	0.019
Bob	<i>credit</i>	0.860	0.343	0.518	0.855	0.895	0.228	0.119

Table 5: Responsibility attributions for Alice and Bob. SHAP values are signed (ϕ_i^g for $P(loan = 0 \mid x)$); desiderata compare magnitudes $|\cdot|$. ATE is a population quantity (identical across the two persons by feature); CATE conditions on the *other* covariate (credit for the gender rows; the person’s gender for the credit rows); ITE abducts the individual’s exogenous noise from the full factual record (covariates *and* the observed denial) and takes the counterfactual contrast on that pinned noise. PNS and PCI values from Table 2; ATE/CATE/ITE computed by direct PSCM evaluation in `docs/source/obcb_computations.ipynb`.

ATE and CATE fail the same two individual-level desiderata that PNS fails, and for related reasons. ATE attributes responsibility at the population level only, so its values are identical across Alice and Bob, immediately violating **D-comp** ($|R(gender \mid Alice)| = |R(gender \mid Bob)| = 0.383$). CATE conditions on observable covariates but inherits PNS’s blind spot via the *check-failed* propagation discussed in Section 2: for Alice it returns 0.045 for gender and 0.180 for credit, the same ranking as PNS_c and the same **D-A-rank** failure. CATE also fails **D-comp**, since $CATE(gender \mid credit=bad) = 0.045$ is the same number for Alice and Bob.

The individual treatment effect (ITE) is the natural endpoint of this ladder: where ATE marginalises all exogenous noise and CATE conditions only on observed covariates, the ITE abducts the unit’s noise from its *full* factual record — covariates *and* the observed denial — and then takes the counterfactual contrast holding that abducted noise fixed, the abduction–action–prediction reading already used for *PN* in Section 2. This buys back exactly the information CATE discards. The ITE separates the two applicants on **D-comp** where CATE cannot: $ITE(gender \mid Alice) = 0.045$ but $ITE(gender \mid Bob) = 0$, because abduction places Bob in the checked-and-failed branch, where being male is not what blocked him — flipping him to female (still bad credit) leaves him denied with certainty. The ITE is also a genuine counterfactual rather than an intervention: because it conditions on the outcome, $ITE(credit \mid Bob) = 0.895$ rather than the interventional $P(loan=1 \mid do(credit=good), gender=male) = 0.90$, as conditioning on Bob’s denial shifts posterior mass toward the unchecked branch. But the ITE does *not* fix **D-A-rank**: for Alice it still returns 0.180 for credit against 0.045 for gender, the same inversion as PNS_c and CATE. Abduction pins the *noise* but not the *mechanism* — the counterfactual still routes through a hypothetical credit check in the 20% of Alice’s posterior in which she was checked, so credit retains spurious responsibility. Recovering the correct ranking requires PCI’s witness mechanism, which pins *check-failed* at its factual value (dominantly the unchecked branch for Alice, where credit is causally disconnected from the loan) rather than merely integrating over its noise. In this sense PCI is the ITE *plus* the structural machinery needed to see the mediated path: it inherits the ITE’s individual-level, counterfactual character and adds the context-fixing the ITE alone lacks.

SHAP fails two desiderata. The primary failure is **D-A-rank**: Alice’s credit outranks her gender ($0.174 > 0.107$). This is an overdetermination effect: *credit=bad* correlates with the *check-failed* mechanism that blocked Alice’s credit evaluation, so SHAP’s observational marginalization inflates

credit attribution above gender’s. PCI includes the factual *check-failed* value directly, isolating gender’s upstream causal role and recovering the correct ranking. PNS fails **D-A-rank** for the same reason; it also fails **D-B1** by returning exactly zero for Bob’s gender (see Table 3).

The second SHAP failure is **D-comp**: $|\phi_{gender}^g| = 0.107$ for both Alice and Bob. But gender played a strictly larger causal role in Alice’s rejection than in Bob’s. For Alice, *gender*=female was sufficient on its own to produce rejection: it blocked the credit evaluation entirely, so her credit was never a factor. For Bob, *gender*=male was a facilitating condition: it triggered the credit check, but *credit*=bad was the actual driver of the denial. SHAP cannot distinguish these because it measures how far each feature value deviates from the population prediction mean: female and male are equidistant deviations in opposite directions, so they receive equal magnitude regardless of their individual causal weight. This is an instance of a deeper limitation: SHAP explains predicted outputs, not individual factual outcomes, which we examine further in Section 4.3.

Causal SHAP coincides exactly with plain SHAP under Option A: both features are exogenous roots, so $P(\text{credit} \mid \text{do}(\text{gender} = g)) = P(\text{credit})$ and vice versa, no dropped feature is a descendant of any coalition member, and $v_{\text{causal}}(\mathcal{S}) = v_{\text{plain}}(\mathcal{S})$ for every coalition. Option A therefore reduces to a single SHAP analysis, and the failures above belong to both methods.

Option B: two roots plus the mediator. Throughout this subsection we abbreviate *gender*, *credit*, and *check-failed* as *g*, *c*, and *cf* in mathematical expressions and table cells; running prose retains the full names. Now we admit *check-failed* as a third model input and apply both methods to the 3-feature game $N' = \{\text{gender}, \text{credit}, \text{check-failed}\}$. This requires defining f on $\{0, 1\}^3$ rather than on $\{0, 1\}^2$. The natural observational extension is again the conditional expectation $\tilde{f}(g, c, cf) = \mathbb{E}[\text{loan} \mid \text{gender} = g, \text{credit} = c, \text{check-failed} = cf]$. This is well defined for three of the four (c, cf) combinations, but not the fourth: a failed check requires that a check happened *and* the credit was bad, so *check-failed* = 1 together with *credit* = good is impossible under the PSCM and has no observational data to condition on. We leave $\tilde{f}$ undefined on this cell.

Both methods need a value function $v(\mathcal{S})$: the expected value of $\tilde{f}$ when the features in $\mathcal{S}$ are held at their factual values $x_{\mathcal{S}}^*$ and the remaining features are averaged out. The methods differ in how that average is taken. Plain SHAP averages over each dropped feature using its own observational marginal $P(X_i)$, with the dropped features treated as independent of one another. Causal SHAP averages over the joint distribution the PSCM produces under the intervention $\text{do}(X_{\mathcal{S}} = x_{\mathcal{S}}^*)$, so the dropped features keep whatever joint dependencies the PSCM imposes on them. Whether either average ever lands on the impossible cell ($c=1, cf=1$) (and so whether $v(\mathcal{S})$ is finite) depends on both the method and on which features are held fixed at which factual values. Table 6 works out all eight coalition values for both individuals.

$\mathcal{S}$	Alice (g, c, cf)=(0, 0, 0)		Bob (g, c, cf)=(1, 0, 1)	
	v_{plain}	v_{causal}	v_{plain}	v_{causal}
$\emptyset$	NaN	0.281	NaN	0.281
$\{g\}$	NaN	0.090	NaN	0.473
$\{c\}$	0.007	0.023	0.007	0.023
$\{cf\}$	0.270	0.270	NaN	NaN
$\{g, c\}$	0.000	0.000	0.014	0.045
$\{g, cf\}$	0.090	0.090	NaN	NaN
$\{c, cf\}$	0.000	0.000	0.025	0.025
N'	0.000	0.000	0.050	0.050

Table 6: Coalition values for the 3-feature game $N' = \{\text{gender}, \text{credit}, \text{check-failed}\}$. Bold **NaN** entries flag coalitions whose value function evaluates $\tilde{f}$ at the impossible cell ($c=1, cf=1$) with positive weight. Plain SHAP and Causal SHAP coincide whenever neither *credit* nor *check-failed* is dropped, since the only structural coupling in the model is between these two; they diverge whenever one is in $\mathcal{S}$ and the other is not.

Plain SHAP. Plain SHAP treats *credit* and *check-failed* as independent, so whenever both are dropped from the coalition the average runs over all four (c, cf) combinations, with the impossible ($c=1, cf=1$) combination receiving weight $P(c=1) \cdot P(cf=1) = 0.5 \cdot 0.275 \approx 0.140$, the same nonzero weight as any other corner. For Alice, both *credit* and *check-failed* are dropped at $\mathcal{S} \in$

$\{\emptyset, \{gender\}\}$, the first two NaN entries in Table 6. For Bob the same two coalitions fail, and two more break for an additional reason: at $S = \{check-failed\}$ and $S = \{gender, check-failed\}$ the value $cf^* = 1$ is held fixed while $credit$ is sampled from its marginal, so the $c=1$ branch enters with weight 0.5 and again hits the impossible cell. Plain SHAP gives undefined Shapley values for both individuals.¹⁶

Causal SHAP for Alice. Causal SHAP samples cf following its structural equation $cf = check \cdot (1 - c)$, with $check \sim \text{Bern}(p_{\text{check}}[g])$, given whatever (g, c) values appear in the integrand. When $c = 1$ the equation forces $cf = 0$ regardless of $check$, so the $(c=1, cf=1)$ combination is never produced. Every Alice entry in the v_{causal} column of Table 6 is therefore finite. Plugging Alice’s column into the Shapley formula gives magnitudes $|\phi_{gender}| = 0.097$, $|\phi_{credit}| = 0.176$, $|\phi_{check-failed}| = 0.007$, so $|\phi_{credit}| > |\phi_{gender}|$ and **D-A-rank** fails: credit outranks gender, exactly the same pathology plain SHAP exhibited on the 2-feature game.

Causal SHAP for Bob. Two coalitions break for Bob: $S = \{check-failed\}$ and $S = \{gender, check-failed\}$. Both hold $check-failed$ at Bob’s factual value $cf^* = 1$ and average $credit$ over its marginal. To see why this is fatal, apply Definition 23 to the first:

$$v_{\text{causal}}(\{cf\}) = \mathbb{E}[\tilde{f}(X) \mid do(cf = 1)] = \frac{1}{4} \sum_{g \in \{0,1\}} \sum_{c \in \{0,1\}} \tilde{f}(g, c, 1).$$

The uniform $\frac{1}{4}$ weights come from $gender$ and $credit$ sitting upstream of $check-failed$ in the causal graph: intervening on $check-failed$ overrides the structural equation that would normally determine it from $credit$ and $check$, but does not alter the marginal distributions of the upstream features. Two of the four summands evaluate $\tilde{f}$ at $(g, c=1, cf=1)$ — the impossible cell — with weight $\frac{1}{4}$ each. So $v_{\text{causal}}(\{cf\})$ is undefined, and so is $v_{\text{causal}}(\{g, cf\})$ for the same reason. These two coalitions appear in the Shapley sum for every feature, so all three of Bob’s Causal SHAP values are NaN (Bob’s columns in Table 6).

Why Bob breaks and Alice does not. The same two coalitions arise for Alice; the difference is the value of cf^* being held fixed. With Alice’s $cf^* = 0$, the $c=1$ summand in the formula above evaluates $\tilde{f}(g, 1, 0) = p_{\text{check}}[g] \cdot \text{loan_prob}[g, 0]$, which is defined because the PSCM does allow $(c=1, cf=0)$: when $credit$ is good, cf is forced to 0 deterministically, so the conditional expectation has a value. With Bob’s $cf^* = 1$, the $c=1$ summand evaluates $\tilde{f}(g, 1, 1)$, the impossible cell, “good credit and a failed check”, which the PSCM rules out, since failing the credit check requires bad credit by the structural equation $cf = check \cdot (1 - c)$.

The mechanism producing this asymmetry is the intervention itself. When the PSCM runs forward, $credit$ and $check-failed$ are linked by $cf = check \cdot (1 - c)$, which keeps $c=1$ and $cf=1$ from co-occurring. When Causal SHAP intervenes via $do(cf = cf^*)$, this link is overridden: $check-failed$ is fixed externally and $credit$ is sampled freely. The override always produces a $(c=1, cf^*)$ combination with positive weight; whether $\tilde{f}$ has a value for that combination depends on which cf^* value Causal SHAP is intervening on.

Promoting the mediator to a feature therefore does not rescue Causal SHAP: it fails the desideratum on Alice and breaks down outright on Bob.

The OBCB analysis exposed two structural failure modes: SHAP’s overdetermination effect on the 2-feature game, and Causal SHAP’s breakdown once the mediator is admitted as a feature under Option B. The second of these depends on a structural feature of the OBCB PSCM: the equation $cf = check \cdot (1 - c)$ ties $credit$ and $check-failed$ together, so Option B’s natural feature set contains one combination of input values (“good credit and a failed check”) that the model is not defined on. To see whether the same families of failure appear without this complication, we now turn to the signal-with-mediation chain $X \rightarrow M \rightarrow Y$, where the mediator is naturally an observable input, $\tilde{f}$ is defined everywhere, and no impossible-cell issue arises. Causal SHAP’s failures there are not artifacts of an undefined $\tilde{f}$ but properties of the method itself.

¹⁶The underlying issue is that plain SHAP does not see that $credit$ and $check-failed$ are linked by the structural equation $cf = check \cdot (1 - c)$, and so cheerfully samples a combination the PSCM rules out.

4.3 Signal with Mediation

Example 24 (Signal with mediation). We work with the following **generative model**: X sends a signal, M mediates it with some noise, and Y is the noisy final outcome:

$$X \sim \mathcal{N}(0.5, 0.25) \quad (9)$$

$$M = X + \varepsilon_M, \quad \varepsilon_M \sim \mathcal{N}(0, 0.1) \quad (10)$$

$$Y = M + \varepsilon_Y, \quad \varepsilon_Y \sim \mathcal{N}(0, 0.1) \quad (11)$$

The causal graph is $X \rightarrow M \rightarrow Y$, a simple mediation chain. All noise terms are independent of each other and of X . Since everything is jointly Gaussian, all conditional expectations are linear functions of the conditioning variables, and no approximations are required.

Let $R(X \rightsquigarrow Y)$ denote the causal responsibility of X for Y . Following the convention of Section 3, the desiderata are stated in terms of absolute magnitudes $|R(\cdot)|$, allowing for directionality. The **qualitative causal responsibility desiderata** for this example are:

$$\mathbf{D}\text{-}\mathbf{XY} \quad |R(X \rightsquigarrow Y)| > 0$$

$$\mathbf{D}\text{-}\mathbf{MY} \quad |R(M \rightsquigarrow Y)| > 0$$

$$\mathbf{D}\text{-}\mathbf{XM} \quad |R(X \rightsquigarrow M)| > 0$$

$$\mathbf{D}\text{-}\mathbf{YX} \quad |R(Y \rightsquigarrow X)| = 0$$

$$\mathbf{D}\text{-}\mathbf{YM} \quad |R(Y \rightsquigarrow M)| = 0$$

$$\mathbf{D}\text{-}\mathbf{MX} \quad |R(M \rightsquigarrow X)| = 0$$

A more subtle desideratum concerns telling apart the roles of the two genuine causes. X reaches Y only through the mediator M , whereas M acts on Y directly; a good attribution method should be able to distinguish these two roles rather than collapse them into the same number. One systematic way to register the distinction is to require that the two causes receive different, rankable scores, and we adopt the convention of assigning the larger score to the cause closer to the outcome:

$$\mathbf{D}\text{-}\mathbf{MXY} \quad |R(M \rightsquigarrow Y)| > |R(X \rightsquigarrow Y)|$$

The direction of the inequality is a convention rather than a law of responsibility; what matters is that the method separates the direct from the indirect path at all, and scoring the closer cause higher is one principled, consistent way to register that separation. We apply this convention throughout.

The key **distributional facts** we will use are:

$$\mathbb{E}[X] = \mathbb{E}[M] = \mathbb{E}[Y] = 0.5, \quad (12)$$

$$(\text{Var}(X), \text{Var}(M), \text{Var}(Y)) = (0.25, 0.35, 0.45), \quad (13)$$

$$(\text{Cov}(X, M), \text{Cov}(X, Y), \text{Cov}(M, Y)) = (0.25, 0.25, 0.35). \quad (14)$$

For each target variable, the optimal predictor under squared loss is the conditional expectation, computable exactly in this Gaussian model:

$$f_Y(X, M) = \mathbb{E}[Y \mid X, M] = M \quad (\{X, M\} \Rightarrow Y) \quad (15)$$

$$f_M(X, Y) = \mathbb{E}[M \mid X, Y] = 0.5X + 0.5Y \quad (\{X, Y\} \Rightarrow M) \quad (16)$$

$$f_X(M, Y) = \mathbb{E}[X \mid M, Y] = 0.5 + \frac{5}{7}(M - 0.5) \quad (\{M, Y\} \Rightarrow X) \quad (17)$$

The distributional facts and the three conditional expectations above are derived in Appendix B (§B.1–§B.2).

4.4 Computations and Desiderata Analysis

We analyze two instances. **Instance 1** sets $X^* = M^* = Y^* = 1$: every variable lies above its mean, the prediction is non-trivial at every target, and SHAP, Causal SHAP, and PCI all return non-zero values. **Instance 2** sets $X^* = M^* = Y^* = 0.5$: every variable sits at its mean, the prediction equals its baseline, and SHAP’s attribution budget $f(x^*) - \mathbb{E}[f(X)]$ collapses to zero. We work through Instance 1 first, then return to Instance 2 to isolate what each method is really tracking.

Computing the three methods. At Instance 1 ($X^* = M^* = Y^* = 1$) all three methods admit closed-form evaluation in this Gaussian model. We defer the full per-target coalition-value and Shapley computations for plain and Causal SHAP, and the per-(C, T) PCI derivation, to Appendix B (§B.3–§B.5); the resulting attributions populate Table 7. One computed fact drives the D-YX discussion below and is worth stating here.

Plain SHAP assigns $\phi_Y = 0.139 > 0$ for target X , even though Y does not cause X . This arises because $v(\{Y\}) = 0.778 > 0.5$: observing $Y=1$ is statistically informative about $X=1$ through the chain $X \rightarrow M \rightarrow Y$. SHAP cannot distinguish “ Y is informative about X ” from “ Y causes X ” because the value function is grounded in observational conditionals, which are symmetric under reversal of causal direction.

Desiderata analysis. Following the convention of Section 3, we use $R(A \rightsquigarrow B)$ for the method-agnostic responsibility and ϕ_A^B exclusively for SHAP/Causal SHAP values; prose statements about desiderata refer to magnitudes $|R(A \rightsquigarrow B)|$. Here A denotes the candidate cause and B the target outcome (the generic Y of Section 3); for the PCI columns B^n, B^s are its necessity/sufficiency worlds, B^* its factual value, and $\bar{c} = |B^n - B^*| - |B^s - B^*|$ the absolute-difference impact of Example 21. Table 7 maps the desiderata from Section 4.3 to the values computed above.

Desideratum	Condition	Plain SHAP	Causal SHAP	PCI ($\mathbf{W}=\emptyset$)	PCI ($\mathbf{W}=3\text{rd}$)
D-XY	$ R(X \rightsquigarrow Y) > 0$	0.250 ✓	0.250 ✓	0.249 ✓	0.186 ✓ [†]
D-MY	$ R(M \rightsquigarrow Y) > 0$	0.250 ✓	0.250 ✓	0.283 ✓	0.319 ✓
D-XM	$ R(X \rightsquigarrow M) > 0$	0.306 ✓	0.375 ✓	0.253 ✓	0.284 ✓
D-YX	$ R(Y \rightsquigarrow X) = 0$	0.139 ×	0.000 ✓	0.000 ✓	0.000 ✓
D-YM	$ R(Y \rightsquigarrow M) = 0$	0.194 ×	0.125 ×	0.000 ✓	0.000 ✓
D-MX	$ R(M \rightsquigarrow X) = 0$	0.219 ×	0.357 ×	0.000 ✓	0.000 ✓
D-MXY	$ R(M \rightsquigarrow Y) > R(X \rightsquigarrow Y) $	0.25=0.25 ×	0.25=0.25 ×	0.283>0.249 ✓	0.319>0.186 ✓ [‡]

Table 7: Desiderata satisfied (✓) and violated (×) by plain SHAP, Causal SHAP, and PCI on the signal-with-mediation example ($X \rightarrow M \rightarrow Y$), instance $X = M = Y = 1$. PCI uses $\bar{c} = \mathbb{E}[|B^n - B^*|] - \mathbb{E}[|B^s - B^*|]$ with Γ_s, Γ_w uniform over non-empty subsets and rejection on $\mathbf{C} \cap \mathbf{T} \neq \emptyset$. $\mathbf{W}=3\text{rd}$ sets the third variable (neither A nor B) as the witness when not in $\mathbf{C}$. [†] 0.186 weights three suspect-containing (C, T) configurations at 1/4 each (the rejection-sampling filter keeps 2/3 of $\Gamma_s \times \Gamma_w$ draws, including a fourth valid pair ($\{M\}, \emptyset$) that does not contain X): one pins M as witness (contribution 0, necessity and sufficiency coincide) and two let M propagate freely (contributions 0.320 and 0.425). [‡] The two values use different witness sets ($\mathbf{W}=\{M\}$ for D-XY, $\mathbf{W}=\{X\}$ for D-MY); this column reports the amplified verdict under an asymmetric witness rule. The $\mathbf{W}=\emptyset$ column above is the corresponding single-game ranking and already passes D-MXY.

Positive results (D-XY, D-MY, D-XM). Both methods correctly assign positive attribution to every direct or upstream cause of the target. Causal prediction models are built from observational conditionals, which do preserve correlation from cause to effect, so these desiderata pose no challenge to either method.

Causal SHAP fixes D-YX, not D-YM. For target X , the interventional value $v_{\text{causal}}(\{Y\}) = 0.5$ equals the baseline because M is not downstream of Y : $P(M \mid \text{do}(Y=1)) = P(M)$, so fixing $Y = 1$ provides no interventional leverage over f_X , and Causal SHAP correctly yields $\phi_Y^X = 0$. For target M , however, $\phi_Y^M = 0.125 > 0$ persists. The prediction model $f_M(X, Y) = 0.5X + 0.5Y$ contains Y as an input because Y is observationally predictive of M ; setting $Y = 1$ raises the model output even under the interventional regime. Causal SHAP replaces the conditioning distribution but cannot remove a feature that the prediction model itself uses.

D-MX worsens under Causal SHAP. Plain SHAP assigns $\phi_M^X = 0.219$; Causal SHAP assigns 0.357. Fixing D-YX reduces $v_{\text{causal}}(\{Y\})$ from 0.778 to 0.5, driving ϕ_Y^X to zero, but reallocates the freed attribution to M . Since the structural noise that actually drives X is unobserved, all attribution concentrates on the only visible predictor, M , even though M does not cause X .

D-MXY fails for both. Since $f_Y = M$, the singleton values $v(\{X\}) = \mathbb{E}[M \mid X=1] = 1$ and $v(\{M\}) = 1$ are identical, giving $\phi_M^Y = \phi_X^Y = 0.25$ for both methods. The Shapley formula cannot distinguish that X reaches Y only through M : when $X = 1$, the expected prediction is already 1 regardless of whether M is observed or not, so both features receive the same marginal contribution.

PCI: structural intervention separates the paths; witnesses sharpen the gap. Table 7 reports PCI in two configurations: without witnesses ($\mathbf{W} = \emptyset$) and with the third variable as witness.

Without witnesses, PCI satisfies all six single-variable desiderata. D-YX, D-YM, and D-MX are zero exactly: intervening on a variable that does not structurally affect the target leaves both the necessity and sufficiency worlds with identical distributions, so $ci = 0$ identically. D-XY, D-MY, and D-XM are positive. D-MXY also **passes** at $\text{PCI}(M \rightarrow Y) \approx 0.283 > \text{PCI}(X \rightarrow Y) \approx 0.249$, but only by a margin of 0.034. The asymmetry comes from the sufficiency world: at $C=\{X\}$, X 's path to Y accumulates $\varepsilon_M + \varepsilon_Y$ in $|Y^s - Y^*|$, while at $C=\{M\}$, M 's path accumulates only ε_Y . The necessity expectations match across paths ($\text{Var}(X) + \text{Var}(\varepsilon_M) + \text{Var}(\varepsilon_Y) = \text{Var}(M) + \text{Var}(\varepsilon_Y) = 0.45$), so the gap is carried entirely by the sufficiency variance asymmetry $\text{Var}(\varepsilon_M)$. A small structural margin like this is fragile: under any perturbation of the noise variances or the contrast function the $\mathbf{W} = \emptyset$ ordering could flip.

Admitting the third variable as a witness amplifies the gap by roughly fourfold. $\text{PCI}(X \rightarrow Y \mid \mathbf{W}=\{M\}) \approx 0.186$ weights three suspect-containing (C, T) configurations at 1/4 each: in two cases M propagates freely (contributing 0.320 and 0.425) and in one case M is pinned as witness in both worlds, equalizing them and contributing 0. $\text{PCI}(M \rightarrow Y \mid \mathbf{W}=\{X\}) \approx 0.319$ weights three cases in which M takes an alternative value; since X does not appear in Y 's structural equation, pinning X has no effect on Y^n or Y^s , and all three cases contribute the same 0.425. The gap widens from 0.034 at $\mathbf{W} = \emptyset$ to $0.319 - 0.186 = 0.133$ at $\mathbf{W} = \{\text{third}\}$: the ordering is now *robust* rather than marginal. The comparison relies on different witness sets for the two cells, so it remains illustrative rather than a single-game ranking.

Instance 2 (baseline). We now return to **Instance 2**: every variable sits at its mean, $X^* = M^* = Y^* = 0.5$, and the model predicts the baseline at every target: $f_Y(0.5, 0.5) = 0.5$, $f_M(0.5, 0.5) = 0.5$, $f_X(0.5, 0.5) = 0.5$. At this instance, SHAP's attribution budget $f(x^*) - \mathbb{E}[f(X)]$ vanishes, while PCI's machinery is unaffected.

desideratum	Plain / Causal SHAP		PCI ($\mathbf{W} = \emptyset$)		PCI ($\mathbf{W} = \{\text{third}\}$)	
	value	status	value	status	value	status
D-XY	0.000	×	0.154	✓	0.115	✓
D-MY	0.000	×	0.189	✓	0.212	✓
D-XM	0.000	×	0.146	✓	0.165	✓
D-YX	0.000	✓	0.000	✓	0.000	✓
D-YM	0.000	✓	0.000	✓	0.000	✓
D-MX	0.000	✓	0.000	✓	0.000	✓
D-MXY	0.000 = 0.000 ×		0.189 > 0.154 ✓		0.212 > 0.115 ✓	

Table 8: Desiderata at the baseline instance $X^* = M^* = Y^* = 0.5$. SHAP collapses to zero on every cell, correctly assigning zero to the non-causal triples by accident, but with no signal to offer on the causal ones. PCI reports the same structural verdicts as at $X^* = M^* = Y^* = 1$.

The pattern in Table 8 reflects the object each method is decomposing. SHAP attributes the quantity $f(x^*) - \mathbb{E}[f(X)]$ across features. At the baseline this gap is zero on every target, so the budget being distributed is zero, and every cell collapses to zero, uniformly, regardless of how the realized outcome was actually produced. The non-causal triples (D-YX, D-YM, D-MX) are satisfied only by this collapse: SHAP is not isolating the absence of a causal path; it is reporting that there is nothing to attribute. The same collapse turns D-XY, D-MY, and D-XM into violations.

PCI never invokes a population expectation. With $\mathbf{S}$ and $\mathbf{W}$ defined relative to a fixed factual instance, the question is whether varying the suspect would move the realized outcome y^* away from itself, not whether $f(x^*)$ deviates from $\mathbb{E}[f(X)]$. The structural mechanism is unchanged at the baseline, so the three causal pairs continue to register positive values; the three non-causal pairs continue to register zero exactly, because intervening on a variable that does not structurally affect the target leaves both worlds with identical distributions. The D-MXY verdict is preserved in both witness configurations, with margin 0.035 at $\mathbf{W} = \emptyset$ (vs. 0.034 at the $X^* = M^* = Y^* = 1$ instance) and 0.097 at $\mathbf{W} = \{\text{third}\}$ (vs. 0.133). The gap is a property of the PSCM's noise variances, not of where the factual sits.

The two instances together make the contrast sharp. At Instance 1, SHAP’s attribution budget $f(x^*) - \mathbb{E}[f(X)]$ is non-zero and SHAP returns non-trivial numbers; at Instance 2 the budget vanishes and every cell collapses. PCI returns the same structural verdicts on all seven desiderata across the two instances. The lesson is not that SHAP gives the wrong answer at the baseline: it correctly reports that the factual prediction does not deviate from its mean. The point is that this is a different question from causal responsibility for y^* : when the prediction equals its baseline, SHAP has no budget to distribute even when the PSCM’s causal mechanism is perfectly intact. PCI asks the responsibility question directly, and answers it whether or not the prediction happens to coincide with its population mean.

4.5 Summary Across Both Examples

The failures above are not artifacts of a particular feature set or parameter choice: each traces to a structural limitation of the SHAP/Causal SHAP framework. Causal SHAP strictly improves on plain SHAP for one desideratum (D-YX), by replacing observational conditioning with interventional distributions; the remaining failures share two sources: prediction models include non-causal features that the conditioning distribution cannot exclude (D-YM, D-MX), and Shapley averaging cannot distinguish direct from indirect causal paths (D-MXY). The OBCB analysis of Section 4.2 exhibits the same family in different surface form: plain SHAP fails **D-A-rank** on the 2-feature game, and Causal SHAP either inherits that failure (Alice) or becomes undefined outright (Bob) once the mediator is admitted as a feature. Across both examples, neither SHAP variant reliably satisfies the desiderata that motivated PCI in Section 3. PCI addresses these gaps through three ingredients the SHAP framework systematically lacks: the witness mechanism, the suspect-set distribution Γ_s , and the use of the realized outcome rather than the prediction baseline; the path-noise asymmetry in the sufficiency world produces a structural gap on D-MXY, which appears already in the witness-free single game and is amplified into a robust margin under a witness rule.

Where we go next. The two examples in this section are small enough to inspect every desideratum individually, but they cover only two causal patterns. Section 5 stress-tests PCI on a synthetic structural model carrying three textbook archetypes — linear necessary-and-sufficient, overdetermined-not-necessary, and preempted (gated) — plus an irrelevance control, and verifies that PCI separates them at a Monte Carlo budget small enough to bootstrap honest error bars. Section 6 then contrasts PCI with gradient-based attribution on a continuous credit-limit example.

5 Synthetic Evaluation with Overdetermination and Undercutting

Where we are. Section 4 has compared PCI with SHAP and Causal SHAP on two small examples and shown where the SHAP framework’s desiderata fail. Those examples cover only two causal patterns: overdetermination through a gating mediator (OBCB) and correlated mediation (signal). This section stress-tests PCI across a wider catalogue of patterns, with an analytic ground truth available throughout: the question is whether PCI correctly separates linear, overdetermined, and preempted contributions on a single structural model, and stays at the noise floor on a variable that plays no role.

To evaluate PCI in a setting that exhibits multiple distinct causal patterns simultaneously, we construct a synthetic structural causal model carrying three archetypes — linear necessary-and-sufficient, overdetermined-not-necessary, and preempted (gated) — together with an irrelevance control.¹⁷

The model has six independent root variables and three deterministic branches feeding the outcome E (the generic outcome Y of Section 3, written E here to match the model diagram; its factual value

¹⁷The classical motivating example is Sally and Billy throwing stones at a bottle: Sally’s stone arrives first and shatters it; Billy’s would have done the same had Sally’s not preempted it. The case is canonical in the actual-causality literature [Halpern, 2016] and motivates the preempted archetype below; we use generic letters rather than personal names to keep variable labelling tight.

is y^* and its necessity/sufficiency worlds Y^n, Y^s):

$$\begin{aligned} L_1, L_2 &\sim \mathcal{N}(0, 1), \quad O_1, O_2 \sim \mathcal{N}(1, 1), \quad P, D \sim \mathcal{N}(0, 1), \\ \text{lin} &= 5L_1 + 10L_2, \\ \text{od} &= \max(5O_1, 5O_2), \\ \text{gate} &= \mathbb{1}\{|L_2| \leq \tau\}, \\ \text{p_branch} &= 5P \cdot \text{gate}, \\ E &= \text{lin} + \text{od} + \text{p_branch}. \end{aligned}$$

We pick $\tau = 0.674$ so the gate is on / off about half the time ($\Pr(|L_2| \leq 0.674) \approx 0.5$ for $L_2 \sim \mathcal{N}(0, 1)$). The DAG is shown in Figure 1. The variable D is sampled but never enters E , serving as an irrelevance control, so any non-zero score for D is the noise floor of the procedure.




Figure 1: Synthetic archetype model. Roots (circles) are independent draws: $L_1, L_2 \sim \mathcal{N}(0, 1)$ (linear, purple); $O_1, O_2 \sim \mathcal{N}(1, 1)$ (overdetermined via max, blue); $P \sim \mathcal{N}(0, 1)$ (gated by $|L_2| \leq \tau$, green); $D \sim \mathcal{N}(0, 1)$ (irrelevance control, grey). Mediators (white ellipses) carry the deterministic functions of their parents. Outcome E (orange double-circle) is the sum $\text{lin} + \text{od} + \text{p_branch}$. The variable D has no outgoing edges by construction.

The PCI evaluation treats the six roots as the suspect set; deterministic mediators are not suspects. Restricting suspects to roots loses no causal information here since downstream values are determined by the roots (formal statement deferred to Section 3).

Causal impact, decomposed. We follow the impact function $ci(y^s, y^n, y^*) = |y^n - y^*| - |y^s - y^*|$ defined in Section 3, but report its two components separately as ci_N and ci_S :

$$\begin{aligned} ci_N(V) &= \mathbb{E}[|Y^n - y^*| \mid V \text{ intervened}] \quad (\text{necessity}), \\ ci_S(V) &= -\mathbb{E}[|Y^s - y^*| \mid V \text{ pinned at factual}] \quad (\text{sufficiency}), \end{aligned}$$

with $ci(V) = ci_N(V) + ci_S(V)$. Higher ci_N means more necessary; higher ci_S (closer to zero) means more sufficient. The raw scores carry a co-intervention baseline that does not vanish for D , so we report *excess* over D within each case:

$$\Delta_N(V) = ci_N(V) - ci_N(D), \quad \Delta_S(V) = ci_S(V) - ci_S(D).$$

D is the noise floor; any other variable's Δ is its causal signal above it.

Two factual cases. We instantiate the model with one batch of 500 events and select two factual observations to read all archetypes off.

Case 1 (preempted, contestable O pair). Filter: $|L_2| > \tau$ (gate off, so $\text{p_branch} = 0$ regardless of P) and $|O_1 - O_2| < \delta$ with $\delta = 1$ (the two overdetermined contributors are close: the winner is barely the winner; we call such a pair *contestable*, since a small perturbation to either contributor

could flip which one wins $\max(5O_1, 5O_2)$). Linear N+S on the L_s , between-variable sufficiency asymmetry on a contestable O pair, and noise floor for both P and D should all be readable from this single observation.

Case 2 (unpreempted, dominant O winner). Filter: $|L_2| \leq \tau$ (gate on, P branch active) and $|O_1 - O_2| > \delta$. The dominant winner is what makes the within-variable S -not- N gap visible: the winner is large enough that alternatives rarely beat it (high Δ_S), and the loser provides a partial floor when the winner is intervened (reduced Δ_N). A contestable pair cannot show this within-variable gap, since pinning either one still lets the partner flip the max.

The two cases together test the structural-role-not-identity claim: the same variable P should sit at the noise floor in Case 1 and mirror L_1 in Case 2.

Desiderata and measurements. Table 9 pairs each archetype expectation with the formal inequality on the excess scores Δ_N, Δ_S , the measured value at the production sample budget ($N = 20\,000$ Monte Carlo draws per intervention), and a bootstrap 95% confidence interval (4 000 resamples, joint across V and D so per-sample correlations are preserved). The noise-floor variable D itself drifts across the two cases: co-intervention variance is structurally larger in the gate-on regime than in the gate-off regime, so $\text{ci}_N(D)$ shifts between regimes by a non-negligible amount. This is exactly why every archetype claim is stated on the excess scores Δ_N, Δ_S rather than raw ci values: subtracting D within each case normalises out the regime-dependent floor, so the archetype patterns can be read off stably regardless of D 's absolute drift.

The threshold from the data, not from a guess. The directional desiderata of the form $\Delta > \epsilon$ or $|\Delta| < \epsilon$ all depend on a single tolerance ϵ . Rather than hand-pick it, we derive it from the Monte Carlo budget. For every Δ score appearing in any desideratum we bootstrap a standard error σ_V and take

$$\epsilon = 1.645 \cdot \sigma_{\max}, \quad \sigma_{\max} = \max_{V, \text{case}} \sigma_V,$$

the one-sided 95% Z-quantile times the largest standard error across all measured Δ s. Under the null hypothesis $\Delta = 0$, fewer than 5% of measurements would exceed ϵ ; a Δ that clears ϵ is therefore not Monte Carlo chatter at the chosen budget. At $N = 20\,000$ samples we measure $\sigma_{\max} \approx 0.092$, giving $\epsilon \approx 0.151$. Smaller budgets give a noisier ϵ that defeats the point (e.g. at $N = 5\,000$, $\sigma_{\max} \approx 0.18$ would push ϵ into the signal range); this fixes the production sample size for the section. Ten of the twelve desiderata listed in earlier drafts survive this honest threshold; the two that do not (cross-regime P flip, and quantitative match of $\Delta_N(P)$ to $\Delta_N(L_1)$ in Case 2) are discussed below as diagnostics.

#	Formal desideratum	Measured	95% CI	Result
1	$\Delta_N(L_2)$ is the largest excess necessity, Case 1	+2.129	[+1.953, +2.300]	✓
2	$\Delta_N(L_2)$ is the largest excess necessity, Case 2	+0.642	[+0.476, +0.813]	✓
3	$\Delta_S(O_w) - \Delta_S(O_l) > \epsilon$, Case 1 (contestable)	+0.347	[+0.216, +0.473]	✓
4	$\Delta_S(O_w) - \Delta_S(O_l) > \epsilon$, Case 2 (dominant)	+0.529	[+0.387, +0.667]	✓
5	$ \Delta_N(O_w) - \Delta_N(O_l) < \epsilon$, Case 1	-0.093	[-0.266, +0.081]	✓
6	$\Delta_S(O_w) - \Delta_N(O_w) > \epsilon$, Case 2 (within-variable S -not- N signature)	+0.487	[+0.256, +0.712]	✓
7	$\Delta_N(L_2) - \Delta_N(O_w) > \epsilon$, Case 2	+1.082	[+0.915, +1.251]	✓
8	$ \Delta_N(P) < \epsilon$, Case 1 (gate off)	+0.112	[-0.058, +0.280]	✓
9	$\Delta_N(L_1) - \Delta_N(P) > \epsilon$, Case 1	+0.670	[+0.499, +0.843]	✓
10	$\Delta_N(P) > \epsilon$, Case 2 (gate on)	+0.250	[+0.079, +0.422]	✓

Table 9: Desiderata satisfaction on the synthetic archetype model. Threshold $\epsilon = 1.645 \cdot \sigma_{\max} \approx 0.151$ is the smallest Δ gap that, at $N = 20\,000$ samples, exceeds Monte Carlo noise at one-sided 95% confidence. Subscripts w/l denote the winner / loser of the $\max(5O_1, 5O_2)$ branch in the given case. Per-row check identifiers and the underlying numerical computations are listed in the companion notebook `responsibility_archetypes.ipynb`.

Figure 2 visualises the same ten claims as a forest plot: each row carries a plain-language statement of the structural expectation; the dot is the point estimate of the relevant Δ score (or difference of Δ scores); the horizontal bar is its bootstrap 95% CI; and the green band is the row's *pass region*,

i.e. the set of Δ values that would confirm that row's claim given ϵ . The pass region depends on the claim type: $\Delta > \epsilon$ claims pass to the right of ϵ ; $|\Delta| < \epsilon$ closeness claims pass inside the strip $(-\epsilon, +\epsilon)$; rank claims (#1 and #2) pass to the right of 0, since they compare $\Delta_N(L_2)$ against the largest competing suspect. A row is counted confirmed when the point estimate lies in its green band; the CI is shown separately so the reader can see when a measurement is statistically tight and when its CI crosses the pass boundary. All ten dots fall inside their pass bands. The tightest row is #10 ($\Delta_N(P) > \epsilon$ in Case 2, gate on): the point estimate $+0.250$ clears $\epsilon \approx 0.151$ by roughly one σ , with its CI lower edge at $+0.079$ still inside the band on the right side of zero.




Figure 2: Forest plot of the ten numbered archetype desiderata. Each row carries a plain-language statement of the structural expectation (the case label C1 / C2 marks the factual observation it applies to); the dot is the bootstrap point estimate of the relevant Δ score, the horizontal bar is its 95% CI, and the green band is the row's *pass region*, the set of Δ values that would confirm that row's claim given the bootstrap-derived threshold $\epsilon \approx 0.151$. The right-hand column annotates the formula and its point estimate. All ten dots fall inside their pass bands. Underlying numerical computations are in the companion notebook `responsibility_archetypes.ipynb`.

Diagnostics that do not survive the bootstrap-derived threshold. Two claims from earlier drafts of this section, both about the preempted variable P , fail at the honest threshold and are reported as diagnostics rather than numbered desiderata. (D1) The cross-regime flip $\Delta_N(P, \text{Case 2}) - \Delta_N(P, \text{Case 1})$ measures $+0.139$ with CI $[-0.081, +0.365]$: the CI straddles zero, so P 's rise across regimes is not, on its own, statistically resolved at the production budget. The qualitative content of this claim is already carried by desiderata #8 (P at the noise floor in Case 1) and #10 (P above ϵ in Case 2): if both pass, the regime flip is a corollary, not an independent claim. (D2) The quantitative match $|\Delta_N(P) - \Delta_N(L_1)| < \epsilon$ in Case 2: the signed gap $\Delta_N(P) - \Delta_N(L_1)$ measures -0.267 with CI $[-0.441, -0.106]$, whose magnitude 0.267 is well above ϵ . At this sample budget, gated P does not quantitatively mirror ungated L_1 ; what we can claim is the weaker, qualitative statement that both clear the noise floor in Case 2, which is exactly desideratum #10 for P and built into Case 2 by construction for L_1 .

Where we go next. The archetype evaluation establishes that PCI recovers the right qualitative pattern on every standard causal motif against an analytic ground truth. Section 6 now turns to a complementary continuous-input comparison: how does PCI behave next to gradient-based attribution (Differential Causal Effect of Butler et al., 2022) when the same continuous cause is rescaled or moved out of the steep region of the response surface, where the gradient itself collapses but causal responsibility arguably should not?

6 Comparison to Differential Causal Effect

Having tested PCI against ground truth in Section 5, we now compare it with gradient-based attribution on a continuous-input credit-limit example.

A natural baseline for continuous-input attribution is to read causal influence off the gradient of an interventional mean function. The motivation is direct: $\mathbb{E}[Y \mid \text{do}(X=1)] - \mathbb{E}[Y \mid \text{do}(X=0)]$ is the binary ATE, and for a continuous cause one “arrives at derivative calculus” [Ness, 2025, Section 7.4.3] by letting the contrast shrink: $\partial_x \mathbb{E}[Y \mid \text{do}(X=x)]$. Generalised to a vector $\mathbf{x} = (x_1, \dots, x_d)$ with interventional mean $m(\mathbf{x}) = \mathbb{E}[Y_{\mathbf{X}=\mathbf{x}}]$, the *causal gradient* is

$$\nabla_{\mathbf{x}} m(\mathbf{x}) = \left(\partial_{x_1} \mathbb{E}[Y_{\mathbf{X}=\mathbf{x}}], \dots, \partial_{x_d} \mathbb{E}[Y_{\mathbf{X}=\mathbf{x}}] \right),$$

and each partial is the *differential causal effect* (DCE) of a single variable at $\mathbf{x}$. Butler et al. [2022] introduced this quantity in the Gaussian Process regression setting, where, with the posterior expected mean expressible as $\sum_n k(x, x^{(n)}) \alpha_n$, the DCE has the closed form $\partial_{x_i} \hat{F} = \sum_n \partial_{x_i} k(x, x^{(n)}) \alpha_n$.

PCI and DCE are not direct competitors (DCE estimates a rate of change, PCI estimates individual-level responsibility), but PCI inherits the territory of continuous attribution, so a contrast on a single example is informative. We highlight three differences:

1. DCE requires $y = F(\mathbf{X}, \epsilon)$ with F differentiable; PCI requires only that the causal model be sampleable, and the inference algorithm for the expanded model is the user’s choice (we use Monte Carlo throughout).
2. DCE inherits the units of both $\mathbf{X}$ and Y , so cross-feature comparison is unit-sensitive. The PCI score integrates over the necessity- and sufficiency-world densities, so changes to input scale leave the score essentially unchanged.¹⁸
3. DCE varies abruptly with the local geometry of F . The PCI score depends on where the factual value sits relative to the necessity/sufficiency densities and is therefore non-zero in flat regions while remaining locally informative.

Example 25 (credit-limit assignment). *We model an applicant’s approved credit limit as a function of age and the time of application. The deterministic component is a sigmoidal rise in age with a small sinusoidal modulation in time-of-day; intuitively, age should dominate. We consider four settings on two axes: time measured in hours vs. minutes (a units rescaling that should not affect attribution), and an age prior centred at 30 vs. 40 (a genuine change in what counts as unusual). The deterministic component is shown in Figure 3. Computational details and code are in docs/source/gradient_based_attribution.ipynb.*

PCI scores. We instantiate ci as the absolute-difference score from Example 21, $ci(y^s, y^n, y^*) = |y^n - y^*| - |y^s - y^*|$, and run the search of Section 3 on each setting, conditioning on age and on time of application as singleton suspects. Figure 4 shows the mechanism for the canonical setting (age-30 prior, hours): when age is the active suspect the necessity-world outcome tracks the deterministic curve while the sufficiency-world outcome stays near factual, and the same picture, with a smaller displacement, holds for time of application. Table 10 reports the mean total scores across all four settings. Two patterns stand out. First, age outscores time of application in every setting, by roughly a factor of 3/2, matching the intuition that the age sigmoid moves the outcome by a much larger absolute amount than the daily modulation does. Second, the scores are essentially unchanged between the hours and minutes columns, confirming the unit-invariance discussed above, whereas the shift from the age-30 to the age-40 prior raises every score, since a more unusual factual age means a larger search-space displacement.

PCI vs. DCE across the input grid. Figure 5 contrasts the two methods on the same grid of factual values (age-30 prior, hours); the other three settings give qualitatively the same picture. The

¹⁸This unit-invariance is contingent on how the alternative-value distribution Δ is specified. The ϵ -excised Δ of Section 3 is invariant to a reparameterisation of the input scale precisely when ϵ is fixed by the *probability mass* excised around the factual value (equivalently, in standardised units), as in the ROPE/effect-size reading given there, rather than as an absolute distance in the raw units of $\mathbf{X}$: an absolute-distance ϵ held fixed across a rescaling would itself become unit-dependent and reintroduce the very sensitivity PCI otherwise avoids.




Figure 3: Deterministic component of the credit-limit model (Example 25), on the hours scale: limit vs. age at several times of day (left) and limit vs. time of day at several ages (right). Age dominates the limit, with a small sinusoidal modulation from time of day; the minute scale is identical modulo a linear reparameterisation (verified in the notebook).




Figure 4: PCI search for the canonical setting (age-30 prior, hours), with age (left) and time of application (right) as the active suspect. Necessity-world (green) and sufficiency-world (orange) limit samples and the total score (purple) are plotted against the necessity-world value of the suspect; the dashed line marks the factual. The necessity world departs from factual along the deterministic response while the sufficiency world stays near it; each panel title gives the mean total score.

top panel plots the difference between the PCI scores for age and for time of application: it is positive everywhere, so age is consistently the more responsible variable, with the gap largest near the steep region of the age-limit sigmoid and growing for ages further from the prior mean. The bottom panels show the DCE for the two inputs, and two differences with PCI stand out. First, on the hour scale the DCE for time of application exceeds the DCE for age over most of the grid and traces a visible sine wave, whereas PCI reports age as the more responsible variable everywhere. Second, switching the time unit from hours to minutes flattens the time DCE by roughly a factor of 60, making the unit sensitivity explicit. Both are direct consequences of DCE’s role as a rate-of-change estimator rather than a responsibility score, and both make it harder to compare across features and across reparameterisations.

Where we go next. Sections 4–6 have positioned PCI against the dominant feature-attribution baselines (SHAP, Causal SHAP, gradient-based DCE) and a synthetic ground-truth catalogue. We now turn to the third leg of the comparison: the formal and empirical link between PCI and *actual causality*. Section 7 states a correspondence theorem identifying, for suitable choices of Γ and Δ , the configurations on which PCI is positive with Halpern’s actual causes; Section 8 compares PCI to Pearl’s *probability of actual causation*, the closest construction in the literature; Section 9 then measures the empirical gap in runtime and verdict recovery between exact AC enumeration and the necessity-only PCI estimator as model size grows.

Table 10: Mean PCI total scores for age and time of application across the four settings (time unit $\times$ age prior), at the factual instance. Age outscores time in every setting; the hours and minutes columns agree (unit-invariance), while the age-40 prior raises all scores relative to age-30 (a more unusual factual age).

Suspect	age-30 prior		age-40 prior	
	hours	minutes	hours	minutes
Age	100.45	97.95	177.87	176.56
Time of application	68.61	66.51	117.97	118.42




Figure 5: **PCI vs. DCE on the credit-limit grid** (age-30 prior, hours). *Top*: difference between the PCI scores for age and for time of application; positive everywhere, largest near the steep region of the age-limit sigmoid and for ages far from the prior mean, so age is always the more responsible variable. *Bottom*: DCE for age (left) and time of application (right) on both unit scales. Age has the larger absolute DCE only inside the narrow steep region of the sigmoid; elsewhere time of application dominates on the hour scale and goes comparatively flat on the minute scale, illustrating DCE’s shape and unit sensitivity.

7 Relation with Actual Causality

Where we are. Sections 4–6 have positioned PCI against feature-attribution baselines. We now turn to PCI’s relation to the causal-attribution tradition it inherits the witness mechanism from — Halpern’s actual causality. This section establishes the *formal* connection (Theorems 28–29 and Proposition 34); Section 8 adds the side-by-side comparison with Pearl’s *probability of actual causation*; and Section 9 measures the empirical gap in runtime and verdict recovery.

PCI is not meant to explicate Halpern’s notion of actual causality; the goal is to combine the intuitions underlying Halpern’s notion with those motivating Pearl’s probabilities of necessity and sufficiency to provide a more general tool for causal attribution. Nevertheless, to show that we preserve the spirit of Halpern’s definition, we now record the connection between actual causality and the joint necessity-sufficiency machinery of Section 3. The goal is to show that there is a natural choice of variable selection distribution Γ and alternative-value distribution Δ under which PCI verdicts agree with actual-causality verdicts. With one caveat: probability-of-causation maximisation will not identify subset-minimal causes that are not cardinality-minimal, a misalignment we characterise and remedy below.

Scope: a necessity-only correspondence. Halpern’s definition involves factivity, context-sensitive necessity, and minimality, but no sufficiency clause analogous to Pearl’s PS or PNS. The correspondence we establish in this section is therefore *necessity-only*: the sufficiency component of the kernel Φ defined below acts as a factivity check (restoring factual values must preserve φ under fixed noise), not as a substantive sufficiency-attribution component. The substantive sufficiency component of PCI, the Y^s factor in Definition 18 and the ci_S term in Section 5, is exercised in the synthetic overdetermination and undercutting evaluation (Section 5), the differential causal effect comparison (Section 6), and the SIR benchmark (Section 10), all of which are settings where AC cannot adjudicate. The narrowness of this section’s correspondence reflects what AC itself permits, not a limitation of PCI.

We first recall Halpern’s definition.

Definition 26 (Actual Cause, after Halpern, 2016). $C = c$ is an actual cause of φ in the causal setting $(M, \mathbf{u})$ if the following three conditions hold:

- **Factivity:** $(M, \mathbf{u}) \models (C = c)$ and $(M, \mathbf{u}) \models \varphi$,
- **Context-sensitive Necessity:** There is a set of variables $T \subseteq V$ and an alternative setting c' of variables in C such that if $(M, \mathbf{u}) \models T = t^*$, then
$$(M, \mathbf{u}) \models [C \leftarrow c', T \leftarrow t^*] \neg \varphi,$$
- **Minimality:** C is minimal; there is no strict subset $C_{\text{sub}} \subset C$ such that $C_{\text{sub}} = c_{\text{sub}}$ satisfies the above two conditions where c_{sub} is the restriction of c to the variables in C_{sub} .

To connect this to PCI, we instantiate the joint necessity-sufficiency machinery (Definition 18) at a single configuration (C, T) and a fixed noise realisation $\mathbf{u}$, and read off the configuration-level contribution. The resulting kernel combines the same three ingredients as Halpern’s definition: a sufficiency check at factual values, a necessity check against alternatives, and a Γ -weight that records how much the configuration is being asked about.

Definition 27 (AC-aligned PCI kernel). Fix a PSCM $M = \langle U, V, F, P_U \rangle$,¹⁹ a noise realisation $\mathbf{u}$, suspect and witness sets $S, W \subseteq V$ with factual values s^*, w^* , a variable selection distribution Γ on $2^S \times 2^W$ with mass function p^Γ , and an alternative-value distribution $\Delta(s^*)$. For an outcome event φ , active suspect set $C \subseteq S$ and active witness set $T \subseteq W$ with $T \cap C = \emptyset$, define

$$\begin{aligned} \Phi(C, T) &:= p^\Gamma(C, T) \cdot \mathbb{I}[(M, \mathbf{u}) \models [C \leftarrow s^* | C, T \leftarrow w^* | T] \varphi] \\ &\cdot \int \mathbb{I}[(M, \mathbf{u}) \models [C \leftarrow c', T \leftarrow w^* | T] \neg \varphi] \Delta_C(s^*)(dc'). \end{aligned} \quad (18)$$

Concretely, $\Phi(C, T)$ is the matched-configuration slice of the joint necessity-sufficiency measure $P_k^{s,n}$ from Definition 18: integration over P_U is replaced by point evaluation at $\mathbf{u}$, and both the sufficiency and the necessity sums are pinned to the same (C, T) pair, mirroring AC’s structure of testing factivity and the alternative against a single suspect/witness pair. The kernel is positive exactly when

¹⁹The kernel and the theorems below take the full PSCM (structural equations F plus noise distribution P_U) as given, since both the sufficiency and the necessity factors require sampling counterfactual outcomes under $\mathbf{u}$. As discussed in Section 1, this is a stronger commitment than approaches that bound PN/PS/PNS from observational data and a causal graph alone, but is the price of pointwise (not just bound) verdicts at the configuration level.

three conditions hold simultaneously: $(\mathbf{C}, \mathbf{T})$ has positive Γ -mass; restoring $\mathbf{C}$ to its factual values while holding $\mathbf{T}$ fixed preserves φ under $\mathbf{u}$ (the factivity-as-sufficiency component); and there exists a Δ -positive-density alternative $\mathbf{c}'$ for which the corresponding intervention overturns φ under $\mathbf{u}$ (the necessity component). The first factor takes the role played by the suspect/witness biases b_s, b_w in earlier work: regularity of Γ — positive p^Γ on every legal configuration — corresponds to the old $-0.5 < b_s, b_w < 0.5$ regularity. We will say that Γ is *regular* when $p^\Gamma(\mathbf{C}', \mathbf{T}') > 0$ for every $(\mathbf{C}', \mathbf{T}')$ with $\mathbf{C}' \subseteq \mathbf{S}, \mathbf{T}' \subseteq \mathbf{W}, \mathbf{C}' \cap \mathbf{T}' = \emptyset$.

We can now record the forward direction.

Theorem 28. *Suppose that $\mathbf{C}$ is an actual cause of φ with witness set $\mathbf{T}$ in $(M, \mathbf{u})$. Take any $\{\mathbf{S} = \mathbf{s}\} \supseteq \mathbf{C}$ as the potential cause set, subject to the constraint that no element of $\mathbf{S}$ is a deterministic function (under M) of any subset of the remaining elements of $\mathbf{S}$,²⁰ and any $\mathbf{W} \supseteq \mathbf{T}$ as the potential witness set. Assume Γ is regular and that $\Delta(\mathbf{s})$ assigns positive density to every $\mathbf{s}' \in \text{dom}(\mathbf{S}) \setminus \{\mathbf{s}\}$ (positive PMF in the discrete case, positive Lebesgue density in the continuous case). In the continuous case, assume additionally:*

- (i) *the structural equations of M downstream of $\mathbf{C}$ are continuous in $\mathbf{c}'$ under fixed $\mathbf{u}$;*
- (ii) *$\neg\varphi$ is open in $\text{dom}(Y)$ (or, more weakly, the boundary of $\neg\varphi$ has measure zero under the pushforward of Y).*

Then

$$\Phi(\mathbf{C}, \mathbf{T}) > 0, \quad (19)$$

and therefore

$$\sum_{\mathbf{T}' \subseteq \mathbf{W}} \Phi(\mathbf{C}, \mathbf{T}') > 0.$$

Proof. Fix $\mathbf{u}, \mathbf{c}'$, and $\mathbf{T}$ as witnesses to the satisfaction of the conditions in Definition 26. The context-sensitive-necessity clause forces $\mathbf{T} \cap \mathbf{C} = \emptyset$, since otherwise the joint intervention $[\mathbf{C} \leftarrow \mathbf{c}', \mathbf{T} \leftarrow \mathbf{t}^*]$ would be ill-defined. It remains to show that each of the three factors in $\Phi(\mathbf{C}, \mathbf{T})$ is strictly positive.

- *Selection weight.* By regularity of Γ , $p^\Gamma(\mathbf{C}, \mathbf{T}) > 0$.
- *Sufficiency indicator.* With exogenous noise $\mathbf{u}$ held constant, factivity gives $(M, \mathbf{u}) \models \varphi$. Restoring $\mathbf{C}$ to $\mathbf{s}^*|_{\mathbf{C}}$ and $\mathbf{T}$ to $\mathbf{w}^*|_{\mathbf{T}}$ leaves the model agreeing with $(M, \mathbf{u})$ on every variable, so the indicator evaluates to 1.
- *Necessity integral.* Context-sensitive necessity provides a specific $\mathbf{c}'$ witnessing $(M, \mathbf{u}) \models [\mathbf{C} \leftarrow \mathbf{c}', \mathbf{T} \leftarrow \mathbf{t}^*] \neg\varphi$. The closure condition on $\mathbf{S}$ and the positive-density assumption on $\Delta(\mathbf{s})$ imply that $\mathbf{c}'$ lies in the support of $\Delta_{\mathbf{C}}(\mathbf{s}^*)$. We treat the two cases separately. *Discrete case:* the singleton $\{\mathbf{c}'\}$ has positive PMF $\Delta_{\mathbf{C}}(\{\mathbf{c}'\}) > 0$ by hypothesis, so the integral is at least $\Delta_{\mathbf{C}}(\{\mathbf{c}'\}) > 0$. *Continuous case:* hypothesis (i) gives continuity of the intervention map $\mathbf{c}' \mapsto Y$ at fixed $\mathbf{u}$; hypothesis (ii) places $Y(\mathbf{c}')$ in the interior of $\neg\varphi$;

²⁰Equivalently, $\mathbf{S}$ is closed under no nontrivial structural equation of M . The constraint excludes admitting both S and a deterministic descendant $H = f(S, \dots)$ as candidates in the same suspect set, but covers all candidate sets standardly considered in the AC literature, where structural descendants of one candidate are not admitted as separate candidates: their contribution is mediated by the equation, and AC's minimality clause already prefers the ancestor alone. The same closure condition discharges a joint-support concern: the next assumption requires $\Delta(\mathbf{s})$ to place positive density on every *joint* alternative configuration $\mathbf{s}' \in \text{dom}(\mathbf{S}) \setminus \{\mathbf{s}\}$, not merely positive marginal density on each component. In a chain $S \rightarrow H$ with $H = f(S)$, joint assignments such as $(S=0, H=1)$ are impossible under the structural equations, so any Δ that placed mass on them would be incompatible with M . Closure rules out exactly this case: when no element of $\mathbf{S}$ is a deterministic function of the remaining elements, the joint domain $\text{dom}(\mathbf{S})$ contains no rectangles forced to zero measure by M , and a Δ with positive PMF or Lebesgue density on every $\mathbf{s}' \neq \mathbf{s}$ is consistent with M on the suspect set. In the stone-throwing benchmark of Section 9, the suspect set is $\{A_i, B_i\}_i$ and the deterministic mediators $\{W_i^a, W_i^b\}_i$ live in the witness set, not the suspect set, exactly to keep the suspect set closure-respecting. Witnesses are pinned at their factual values and are not subject to a joint-support requirement on alternatives, so they need no analogous condition.

pulling back an open neighbourhood within $\neg\varphi$ yields an open neighbourhood of $\mathbf{c}'$ on which the indicator equals 1, and this neighbourhood has positive Lebesgue measure and hence positive $\Delta_{\mathbf{C}}$ -mass by the positive-density hypothesis. In either case the integral is strictly positive.

The product of three strictly positive factors is strictly positive, giving $\Phi(\mathbf{C}, \mathbf{T}) > 0$. Summing over $\mathbf{T}' \subseteq \mathbf{W}$ adds non-negative terms to a positive one, preserving positivity. The proof uses factivity and context-sensitive necessity but not minimality: the same conclusion holds whenever $\mathbf{C}$ satisfies factivity and context-sensitive necessity in $(M, \mathbf{u})$, a fact we will use in Proposition 34. $\square$

Remark (when (i)–(ii) bite). In settings with continuous outcomes and an open event such as a threshold $\{Y > y_0\}$ (for instance the SIR overshoot benchmark of Section 10), both hypotheses are satisfied trivially. Binary-output models, where Y takes values in $\{0, 1\}$, satisfy (ii) automatically (every subset of a discrete codomain is open) but may not satisfy (i) when the binary output comes from thresholding a continuous score. In such cases the proof falls back to the discrete branch, which only requires positive PMF on the witnessing $\mathbf{c}'$.

The opposite direction follows by reading the proof in reverse on the necessity factor.

Theorem 29. *Under the same hypotheses on $\{\mathbf{S} = \mathbf{s}\}$, Γ , and Δ as in Theorem 28, if*

$$\sum_{\mathbf{T}' \subseteq \mathbf{W}} \Phi(\mathbf{C}, \mathbf{T}') > 0,$$

then both factivity $((M, \mathbf{u}) \models \varphi)$ and the context-sensitive-necessity condition in Definition 26 hold for $\mathbf{C}$ in $(M, \mathbf{u})$.

Proof. Since Φ is non-negative, the assumption forces $\Phi(\mathbf{C}, \mathbf{T}') > 0$ for at least one $\mathbf{T}' \subseteq \mathbf{W}$. By Definition 27, each factor of $\Phi(\mathbf{C}, \mathbf{T}')$ is strictly positive. The sufficiency indicator equals 1, i.e. $(M, \mathbf{u}) \models [\mathbf{C} \leftarrow \mathbf{s}^* |_{\mathbf{C}}, \mathbf{T}' \leftarrow \mathbf{w}^* |_{\mathbf{T}'}] \varphi$; since pinning variables to their factual values under fixed $\mathbf{u}$ leaves every variable unchanged, this is equivalent to $(M, \mathbf{u}) \models \varphi$, giving factivity. The necessity integral being positive provides a $\Delta_{\mathbf{C}}$ -positive-density set of values $\mathbf{c}'$ for which $(M, \mathbf{u}) \models [\mathbf{C} \leftarrow \mathbf{c}', \mathbf{T}' \leftarrow \mathbf{w}^* |_{\mathbf{T}'}] \neg\varphi$; any such $\mathbf{c}'$, together with $\mathbf{T}'$, witnesses the existential in the context-sensitive-necessity clause of Definition 26. $\square$

Remark (deterministic intermediates and joint support). The hypothesis on $\{\mathbf{S} = \mathbf{s}\}$ is what rules out the following obstruction. Suppose the suspect set were closed under the structural equations of M , e.g., both Sally’s throw S and the bottle’s hit $H = f(S, \dots)$ admitted as candidates. Then the joint assignment $(S=0, H=1)$ is structurally impossible, and any alternative-value distribution consistent with M assigns it density zero, so the joint full-support condition required to recover the witnessing $\mathbf{c}'$ from $\Phi > 0$ fails. Restricting suspects to variables that are not deterministic functions of the others under M removes this obstruction. The constraint is also conservative with respect to the AC literature: in the canonical stone-throwing setup, the candidate set is $\{S_{\text{Sally}}, S_{\text{Bill}}\}$, with H as the outcome rather than a candidate, and AC’s own minimality clause already prefers ancestor candidates over their deterministic descendants when both are admitted. Sections 5 and 9 adopt the same convention experimentally — the continuous synthetic benchmark uses the six roots as suspects and the discrete scaled-throwing benchmark uses $\{A_i, B_i\}_i$ as suspects with the deterministic mediators $\{W_i^a, W_i^b\}_i$ as witnesses — in both cases deterministic intermediates are excluded from the suspect set. Cases where a user genuinely wants to attribute responsibility separately to a deterministic intermediate (rather than to its structural ancestors) lie outside the scope of these theorems and require an alternative reading of intervention semantics; we treat that as a separate problem.

With minimality, the situation is somewhat more complicated. Our choice of Γ will have an impact on which configurations maximise Φ . Moreover, Φ compares configurations by Γ -weight rather than by the subset relation, while Definition 26 formulates minimality strictly in terms of subsets, so some misalignment is to be expected. Some useful observations can be made nonetheless.

Definition 30 (Necessity dilution). *A pair (M, Δ) satisfies necessity dilution on $\mathbf{S}$ if, writing $N(\mathbf{C}, \mathbf{T})$ for the necessity integral in Equation (18),*

$$N(\mathbf{C}^\sharp, \mathbf{T}^\sharp) \geq N(\mathbf{C}^\dagger, \mathbf{T}^\dagger)$$

for every $\mathbf{C}^\sharp \subsetneq \mathbf{C}^\dagger \subseteq \mathbf{S}$ and witnesses $\mathbf{T}^\sharp, \mathbf{T}^\dagger$ disjoint from the corresponding suspect set.

The first is that there is a simplified setting in which a positive value of Φ , under factivity, entails minimality with respect to the same parameters. Three conditions suffice:

- Γ_w is uniform over $2^{\mathbf{W}}$;
- Γ_s is *strictly cardinality-decreasing*, i.e. $p^{\Gamma_s}(\mathbf{C}_1) > p^{\Gamma_s}(\mathbf{C}_2)$ whenever $\mathbf{C}_1 \subsetneq \mathbf{C}_2$ (the modern counterpart of the old null witness bias and positive suspicion bias);
- the pair (M, Δ) satisfies necessity dilution on $\mathbf{S}$ (Definition 30).

The strategy used in the proof entails that we can safely marginalise over witnesses as well.

Theorem 31. *Consider the set of optimisers $\mathcal{O}^* = \arg \max_{\mathbf{C} \subseteq \{\mathbf{S}=\mathbf{s}\}, \mathbf{T} \subseteq \mathbf{W}} \Phi(\mathbf{C}, \mathbf{T})$. Assume $(\mathbf{C}^\dagger, \mathbf{T}^\dagger) \in \mathcal{O}^*$ with $\Phi(\mathbf{C}^\dagger, \mathbf{T}^\dagger) > 0$, that factivity is satisfied for $\mathbf{C}^\dagger$ (as in Definition 26), and that no element of $\{\mathbf{S} = \mathbf{s}\}$ is a deterministic function under M of the others. Let Γ_w be uniform on $2^{\mathbf{W}}$ and let Γ_s be strictly cardinality-decreasing, and assume that (M, Δ) satisfies necessity dilution on $\mathbf{S}$. Then $\mathbf{C}^\dagger$ is an actual cause of φ according to Definition 26.*

Proof. Factivity is assumed; by Theorem 29 the context-sensitive-necessity condition holds for $\mathbf{C}^\dagger$ in $(M, \mathbf{u})$. So the only way $\mathbf{C}^\dagger$ can fail to be an actual cause is by failing minimality. Suppose for contradiction that minimality fails: there is $\mathbf{C}^\sharp \subsetneq \mathbf{C}^\dagger$ satisfying factivity and context-sensitive necessity for φ with some witness $\mathbf{T}^\sharp$. By Theorem 28 (its proof requires only factivity and necessity, not minimality), $\Phi(\mathbf{C}^\sharp, \mathbf{T}^\sharp) > 0$ as well, with the sufficiency indicator and the necessity integral both equal to 1 and a strictly positive value, respectively. Both pairs therefore satisfy $\Phi(\mathbf{C}, \mathbf{T}) = p^\Gamma(\mathbf{C}, \mathbf{T}) \cdot N(\mathbf{C}, \mathbf{T})$.

By the construction of Γ from independent marginals (Section 3),

$$p^\Gamma(\mathbf{C}, \mathbf{T}) \propto p^{\Gamma_s}(\mathbf{C}) p^{\Gamma_w}(\mathbf{T}) \mathbb{I}[\mathbf{T} \cap \mathbf{C} = \emptyset].$$

The $\mathbb{I}[\cdot]$ factor is 1 for both pairs (otherwise the contradiction would have been immediate). Uniformity of Γ_w on $2^{\mathbf{W}}$ makes $p^{\Gamma_w}(\mathbf{T}) = 2^{-|\mathbf{W}|}$ identically, so the witness factor cancels. Strict cardinality-decrease of Γ_s together with $\mathbf{C}^\sharp \subsetneq \mathbf{C}^\dagger$ gives $p^{\Gamma_s}(\mathbf{C}^\sharp) > p^{\Gamma_s}(\mathbf{C}^\dagger)$, and necessity dilution gives $N(\mathbf{C}^\sharp, \mathbf{T}^\sharp) \geq N(\mathbf{C}^\dagger, \mathbf{T}^\dagger)$. Multiplying these (and noting $N(\mathbf{C}^\sharp, \mathbf{T}^\sharp) > 0$ to convert weak into strict in the composite) gives $\Phi(\mathbf{C}^\sharp, \mathbf{T}^\sharp) > \Phi(\mathbf{C}^\dagger, \mathbf{T}^\dagger)$, contradicting $(\mathbf{C}^\dagger, \mathbf{T}^\dagger) \in \mathcal{O}^*$. $\square$

Corollary 32. *Keep the notation and hypotheses of Theorem 31. Consider the set of optimisers $\mathcal{O}^\Sigma = \arg \max_{\mathbf{C} \subseteq \{\mathbf{S}=\mathbf{s}\}} \sum_{\mathbf{T} \subseteq \mathbf{W}} \Phi(\mathbf{C}, \mathbf{T})$. Assume $\mathbf{C}^\dagger \in \mathcal{O}^\Sigma$ with $\sum_{\mathbf{T} \subseteq \mathbf{W}} \Phi(\mathbf{C}^\dagger, \mathbf{T}) > 0$ and that factivity is satisfied for $\mathbf{C}^\dagger$. Then $\mathbf{C}^\dagger$ is an actual cause of φ according to Definition 26.*

Proof. The summation hypothesis entails the existence of some $\mathbf{T}^\dagger \subseteq \mathbf{W}$ with $\Phi(\mathbf{C}^\dagger, \mathbf{T}^\dagger) > 0$. Because the slice $\{(\mathbf{C}^\dagger, \mathbf{T}) : \mathbf{T} \subseteq \mathbf{W}\}$ is contained in the joint search space $2^{\mathbf{S}} \times 2^{\mathbf{W}}$, we have $(\mathbf{C}^\dagger, \mathbf{T}^\dagger) \in \mathcal{O}^*$ as well. The conditions of Theorem 31 are satisfied, so $\mathbf{C}^\dagger$ is an actual cause of φ . $\square$

When necessity dilution holds. The hypothesis is a property of the pair (M, Δ) , not of Γ . It says that adding suspects to a candidate cause does not introduce new Δ -mass of flipping alternatives, and three settings make it automatic. (i) *Single-pivot models*: one variable $X^* \in \mathbf{S}$ is pivotal under $(M, \mathbf{u})$ and the others are structurally idle, so N peaks at $\{X^*\}$ and falls off as more suspects are added. (ii) *Factored Δ without Boolean-OR redundancy*: $\Delta_{\mathbf{C}}$ is a product of marginals $\bigotimes_{X \in \mathbf{C}} \nu_X$ with $\nu_X(\text{dom}(X)) \leq 1$, and the flip event factorises coordinate-wise; $N(\mathbf{C})$ is then a product of factors each at most 1, decreasing in $|\mathbf{C}|$. (iii) *Conjunctive outcome events*: if $\neg\varphi$ requires flipping every variable in a required set $\mathbf{R} \subseteq \mathbf{S}$, the flip region's $\Delta_{\mathbf{C}}$ -mass decreases as $\mathbf{C}$ grows beyond $\mathbf{R}$. The hypothesis fails in disjunctive overdetermination (e.g. $\varphi : A \vee B = 1$ with both A, B factual): no proper subset of $\mathbf{S}$ flips φ , so N is zero on subsets and positive on the full set: the opposite of dilution. In such cases no proper subset of $\mathbf{S}$ is an actual cause anyway, so Theorem 31 is not the appropriate tool; Proposition 34 is.

We can now return to the misalignment we mentioned earlier. One can imagine a reasoning mode in which one looks for causes of an outcome by inspecting the integrated probability $P(\mathbf{C} \rightsquigarrow \varphi) :=$

$\int \sum_{\mathbf{T}} \Phi(\mathbf{C}, \mathbf{T}) P_{\mathbf{U}}(d\mathbf{u})$, not knowing what the values of $\mathbf{u}$ are but knowing that φ occurred. It is worth pointing out that $\arg \max_{\mathbf{C} \subseteq \mathbf{S}} P(\mathbf{C} \rightsquigarrow \varphi)$ might fail to include some $\mathbf{C}$ that is an actual cause of φ for some setting $\mathbf{u}$.

Observation 33. Define $\Xi^{\max} = \arg \max_{\mathbf{C} \subseteq \mathbf{S}} P(\mathbf{C} \rightsquigarrow \varphi)$. It is not the case that if $\mathbf{C}^\dagger$ is an actual cause in $(M, \mathbf{u})$ for some $\mathbf{u}$, $\mathbf{C}^\dagger \in \Xi^{\max}$.

Proof. Consider a model with two binary parents A, B uniformly distributed and a binary child $Y = A \vee B$. Take $\varphi : Y = 1$ and the suspect set $\mathbf{S} = \{A, B\}$ with $\mathbf{W} = \emptyset$. Compare $\mathbf{C}^\dagger = \{A, B\}$ at noise $\mathbf{u}_{11}$ ($A = B = 1$) and $\mathbf{C}^\sharp = \{A\}$ at noise $\mathbf{u}_{10}$ ($A = 1, B = 0$).

- In $\mathbf{u}_{11}$, $\mathbf{C}^\dagger$ is an actual cause: setting both parents to 0 yields $Y = 0$, no proper subset achieves this, and the indicators in Φ are 1. With Γ_w uniform on $2^\emptyset$ (a single configuration of weight 1) and a strictly cardinality-decreasing Γ_s , $\Phi(\mathbf{C}^\dagger, \emptyset) \propto p^{\Gamma_s}(\{A, B\})$.
- In $\mathbf{u}_{10}$, $\mathbf{C}^\sharp$ is an actual cause: intervening to $A = 0$ flips Y to 0, the singleton is minimal, and again the indicators are 1. So $\Phi(\mathbf{C}^\sharp, \emptyset) \propto p^{\Gamma_s}(\{A\})$.

Integrating uniformly over the four noise realisations $\{\mathbf{u}_{ij}\}$, $P(\mathbf{C}^\dagger \rightsquigarrow \varphi) \propto p^{\Gamma_s}(\{A, B\})$ (only $\mathbf{u}_{11}$ contributes) while $P(\mathbf{C}^\sharp \rightsquigarrow \varphi) \propto 2p^{\Gamma_s}(\{A\})$ ($\mathbf{u}_{10}$ and $\mathbf{u}_{11}$ both contribute). Whether $\mathbf{C}^\dagger$ or $\mathbf{C}^\sharp$ ends up in $\Xi^{\max}$ depends on the precise shape of Γ_s : under sharply decreasing Γ_s , $\mathbf{C}^\sharp$ wins; under shallow decrease, $\mathbf{C}^\dagger$ may. Both cannot belong to $\Xi^{\max}$ simultaneously in general, so some actual causes are dropped. $\square$

The reason is structural: actual cause sets may be subset-minimal without being cardinality-minimal, while integrated-probability maximisation is governed by Γ_s , which (under the cardinality-decreasing parameterisation that aligns with AC’s minimality) penalises larger sets. The misalignment is therefore not a defect of Φ but a mismatch between AC’s notion of minimality (subset relation) and Γ -weighted scoring.

The fix is to drop the maximisation step and use Φ directly to enumerate candidate causes, then filter for subset-minimality. The filter recovers exactly the actual causes in $\mathbf{S}$.

Proposition 34. Under the hypotheses of Theorem 28, define

$$\Xi := \left\{ \mathbf{C}^\dagger \subseteq \mathbf{S} : \sum_{\mathbf{T} \subseteq \mathbf{W}} \Phi(\mathbf{C}^\dagger, \mathbf{T}) > 0, \nexists \mathbf{C}' \subsetneq \mathbf{C}^\dagger \text{ with } \sum_{\mathbf{T} \subseteq \mathbf{W}} \Phi(\mathbf{C}', \mathbf{T}) > 0 \right\}.$$

Then Ξ contains exactly the actual causes of φ in $(M, \mathbf{u})$ that are subsets of $\mathbf{S}$.

Proof. Ξ -membership $\Rightarrow$ actual cause. Let $\mathbf{C}^\dagger \in \Xi$. Theorem 29 applied to $\sum_{\mathbf{T}} \Phi(\mathbf{C}^\dagger, \mathbf{T}) > 0$ gives both factivity and context-sensitive necessity. For minimality, suppose $\mathbf{C}_{\text{sub}} \subsetneq \mathbf{C}^\dagger$ satisfied factivity and context-sensitive necessity. By the proof of Theorem 28 (which uses only factivity and necessity, not minimality), we would have $\sum_{\mathbf{T}} \Phi(\mathbf{C}_{\text{sub}}, \mathbf{T}) > 0$, contradicting the no-positive-proper-subset clause of Ξ .

Actual cause $\Rightarrow \Xi$ -membership. Let $\mathbf{C}^\dagger \subseteq \mathbf{S}$ be an actual cause of φ in $(M, \mathbf{u})$. Theorem 28 gives $\sum_{\mathbf{T}} \Phi(\mathbf{C}^\dagger, \mathbf{T}) > 0$. For the no-positive-proper-subset clause: suppose $\mathbf{C}' \subsetneq \mathbf{C}^\dagger$ had $\sum_{\mathbf{T}} \Phi(\mathbf{C}', \mathbf{T}) > 0$. Then by Theorem 29, both factivity and context-sensitive necessity would hold for $\mathbf{C}'$. So $\mathbf{C}'$ would witness a violation of $\mathbf{C}^\dagger$ ’s minimality, contradicting the assumption that $\mathbf{C}^\dagger$ is an actual cause. $\square$

Proposition 34 converts the cardinality–subset misalignment into a procedure: enumerate $\{\mathbf{C} \subseteq \mathbf{S} : \sum_{\mathbf{T}} \Phi(\mathbf{C}, \mathbf{T}) > 0\}$ (which by Theorem 29 are exactly the cause sets satisfying context-sensitive necessity), filter for factivity and subset-minimality, and read off the actual causes. Some idiosyncratic features of the AC notion (in particular, the subset-minimality clause, which interacts poorly with Γ -weighted scoring) are part of what motivated the more scalable PCI formulation of Section 3 in the first place; the filter procedure here is the right tool when AC verdicts specifically are wanted, but expectation-based PCI attributions are what scale to ML-grade models.

Summary of guarantees. The section delivers, under explicit hypotheses on $\mathbf{S}$, Γ , and Δ : (i) a forward direction (Theorem 28) “actual cause $\Rightarrow \Phi > 0$ ”; (ii) a backward direction (Theorem 29) “ $\sum_{\mathbf{T}} \Phi > 0 \Rightarrow$ context-sensitive necessity”; (iii) minimality preservation under cardinality-decreasing Γ_s (Theorem 31, Corollary 32); (iv) a counterexample (Observation 33) showing that integration over noise misaligns with subset-minimality; and (v) a corrective filter procedure (Proposition 34) that recovers AC verdicts exactly. The empirical counterpart of the correspondence, a benchmark of the necessity-only PCI estimator against exact subset enumeration on a scaled stone-throwing problem, is reported in Section 9.

Where we go next. Before turning to the empirical benchmark, the next section locates PCI against the construction in the literature closest to its expectation form: Pearl’s *probability of actual causation*. The two share the same surface idea, “how often, under the noise, does the actual-cause verdict fire?”, and they coincide on standard examples, but they disagree on graded responsibility patterns that PCI registers and Pearl’s binary AC verdict discards. Section 8 walks through Pearl’s canonical desert-traveller illustration and a weak-poison variant to make the distinction concrete.

8 Relation with the Probability of Actual Causality

Now we are in the position to discuss the relation between PCI and Pearl’s probability of actual causation. On Pearl’s Definition 10.3.5 [Pearl, 2009, Ch. 10],

$$P(\text{caused}(x, y \mid e)) = \frac{P(U_{xy} \cap U_e)}{P(U_e)}, \quad U_{xy} = \{\mathbf{u}: \text{AC}(x, y; M, \mathbf{u})\}, \quad (20)$$

is the posterior probability, given the evidence, that we are in a noise state for which the actual-cause verdict holds, with $\text{AC}(x, y; M, \mathbf{u})$ a chosen actual-causation predicate: the causal-beam definition of [Pearl, 2009, Ch. 10] or the Halpern–Pearl AC1–AC3 [Halpern, 2016].²¹ The same construction underwrites Chockler and Halpern’s degree of blame [Chockler and Halpern, 2004], which substitutes a graded responsibility share inside the expectation. Because (20) is the closest construction in the literature to PCI’s expectation in Definition 18, we now ask in detail how it differs from PCI and how PCI improves on it.

We proceed in three steps. We replay Pearl’s canonical desert-traveller illustration and Pearl’s binary actual-cause verdicts; compute PCI on the same example and show it recovers Pearl’s ranking with graded magnitudes; and introduce a weak-poison variant that exhibits three intuitive responsibility patterns, two of which Pearl’s binary verdict cannot capture but PCI’s graded reading does, and one (within-scenario ranking) Pearl preserves. Then we move to discuss other differences between PCI and Pearl’s probability of causation, which make the former more general and more scalable.

8.1 Pearl’s Worked Example: The Desert Traveller

We start with the canonical illustration used by Pearl [Pearl, 2009, §10.3.3].

Example 35 (Desert traveller). *A traveller has two enemies. The shooter (enemy 2) shoots and empties the canteen ($X=1$); the poisoner (enemy 1), unaware, puts cyanide in the water ($P=1$). The traveller dies ($Y=1$). Who is the actual cause of death?*

The example involves one source of uncertainty packed into a binary noise u :

- $u = 0$: traveller drank the poisoned water before the canteen was emptied; cyanide killed him.
- $u = 1$: canteen was empty before he could drink; dehydration killed him.

There are at least two intuitions here that we would like to recover. One, in any particular case, we want to be able to identify the actual cause. The other, perhaps a bit weaker, we still don’t want to

²¹All computations behind the comparison in this section, Pearl’s original desert-traveller and the weak-poison variant, are collected in the companion notebook `docs/source/desert_traveler.ipynb`, which runs the PCI Monte Carlo search of Section 3 on a scenario-specific structural model for each forensic posterior alongside a parallel hand-rolled enumeration; the two agree within Monte Carlo noise on every value reported below.

say that the dormant cause is completely blameless. Actual causality gives us the first half of the story, assigning full responsibility to one source here. Pearl’s beam analysis [Pearl, 2009, Ch. 10] (equivalently Halpern–Pearl AC1–AC3 of Halpern, 2016) gives the actual-cause verdicts at each deterministic state:

- In $u = 0$: $P=1$ is the actual cause; $X=1$ is not.
- In $u = 1$: $X=1$ is the actual cause; $P=1$ is not.

So $U_{x,y} = \{u = 1\}$ and $U_{p,y} = \{u = 0\}$. Two natural forensic reports each push Pearl’s posterior to certainty (Table 11): toxicology finding cyanide in the body is incompatible with $u = 1$, so $P(\text{caused poisoner} \mid e_A) = 1$; a forensic report of no cyanide is incompatible with $u = 0$, so $P(\text{caused shooter} \mid e_B) = 1$.

Forensic posterior	$P(\text{caused shooter})$	$P(\text{caused poisoner})$
e_A : cyanide detected ($u = 0$)	0	1
e_B : no cyanide ($u = 1$)	1	0

Table 11: Pearl’s probability of actual causality $P(\text{caused} \mid e)$ on the basic desert traveller. The verdict saturates at 1 for the cause whose path operated in the forensically identified scenario, and at 0 for the dormant one.

8.2 PCI on the Basic Desert Traveller

To compare directly with Pearl, we run PCI under the same forensic conditioning: one structural model per scenario (endogenous variables $X, P, c, d, Y \in \{0, 1\}$, with c the cyanide path operating and d the dehydration path), with u pinned at the scenario’s forensic realisation. Suspects are $\mathbf{S} = \{X, P\}$ with singleton candidate causes; the witness pool is the mediators $\{c, d\}$ under a cardinality-uniform Γ ; Δ deterministically flips $1 \rightarrow 0$; the non-cause suspect is left to the SCM rather than intervened on; and the noise distribution P_U is the forensic posterior, a point mass at the scenario’s u (we do *not* condition on the outcome Y itself). The full structural model and component specification are given in Appendix C.

For each candidate cause C we report three quantities, all in $[0, 1]$,²² higher meaning more responsibility, jointly averaged over the four sources of randomness (Γ , Δ , P_U , and the SCM-driven non-cause suspect):

$$\begin{aligned}
N_C &= \mathbb{E}[\mathbb{1}\{Y^n \neq y^*\}] && \text{(necessity — removal flips } Y), \\
S_C &= \mathbb{E}[\mathbb{1}\{Y^s = y^*\}] && \text{(sufficiency — holding at factual sustains } Y), \\
J_C &= \mathbb{E}[\mathbb{1}\{Y^n \neq y^*\} \mathbb{1}\{Y^s = y^*\}] && \text{(joint PNS-style).}
\end{aligned} \tag{21}$$

Under the spec above (computation in the companion notebook), the results are:

Forensic posterior	Cause	N	S	J	Pearl $P(\text{caused})$
e_A : $u = 0$ (cyanide)	shooter	1/4	11/12	5/24	0
	poisoner	1/3	1	1/3	1
e_B : $u = 1$ (dehydration)	shooter	1/3	1	1/3	1
	poisoner	1/4	11/12	5/24	0

Table 12: PCI and Pearl on the basic desert traveller, under the scenario’s forensic posterior. PCI’s joint score J ranks the same cause Pearl’s $P(\text{caused})$ does in each scenario; PCI returns a graded reading where Pearl returns the binary $\{0, 1\}$.

²²These are evaluated for a fixed candidate cause C , hence normalised to $[0, 1]$; this differs from the sub-probability $\mathbb{E}[c_i]$ of Definition 19, whose total mass is $\Pr_{\Gamma}[X_k \in \mathbf{C}] \leq 1$ because it sums only over suspect sets that contain X_k .

Three points to read off Table 12:

(1) PCI and Pearl agree on the ranking. In each scenario, J is strictly larger for the cause Pearl identifies, and strictly smaller for the other. PCI’s ordering matches the actual-cause verdict at each forensic posterior.

(2) PCI is graded; Pearl is binary. Pearl’s $P(\text{caused})$ saturates at 0 or 1. PCI’s J never reaches 1 here because the Γ -averaging over witness subsets dilutes each cause’s score: the “best” witness subset for the operative cause does flip the outcome deterministically and gives $J = 1$ at that single configuration, but the cardinality-uniform average over the witness subsets (weights $\frac{1}{3}, \frac{1}{6}, \frac{1}{6}, \frac{1}{3}$ over $\emptyset, \{c\}, \{d\}, \{c, d\}$) dilutes this to $1/3$; the per-subset breakdown is in the companion notebook.

(3) The dormant cause is not at zero. PCI assigns $5/24 \approx 0.21$ to the dormant cause. This is not noise: in some witness configurations the dormant cause’s removal does flip Y , because the non-cause suspect is marginalised and the model is not artificially pinned to “this scenario” beyond the noise realisation itself. Pearl’s binary indicator throws this graded information away.

8.3 PCI Expectation vs the AC Machinery

The agreement above is reassuring, but the more interesting observation is what changes underneath. AC’s necessity clause is an *existential*: *there exists* a witness set $\mathbf{T}$ and an alternative value c' such that the intervention flips Y . To verify the existential one in principle enumerates the candidate $(\mathbf{T}, c')$ pairs: this is the main source of AC’s combinatorial hardness. Pearl’s definition then averages the indicator of that already-enumerated predicate over noise states, inheriting the enumeration unchanged.

PCI collapses both moves into one expectation: Φ is averaged over the joint $(\mathbf{C}, \mathbf{T}, \mathbf{u})$ under $\Gamma \otimes P_{\mathbf{U}}$. The existential “does *some* witness work?” becomes the quantitative “*what fraction* of witnesses work?”, the information AC uses only one piece of, exposed as a graded score with sampling-based estimation.

In the desert traveller this is small change: four candidate witness sets, all trivially enumerable. The point is the scaling. In a model with n candidate witness variables, AC’s existential demands checking up to 2^n subsets per noise state, and Pearl’s posterior over the AC indicator inherits that cost per evaluation. PCI’s expectation can be estimated at a Monte Carlo budget the user controls, with variance independent of the problem’s combinatorial size. The desert-traveller agreement is the small- n end of a curve whose large- n end is the computational gap that motivates PCI in the first place.

So far we have two conceptual differences: PCI’s graded reading and its Monte Carlo scaling. We have already seen that this helps to recover the intuition that the shooter of a poisoned victim, while less blameworthy is nevertheless not completely blameless. Now we turn to another example, riding on the fact that actual causality is devoid of a sufficiency component, where a small change to the model makes Pearl’s binary indicator structurally unable to track the responsibility patterns we think a responsibility attribution method should capture.

8.4 Weak-Poison Variant: Two Intuitions Pearl Misses

We modify the desert traveller so the cyanide path is unreliable while the dehydration path stays deterministic. The change is small, but it surfaces three intuitions that a graded responsibility metric should capture and a binary actual-cause indicator structurally cannot.

Example 36 (Desert traveller, weak poison). *Same enemies as in Example 35. The cyanide is a small dose that turns out fatal only with probability $\alpha = 0.1$, governed by an independent exogenous coin $\xi \sim \text{Bern}(\alpha)$. The structural equations are*

$$c = p(u' \vee x'), \quad v_C = c \cdot \xi, \quad d = x(u \vee p'), \quad y = v_C \vee d,$$

where primes here denote Boolean complement ($x' = 1 - x$, and likewise u', p'), not the alternative-value c' of the necessity intervention, and v_C is the new “cyanide-was-fatal” mediator: drinking the poison ($c = 1$) only kills when the fatality coin also operates ($\xi = 1$). $\alpha = 1$ recovers Pearl’s original.

Three responsibility intuitions for an attribution method. Before any computation, the three intuitions that we expect an attribution method to recover in this case are:

1. **Adding noise to a path should lower its own cause’s responsibility.** The cyanide path is now unreliable. In the scenario where cyanide killed (Scenario A, $u = 0, \xi = 1$), the poisoner is still the actual cause, but *less* responsible than in the basic deterministic version, because the killing depended on a coin landing favourably.
2. **Within a scenario, the operative cause should still win.** In Scenario A, even after softening, the poisoner should rank above the shooter. In Scenario B, the shooter should rank above the poisoner.
3. **A reliable cause in its own scenario should beat an unreliable cause in its own scenario.** Comparing across scenarios: the shooter in Scenario B (dehydration deterministic) should outrank the poisoner in Scenario A (cyanide stochastic). Their respective paths are operative in their own scenarios, but one is reliable and the other is not.

Pearl’s $P(\text{caused})$ captures (2) (the within-scenario ranking), but misses (1) and (3). The AC predicate fires deterministically once we condition on the forensic posterior: in Scenario A, $u = 0$ and $\xi = 1$ jointly imply the cyanide path was the operative chain, so $P(\text{caused poisoner} \mid e_A) = 1$; in Scenario B, $u = 1$ implies the dehydration chain operated, so $P(\text{caused shooter} \mid e_B) = 1$. Both scenarios saturate at 1 for their respective operative cause, identical to the basic version (Table 13). Pearl’s binary indicator is structurally indifferent to whether the cyanide path is deterministic or stochastic, and indifferent to a cross-scenario comparison.

Forensic posterior	$P(\text{caused shooter})$	$P(\text{caused poisoner})$
e_A : cyanide killed ($u = 0, \xi = 1$)	0	1
e_B : dehydration killed ($u = 1$)	1	0

Table 13: Pearl’s $P(\text{caused})$ on the weak-poison variant. Identical to the basic version (Table 11); the binary AC indicator is insensitive to α .

What PCI does on the same model. We run PCI exactly as in Section 8.2, with one mechanical change: the witness pool becomes $\mathbf{W} = \{c, d, v_C\}$ (the new mediator v_C is included), and the forensic posterior in Scenario A pins both $u = 0$ and $\xi = 1$ (cyanide-was-fatal is directly forensically observable). Scenario B’s posterior pins $u = 1$; ξ is structurally irrelevant since the cyanide path is dormant.

Table 14 reports PCI’s three quantities alongside Pearl’s verdict. The values are computed in the companion notebook and verified by a hand-rolled enumeration matching the framework.

Forensic posterior	Cause	N	S	J	Pearl $P(\text{caused})$
e_A : $u = 0, \xi = 1$ (cyanide killed)	shooter	≈ 0.167	≈ 0.958	≈ 0.146	0
	poisoner	≈ 0.208	1	$\approx \mathbf{0.208}$	1
e_B : $u = 1$ (dehydration killed)	shooter	1/2	1	1/2	1
	poisoner	1/4	≈ 0.750	≈ 0.125	0

Table 14: PCI and Pearl on the weak-poison desert traveller, under the scenario’s forensic posterior. PCI’s J ranks the same cause Pearl’s binary verdict does in each scenario; the magnitudes also reflect the asymmetry between the deterministic dehydration path and the stochastic cyanide path.

The three intuitions are all captured by J . **(1) Adding noise lowers the noisy cause’s responsibility.** The poisoner in Scenario A drops from $J = 1/3$ in the basic version (Table 12) to $J \approx 0.208$ in the weak version (Table 14). The drop comes through the sufficiency factor: when we hold P at its factual value, the witness mechanism averages over configurations where v_C is not pinned and the cyanide step’s ξ coin still acts. Even at $\xi = 1$ in the forensic posterior, the witness integration

sees the fragility, and J falls. **(2) Within a scenario, the operative cause still wins.** In Scenario A, $J_{\text{poisoner}} \approx 0.208 > J_{\text{shooter}} \approx 0.146$. Cyanide was the operative path, and PCI tracks that. In Scenario B, $J_{\text{shooter}} = 1/2 > J_{\text{poisoner}} \approx 0.125$. **(3) Reliable beats unreliable, across scenarios.** The shooter in Scenario B ($J = 1/2$) outranks the poisoner in Scenario A ($J \approx 0.208$). The dehydration path is deterministic; conditional on $u = 1$, holding X at factual sustains Y with probability 1 regardless of the witness regime, so the shooter is a textbook responsible cause. The cyanide path is fragile; conditional on $u = 0, \xi = 1$, the sufficiency factor still sees the ξ coin under unpinned witness regimes, so the poisoner cannot hit the same ceiling. Pearl’s PAC misses the desiderata for the following reasons. (1) requires sensitivity to mechanism noise, which the AC indicator does not have once the forensic posterior is applied; (3) requires cross-scenario comparison of *strength of responsibility*, which Pearl’s $P(\text{caused})$ does not produce because each posterior independently saturates at 0 or 1.

8.5 Further Differences Between Pearl’s Probability of Causation and PCI

Breadth of aggregation. Def. 10.3.5 aggregates over one dimension: the noise $\mathbf{u}$, conditional on evidence e , for a claim $(X = x, Y = y)$ already fixed. PCI aggregates over four: the suspect set $\mathbf{C}$ and witness set $\mathbf{T}$ via Γ , the alternative value $\mathbf{c}'$ via Δ , and the noise via $P_{\mathbf{U}}$. This is what supports per-feature comparison: the same PCI expectation, evaluated over varying $\mathbf{C}$ inside a fixed $\mathbf{S}$, returns the relative responsibility of each candidate cause. Def. 10.3.5 cannot perform that comparison without a separate evaluation per claim, each inheriting the AC-predicate cost discussed above.

Continuous variables. The AC predicates inside Def. 10.3.5 are event-shaped: they read off $\text{AC}(x, y; M, \mathbf{u})$ for propositional x, y , and extensions to continuous variables require discretising the relevant event before the predicate applies. PCI’s *ci* function is continuous-native (the absolute-difference impact score of Section 3), and the synthetic benchmark of Section 5 exercises that continuity directly.

Lineage in Pearl’s PN/PS. Def. 10.3.5 takes a binary actual-causation predicate — itself a structural-equation construction that does not appeal to PN/PS at all — and lifts it to a probability through $P(\mathbf{u} \mid e)$; PN, PS, and PNS appear nowhere in the construction. PCI, by contrast, generalises PN and PS directly: the necessity factor Y^n is the counterfactual operation underlying PN, the sufficiency factor Y^s is the counterfactual operation underlying PS, both indexed by the cause–witness–alternative–noise tuple, with Γ and Δ exposing the design choices PN/PS makes implicitly. So Pearl’s Def. 10.3.5 and PCI are not competing implementations of the same idea: they inherit from different parts of Pearl’s own toolkit, with PCI playing the role of a context-sensitive, multi-variable extension of the PN/PS machinery rather than a probabilistic wrapper around AC.

Pearl’s Def. 10.3.5 and PCI share the same explanatory target, probabilistic responsibility built on top of structural information, but instantiate it through different mechanisms. Pearl’s definition averages a binary AC indicator against a posterior over noise; PCI integrates a continuous necessity–sufficiency kernel against a joint distribution over suspect sets, witness sets, alternative values, and noise. The basic desert traveller shows the two agree on ranking with PCI providing graded magnitudes; the weak-poison variant shows three intuitions that PCI’s graded reading captures and Pearl’s binary indicator structurally cannot. The cases where sufficiency does even more graded work — the continuous signal-mediation walkthrough of Section 4.3, the synthetic archetypes of Section 5, and the SIR benchmark of Section 10 — are where these constructions answer recognisably different questions, and the question PCI answers is the one we need for ML-grade causal attribution.

Where we go next. Sections 7–8 have laid out the formal relation between PCI and actual causality (and its Pearlian probabilistic wrapper). Section 9 now turns to the *empirical* side of that comparison: a generalised stone-throwing problem where exact subset enumeration of actual causes scales as $(2^{2n} - 1)^2$ cause–witness pairs and is therefore the natural pressure test for the necessity-only PCI estimator. We report runtime, search-space size, and attribution-accuracy curves as the problem size grows.

9 Scaling Actual-Cause Computation: An Empirical Comparison

The correspondence in Theorems 28–29 and Proposition 34 is a statement about verdicts: under the stated hypotheses, AC actual causes coincide with the support of $\sum_{\mathbf{T}} \Phi$. This section reports an empirical companion to the correspondence. We ask whether the necessity-only PCI kernel of Definition 27, instantiated with a finite Monte Carlo estimator, recovers AC verdicts at problem sizes where exact subset enumeration is no longer tractable. The full implementation, raw outputs and plotting code are in the companion notebook `docs/source/actual_causality_benchmark.ipynb`.

Generalised throwing problem. For a size parameter n we replicate the canonical Sally/Billy structure n times in parallel. Binary suspects $A_1, \dots, A_n, B_1, \dots, B_n \in \{0, 1\}$ feed deterministic mediators $W_i^a = A_i$, $W_i^b = B_i \wedge \neg A_i$ that capture overdetermination via preemption: B_i contributes only if A_i does not. The outcome is the conjunction $Y = \bigwedge_{i=1}^n (W_i^a \vee W_i^b)$, and the model size grows as $4n+1$ variables. Ground-truth responsibility at site i is read off during the forward pass: A_i is responsible if $W_i^a = 1$, B_i is responsible if $W_i^b = 1$. The structure scales the canonical example without softening it — overdetermination, undercutting, and witness-mediated symmetry-breaking are present at every site — while keeping a closed-form ground truth available for evaluation.

The variable partition takes the roots $\{A_i, B_i\}_i$ as suspects and the deterministic mediators $\{W_i^a, W_i^b\}_i$ as witnesses, with no suspect a deterministic function of the others. This partition satisfies the closure condition required by Theorem 28 (see the Remark on deterministic intermediates and joint support, which addresses precisely the scenario in which both a root and its structural descendant are admitted as suspects). The benchmark is therefore within the scope of the formal correspondence; in particular, joint full support of the alternative-value distribution holds on the admissible suspect set $\{A_i, B_i\}_i$ even though the model itself is deterministic, because witnesses are not themselves part of the alternative-value distribution.

Exact subset enumeration (AC reference). The exact procedure mechanises Definition 26: for each non-empty cause subset $\mathbf{C} \subseteq \{A_i, B_i\}_i$ and each non-empty witness subset $\mathbf{T} \subseteq \{W_i^a, W_i^b\}_i$, we intervene $[\mathbf{C} \leftarrow 1 - \mathbf{c}^*, \mathbf{T} \leftarrow \mathbf{w}^*]$, record whether the outcome flips, and accept $\mathbf{C}$ as actual cause if some witness $\mathbf{T}$ achieves a flip and no proper subset of $\mathbf{C}$ does. The search space has $(2^{2n} - 1)^2$ cause-witness pairs (9, 225, 3969, 65025 for $n = 1, 2, 3, 4$), which already determines the ceiling reached by the exact method.

Approximate PCI estimator. The PCI estimator instantiates the kernel of Definition 27 via the Monte Carlo search procedure of Section 3. Algorithm 1 is the explicit procedure used to produce the curves in Figures 6–8.

The algorithm draws N intervention configurations under shared noise. The active suspect set $\mathbf{C}_j$ (line 10) is drawn at uniformly chosen cardinality $k_s \in \{1, \dots, n_s\}$; the dropped witness set D_j (line 11) is drawn the same way at $k_w \in \{0, \dots, n_w\}$, and the active witnesses on line 12 are the complement $\mathbf{W} \setminus D_j$, so n_w bounds the number of witnesses *dropped*, not the number kept, and the active witness set has size $|\mathbf{W}| - n_w$ or larger. The witness ablation flag `wit = off` is handled once on line 2 by emptying $\mathbf{W}$; this collapses the witness machinery on lines 11–12 and 14–15 to no-ops, producing the dotted accuracy curves of Figure 8.

Alternative values on line 13 are bit flips: on $\{0, 1\}$ the ϵ -excised alternative-value distribution has a single support point and ϵ plays no role; the explicit sampling form is preserved so that the same routine runs on continuous benchmarks. The two interventional worlds on lines 14–15 share the noise realisation $\mathbf{u}_j$ drawn on line 8, the standard counterfactual coupling, which lets the per-suspect score on lines 16–18 be read off as the nan-mean of $|y_j^n - y^*|$ over the draws in which $X \in \mathbf{C}_j$. The sufficiency outcome y_j^s is computed but not consumed: the AC verdict against which we benchmark has no sufficiency clause, so we read only the necessity world. The joint estimator with sufficiency active is exercised in Sections 5, 6, and 10, where AC has no comparable verdict to benchmark against. At each site $i = 1, \dots, n$, line 24 returns the suspect, A_i or B_i , with the higher mean.

Benchmark protocol. For each problem size n we sample up to ten distinct factive worlds (with $Y = 1$), instantiate the model, run both methods to verdicts, and compare to the analytic ground truth. Each method is granted a 60-minute global budget per run; if the budget is exhausted mid-loop, the largest completed n caps the curve. The approximate estimator is exercised in five configurations

Algorithm 1 Approximate PCI estimator on the scaled throwing problem.

Require: PSCM M ; factual world $(\mathbf{s}^*, \mathbf{w}^*, y^*)$; cardinality bounds n_s, n_w ; sample budget N ; witness flag $\text{wit} \in \{\text{on}, \text{off}\}$
Ensure: per-site verdicts $v_i \in \{A_i, B_i\}$ for $i = 1, \dots, n$

```
1: if  $\text{wit} = \text{off}$  then
2:    $\mathbf{W} \leftarrow \emptyset$  ▷ no witness candidates
3: end if
4: for each  $X \in \mathbf{S}$  do
5:    $L[X] \leftarrow []$  ▷ per-suspect score list
6: end for
7: for  $j = 1, \dots, N$  do
8:   draw  $\mathbf{u}_j \sim P_{\mathbf{U}}$ 
9:   sample  $k_s \sim \text{Unif}\{1, \dots, n_s\}$ ,  $k_w \sim \text{Unif}\{0, \dots, n_w\}$ 
10:  sample  $\mathbf{C}_j \subseteq \mathbf{S}$  uniformly with  $|\mathbf{C}_j| = k_s$ 
11:  sample  $D_j \subseteq \mathbf{W}$  uniformly with  $|D_j| = k_w$  ▷ witnesses dropped
12:   $\mathbf{T}_j \leftarrow \mathbf{W} \setminus D_j$  ▷ active witnesses
13:   $\mathbf{c}'_j \leftarrow \mathbf{1} - \mathbf{s}^*|_{\mathbf{C}_j}$  ▷ bit-flip on  $\{0, 1\}$ 
14:   $y_j^s \leftarrow M.\text{run}(\mathbf{u}_j, \text{do}(\mathbf{C}_j = \mathbf{s}^*|_{\mathbf{C}_j}, \mathbf{T}_j = \mathbf{w}^*|_{\mathbf{T}_j}))$  ▷ sufficiency
15:   $y_j^n \leftarrow M.\text{run}(\mathbf{u}_j, \text{do}(\mathbf{C}_j = \mathbf{c}'_j, \mathbf{T}_j = \mathbf{w}^*|_{\mathbf{T}_j}))$  ▷ necessity, shared  $\mathbf{u}_j$ 
16:  for each  $X \in \mathbf{C}_j$  do
17:    append  $|y_j^n - y^*|$  to  $L[X]$ 
18:  end for
19: end for
20: for each  $X \in \mathbf{S}$  do
21:    $\text{score}[X] \leftarrow \text{nanmean}(L[X])$ 
22: end for
23: for  $i = 1, \dots, n$  do
24:    $v_i \leftarrow \arg \max(\text{score}[A_i], \text{score}[B_i])$ 
25: end for
26: return  $(v_1, \dots, v_n)$ 
```

that vary along three axes identified in Section 3: (i) initial sample budget $N_0 \in \{250, 500\}$, scaled linearly with n by a factor $\alpha \in \{0.5, 1\}$ so that the total sample at size n is $\alpha \cdot N_0 \cdot n$; (ii) presence of witness interventions ($\text{wit} \in \{\text{on}, \text{off}\}$), ablating $\mathbf{T} \leftarrow \mathbf{w}^*$ to test whether witness pinning is load-bearing; (iii) cardinality bounds n_s, n_w , fixed at 4 in the default runs (answering “does some witness set of size $\geq |\mathbf{W}| - 4$ work?”) and grown linearly as $4n$ in the dynamic-set-size configuration (recovering the unrestricted subset search at the cost of higher variance). The cardinality bound is a tunable approximation knob, not a hidden assumption: the dynamic-set-size run measures the empirical cost of bounding n_s, n_w at 4 on this problem, and the gap between the two configurations is what the ablation surfaces. The conceptual case for constraining cardinality at all, that uniform sampling over the full powerset biases attributions toward large-coalition interventions and is analogous to declining to enumerate all n -way regression interaction terms, is given in Section 3.

Results. Three observations summarise the experiment. First, runtime (Figure 6): the exact procedure times out at $4n + 1 = 17$ variables ($n = 4$) once the cause-witness search space crosses sixty-five thousand pairs, while the approximate methods continue to roughly 73–145 variables depending on sample budget, with the frugal $\alpha=0.5$ run reaching the largest sizes. Second, search-space size (Figure 7, log scale): the exact method visits the full powerset, while the approximate methods visit at most a few hundred unique configurations per problem size: the gap is several orders of magnitude and widens with n . Third, accuracy (Figure 8): the exact method is correct by construction up to its early demise; the witness-using approximate methods stay above 0.85 correct-attribution rate up to roughly 60 variables and degrade gracefully beyond that, with the dynamic-set-size run sustaining among the highest accuracies at moderate problem sizes. Witness-free runs (dotted) underperform throughout, starting near 0.67 already at $n = 1$. The ablation confirms a structural claim of Section 3: pinning factual witness values is doing real work, not just regularising.




Figure 6: Per-world runtime against problem size. The exact method (orange) times out at 17 variables; approximate methods continue, with reach controlled by total sample budget. Five approximate configurations vary initial sample size, the linear-growth factor α , the witness ablation, and dynamic cardinality bounds; see Benchmark protocol.




Figure 7: Number of cause-witness configurations explored per problem size, log scale. The exact method visits the full powerset; the approximate methods visit a few hundred unique configurations per problem size regardless of n , a gap of several orders of magnitude that grows with n .




Figure 8: Correct-attribution rate against problem size. Each approximate configuration is lightly smoothed (Gaussian filter, $\sigma = 1$) with its raw per-size points overlaid, and is shape-coded in the legend (marker, colour, and line style). The exact method is correct up to 17 variables. Configurations using witnesses stay above 0.85 up to roughly 60 variables and degrade gracefully thereafter; the witness-free configurations (wit off) underperform throughout, starting near 0.67. Dynamic-set-size sampling (dyn on) is among the highest at moderate problem sizes.

Scope of the conclusion. The benchmark establishes that the necessity-only PCI estimator tracks AC verdicts on a faithful scaling of the canonical example with several orders of magnitude fewer evaluated configurations. The problem itself, however, is structured: responsibility attribution is independent across sites, exactly one of A_i, B_i is responsible at each site given $C=1$, supersets of any witness set witness the same attribution (so a small witness-cardinality bound costs little), and all variables are binary. These regularities account for some of the favourable scaling: in particular, the cardinality-bounded sampling losing little to dynamic-set-sizes sampling on this problem. Two extensions are natural follow-ups. First, a comparison against SHAP and causal SHAP on the same scaled-throwing problem is deferred to a companion benchmark (cf. Section 5 for the corresponding comparison on the continuous overdetermination example); the sufficiency component of PCI is what distinguishes it from those baselines, and a clean head-to-head requires the joint kernel rather than the necessity-only restriction used here. Second, benchmarks that break each of the structural regularities above, particularly continuous-variable extensions where AC’s mechanical enumeration is no longer even well-defined, would test the estimator outside its current favourable regime. As a sanity check that the correspondence in Theorems 28–29 is empirically operational at scale, the present comparison is sufficient.

Where we go next. The AC benchmark exercises the necessity-only specialisation of PCI on a discrete-outcome problem with a closed-form ground truth. Section 10 shifts to the regime where AC has no verdict to compare against: a Bayesian dynamical SIR model with a continuous outcome and asymmetrically interacting policies. The point is to exercise the joint necessity-sufficiency machinery on threshold events that the actual-causality machinery cannot adjudicate, and to show that PCI separates lockdown and masking where but-for analysis collapses them.

10 Dynamical SIR Benchmark: PCI on Continuous Outcomes

The actual-causality benchmark of Section 9 exercises the necessity-only specialisation of PCI on a discrete outcome with a closed-form ground truth. This section shifts to the regime where AC has no verdict to compare against: a Bayesian dynamical model with a continuous outcome and asymmetrically interacting causes. The benchmark mirrors a public ChiRho tutorial²³ in setup and query, which uses ChiRho’s `SearchForExplanation` handler to compute classical actual-cause probabilities. We keep the same model and the same query but replace the explanatory machinery with the PCI thin-search sampler. The full implementation, raw outputs and plotting code are in the companion notebook `docs/source/sir_benchmark.ipynb`.

Model. The dynamics are the standard SIR system $\dot{S} = -\beta SI$, $\dot{I} = \beta SI - \gamma I$, $\dot{R} = \gamma I$, with Bayesian priors $\beta \sim \text{Beta}(18, 600)$ and $\gamma \sim \text{Beta}(1600, 1600)$. Two non-pharmaceutical policies, each enacted with prior probability $1/2$, modulate the transmission rate via an intervention strength $l \in [0, 1]$ that scales β_0 to $(1-l)\beta_0$. Lockdown alone has efficiency 0.6; masking efficiency depends on lockdown (0.1 under lockdown, 0.45 on its own); the joint efficiency is the clamped sum, capped at 0.95. Lockdown is enacted at $t = 1$, masking at $t = 1.5$. The outcome is the *overshoot*, the peak-to-final S gap of the simulated trajectory, and the undesirable event is $\text{os_too_high} = \mathbb{1}[\text{overshoot} > 24]$ on a 100-person population.

But-for analysis fails to separate the suspects. Conditioning on each pair of policy decisions and drawing 100 predictives yields $\Pr(\text{overshoot} > 24) \approx 0.07$ with no intervention, 0.67 with both policies, but 0.82 with lockdown only and 0.87 with mask only — so, since a higher overshoot probability marks a *weaker* policy, mask-only is the weakest of the three intervention regimes. The mechanism is the asymmetric joint efficiency: combining the two policies caps the transmission reduction at 0.95, but neither policy alone reaches that cap, and mask-only is comparatively weak unless lockdown is absent (which shifts mask efficiency from 0.1 to 0.45). But-for cannot disentangle this context dependence: it treats each policy as a binary cause and reports an aggregate effect, with no way to express that lockdown’s causal contribution depends on whether masking is in force. Figure 9 visualises the four interventional regimes: the trajectory bands collapse toward the un-intervened baseline only under both policies, while either policy on its own leaves the infectious peak essentially intact, mirroring the near-equal but-for overshoot probabilities.

What a correct diagnosis should say. The but-for numbers are symmetric, but the model is not, and the asymmetry points to a clear qualitative reading that any responsibility method ought to recover. First, *lockdown is the proximate cause of excessive overshoot*: it is the policy that does the real work of suppressing the infectious peak, and the joint regime clears the 24-overshoot threshold only because lockdown is in force. Second, *masking is at most a context-sensitive cause*: its contribution depends on whether lockdown is also active. Masking is comparatively effective only when lockdown is absent (efficiency 0.45 versus 0.1 under lockdown), so once lockdown is present masking adds little. A good attribution method should therefore rank lockdown above mask and, more finely, flag mask’s causal status as contingent on context rather than robust. These two intuitions are the target of the analysis below; they follow from the model’s asymmetric joint efficiency, not from any particular tool, and they coincide with the verdict the ChiRho tutorial reaches via classical actual-cause probabilities. We state them here so the comparison that follows does not depend on consulting that tutorial.

PCI thin-search procedure. The suspects are $\{\text{lockdown}, \text{mask}\}$. The witnesses are drawn from the policy efficiencies $\{\text{lockdown_efficiency}, \text{mask_efficiency}, \text{joint_efficiency}\}$, with the suspect itself excluded from the witness set on each draw. The factual world fixes both policies on ($\text{lockdown} = 1, \text{mask} = 1$), so the factual outcome $y^* = \text{overshoot}$ is read from the same trajectory. For each suspect the sampler draws 2,500 regimes, scoring each with the $|y^* - y_{\text{nec}}| - |y^* - y_{\text{suff}}|$ statistic and averaging across regimes. The *nec* term rewards regimes where removing the suspect moves the outcome *away* from y^* ; the *suff* term rewards regimes where fixing the suspect at factual keeps the outcome *close* to y^* .

The joint necessity-sufficiency plane. Before turning to numbers, it helps to see what the search produces. For each suspect the sampler returns, per regime, a pair of outcomes: the *necessity*-

²³https://basisresearch.github.io/chirho/explainable_sir.html




Figure 9: SIR trajectories and overshoot distributions under the four interventional regimes (no interventions, both policies, lockdown only, mask only). Left: posterior bands for susceptible (blue), infectious (red) and recovered (green) over time, with the lockdown enacted at $t = 1$ and the mask mandate at $t = 1.5$. Right: per-regime histograms of the overshoot, with the 24-overshoot threshold marked. Either policy on its own leaves the infectious peak essentially intact; only the joint regime suppresses it appreciably, yet the lockdown-only and mask-only overshoot probabilities are near-indistinguishable: the asymmetric joint efficiency that PCI disentangles.

world overshoot y_{nec} (the focal suspect set to its alternative, any other active suspects in $\mathbf{C} \setminus \{X\}$ intervened, witnesses pinned at factual) and the *sufficiency-world* overshoot y_{suff} (the focal suspect held at factual, other active suspects intervened, witnesses at factual). These are the necessity- and sufficiency-world outcomes y^n, y^s of Section 3; we write $y_{\text{nec}}, y_{\text{suff}}$ here to match the figure axes. Plotting one point per regime in the $(y_{\text{nec}}, y_{\text{suff}})$ plane gives a per-suspect cloud (Figure 10) that exposes the score geometry before any averaging. Three reference lines orient the plane: the dotted red cross-hair marks the factual outcome y^* at the both-policies-on world (lockdown = mask = 1); because that world is the high-overshoot scenario, y^* sits near the upper-right of the cloud rather than at its centre. The dash-dot blue line marks the 24-overshoot threshold, and the dashed black diagonal is $y_{\text{nec}} = y_{\text{suff}}$. The per-regime score $|y^* - y_{\text{nec}}| - |y^* - y_{\text{suff}}|$ is positive exactly when a point lies *further from the cross-hair along the necessity axis than along the sufficiency axis*: necessity pulls the outcome away from factual (evidence the suspect was needed), while sufficiency keeps it close (evidence the suspect alone sustains the outcome). A cause therefore shows up as a cloud displaced toward smaller y_{nec} at fixed y_{suff} — the *cause signature* on this geometry — whereas a non-cause hugs the diagonal. Reading the two panels this way, the lockdown cloud is clearly displaced below the diagonal toward smaller y_{nec} , while the mask cloud sits closer to it: the visual form of the verdict the next paragraph quantifies.

Single-world results. PCI assigns lockdown roughly $2.0\times$ the responsibility of mask:




Figure 10: Joint necessity-sufficiency overshoot density per suspect. Each point is one sampled regime. Dotted red cross-hair: factual y^* at lockdown = mask = 1 (the high-overshoot anchor); dash-dot blue: the 24-overshoot threshold; dashed black: $y_{nec} = y_{suff}$. A regime contributes positively to the PCI total iff it sits further from the red cross-hair along the necessity axis than along the sufficiency axis. The lockdown density is shifted toward smaller y_{nec} relative to y_{suff} (the cause signature on this geometry); the mask density sits closer to the diagonal.

suspect	$ y^* - y_{nec} $	$- y^* - y_{suff} $	total
lockdown	8.12	-6.58	1.54
mask	7.43	-6.67	0.76

The gap is driven entirely by the necessity term: removing lockdown moves the overshoot further from y^* than removing mask does (8.12 vs 7.43 in mean absolute distance), while the sufficiency terms agree to within 0.1. This is consistent with the but-for finding that either policy *alone* is roughly equally bad. What distinguishes the suspects is that lockdown’s removal *under context* (with witnesses pinned to factual) pulls the world further from y^* than mask’s removal does — the lockdown cloud’s displacement below the diagonal toward smaller y_{nec} in Figure 10, against the mask cloud’s near-diagonal position.

Robustness across factual worlds. We replicate the analysis across 10 factual worlds drawn from the prior (rather than the single hand-set world above), running thin search on each and averaging. Lockdown’s mean total exceeds mask’s in the majority of worlds; the verdict is not an artifact of the chosen factual.

Effect of context. The ChiRho tutorial isolates one downstream variable at a time, asking whether the verdict survives when that variable is held fixed at factual versus left free. We replicate the experiment by partitioning the PCI regimes by whether the partner-policy efficiency variable was sampled into the witness set (Figure 11). For lockdown, with mask_efficiency *fixed* as a witness the score is 0.81; *free*, it rises to 2.67. For mask, with lockdown_efficiency *fixed* the score is -0.48 (mask fails to register as a cause at all under that context regime) and *free* it rises to 2.72. This experiment partitions the regimes differently from the single hand-set world reported earlier, so the fixed/free scores are not expected to average to that world’s totals (lockdown 1.54, mask 0.76). Lockdown therefore registers as a cause in both context regimes, while mask’s status as a cause is contingent on its partner being allowed to adapt. PCI thus expresses not just *which* policy is the dominant cause but *how robustly* that causal claim holds across counterfactual contexts: a finer-grained verdict than the binary “actual cause / not” that classical AC produces.

Takeaway. On the dynamical SIR setting, PCI recovers both intuitions set out at the start of this section — lockdown is the proximate cause of excessive overshoot and masking only a context-sensitive one, despite both policies appearing similar under but-for analysis — and expresses them on a continuous, decomposable scale. The necessity-vs-sufficiency decomposition distinguishes suspects whose isolation effects are similar but whose removal-under-context effects diverge; the

Effect Of Fixing The Partner-Policy Efficiency As Witness




Figure 11: Different-contexts experiment. Bars compare PCI total score (left) and $\Pr(\text{overshoot} > 24 \mid \text{necessity})$ (right) when the partner-policy efficiency variable is fixed as a witness versus left free. Lockdown’s score is positive in both regimes; mask’s score flips sign across regimes, indicating that mask is a context-sensitive rather than a robust cause.

contexts experiment further distinguishes robust from context-sensitive causes. The same machinery applies to any ChiRho probabilistic program with a continuous outcome and a tractable enumeration over interventional regimes.

Where we go next. The SIR benchmark scales PCI to a research-grade Bayesian dynamical model and demonstrates that the joint kernel captures patterns that AC cannot adjudicate. Section 11 pushes the demonstration one step further: a deployed automated valuation model trained on millions of points, with deep neural-network components and a sparse variational GP outcome head. The question there is no longer whether PCI agrees with a known ground truth (on a proprietary production model none is available), but whether the computation is feasible at all, and whether the qualitative SHAP-vs-PCI divergences observed on the synthetic side persist on a real trained system.

11 Scaling PCI to a Real-World Automated Valuation Model

Where we are. The synthetic archetypes of Section 5, the AC benchmark of Section 9, and the SIR benchmark of Section 10 have all controlled the model and the ground truth, so that PCI could be judged against either an analytic verdict or a competing AC enumeration. This section drops both: we apply PCI to a deployed production model on which neither ground truth nor exact AC enumeration is available, and ask whether the computation remains feasible and whether the qualitative SHAP-vs-PCI divergence persists.

In this section, we apply PCI to a valuation model for residential housing. Disclosure constraints on the trained model and its training data prevent us from publishing a quantitative accuracy comparison against ground truth on this system; the quantitative validation of PCI therefore lives entirely in the controlled synthetic benchmark of Section 5, against an analytic ground truth that the production model does not admit. The role of this section is correspondingly narrow: to report (i) that PCI is computationally feasible at production scale on a model with millions of training points, and (ii) that the qualitative SHAP-vs-PCI divergence observed in Section 4 persists when the model is a real trained one. What is reportable here is the shape of the two attribution distributions and the structural pattern of where attribution mass lands, not the substantive content of the features they assign to.

Setting. The unit of analysis is a residential property transaction. PCI is run against the full country-level production AVM; what we show in this section, however, is an illustrative prototype based on the state of New York rather than the full national model structure, which we cannot disclose in detail. The NY prototype shares the architecture and the causal-kernel construction described below with the full system but is smaller in scope and DAG complexity. The computational-feasibility claim refers to the full national system; the qualitative SHAP-vs-PCI behaviour we illustrate is a property of the shared architecture and carries over from the prototype. The outcome

variable y is the (log-epsilon-standardized) sale price. Transactions are described by a set of categorical variables $x_{cat} = x_1, \dots, x_k$ and continuous (transformed) variables²⁴ $x_{con} = x_{k+1}, \dots, x_n$.

Causal model. A causal DAG $\mathcal{G}$ expresses our causal assumptions about the relationships between variables. It was written manually in collaboration with domain experts and refined by adding edges to break implied conditional independencies that are not upheld by the data. The NY-prototype DAG has 25 nodes and 99 edges; topology with anonymised node labels is shown in Figure 12; the full national DAG is larger and is not shown. As demonstrated in Section 9, even the prototype is already large enough that exact actual-causality enumeration is computationally infeasible, and the same conclusion applies a fortiori to the national system.

Variables x_i are drawn from distributions Dist_i parameterized by functions $f_i(\text{pa}_i, \varphi_i)$ of the parent variables pa_i ²⁵: $x_i \sim \text{Dist}_i(f_i(\text{pa}_i, \varphi_i))$. We call these distributions *causal kernels*. The distributions are either (truncated) normal or categorical, chosen based on the domain of the variable that they model. The functions f_i are deep neural networks with parameters φ_i . The parameters φ_i are optimized using cross-entropy loss for categorical distributions and average negative log likelihood for continuous distributions. The outcome is modeled by a sparse variational Gaussian Process (SVGP; Titsias, 2009, Hensman et al., 2013), $x_{gp} = (x_{con}, \eta(x_{cat}, \varphi_{cat}))$, $y \sim \text{SVGP}(\mu(x_{gp}, \varphi_\mu), \kappa(x_{gp}, \varphi_\kappa))$, where η is a linear embedding of the categorical variables into the continuous space, jointly optimized with the GP.

Feasibility at production scale. To apply PCI we transform the model as described in Section 3 and evaluate the impact of features from a list of potential causal suspects, using the absolute difference impact score of Example 21. We draw Monte Carlo samples from the expanded model. Conditioning on a given suspect X_i being active gives us a sample of the scores corresponding to that feature, which we use to estimate the expected value thereof. The estimation converges at around 25,000 samples; for a batch of 50 data points this takes around 60 minutes on commodity hardware. Sampling needs to be performed only once; the samples can then be reused to compute the attribution score of any suspect. This establishes that the method is computationally tractable at production scale, not a foregone conclusion, since the expanded-model construction in Section 3 introduces additional latent dimensions per suspect and per witness.

Qualitative comparison to SHAP. We compare PCI attributions to SHAP scores [Lundberg and Lee, 2017] on the same trained model. SHAP heavily weights variables that are downstream of many other variables in the causal graph, effectively assigning attribution to what amount to summary statistics of the outcome. PCI distributes attribution more evenly across the upstream causal structure. The pattern is consistent with what Section 4 establishes on controlled examples: SHAP’s observational marginalisation allocates weight differently from intervention-and-witness-based PCI scores, and on a deep DAG this difference manifests as concentration on summary-statistic-like nodes. Figure 13 shows the two attribution distributions with anonymised feature labels; what is reportable here is the shape of the two distributions, not which features they assign to.

Where we go next. Section 12 closes the paper by articulating the relationship between PCI and the two traditions it inherits from, Halpern’s actual causality and Pearl’s PN/PS/PNS, against the background of what every preceding section has shown, and points out the limitations of the present construction along with the natural follow-up directions.

12 Discussion and Conclusions

We are now in a position to articulate the main differences between PCI and the two approaches that inspired it, against the background of what the rest of the paper has shown. The main lesson from the literature on actual causality is that intuitive causal attribution often requires holding some endogenous variables, members of a “witness set”, fixed at their factual values. This ensures context sensitivity by controlling complicating causal paths. We treat witnesses somewhat differently: instead of looking for a single witness set that reveals a causal relationship, we use a distribution

²⁴Usually log-epsilon-standardized, standardized-truncated, or standardized.

²⁵For root nodes, pa_i is empty.




Figure 12: Topology of the AVM causal DAG (NY-prototype version), with node labels anonymised. The graph has 25 nodes and 99 edges; the full national DAG is larger and is not shown. Several variables sit deep in the DAG with high in-degree; these are the nodes towards which SHAP allocates disproportionate attribution mass (see Figure 13).




Figure 13: Attribution distributions on the AVM, with feature labels anonymised. **Left:** PCI attribution mass spread across upstream causal variables (suspects 1–15, ranked by mean total score). **Right:** SHAP attribution mass concentrated on a small number of downstream variables. The qualitative pattern matches the controlled-setting comparison of Section 4.

over witness sets and integrate the causal impact score across them. As in many probabilistic generalisations, this focuses on distributions rather than point estimates, and it lets PCI account for the number of witness sets and the strength of within-witness-set effects. Moreover, the construction admits off-the-shelf probabilistic inference rather than worst-case subset enumeration. The relationship between the two views is now formal rather than gestural: under matching choices of Γ and Δ , AC verdicts coincide with the support of $\sum_{\mathbf{T}} \Phi$ (Theorems 28–29 and Proposition 34), and the empirical companion in Section 9 confirms this at scales where AC enumeration is intractable.

The main concept borrowed from Pearl’s probability of necessity and sufficiency is the joint distribution over outcomes from two interventional regimes, pushed forward through a user-chosen causal impact function. The key difference is that we use witness sets to allow for context-sensitivity and to break symmetries present in otherwise pathological cases like overdetermination and undercutting. Section 4 works through the binary instantiation in which the necessity and sufficiency components coincide with PN and PS, and Section 5 reads the two components apart on a continuous synthetic model where each is needed to surface a different causal archetype.

The further generalisations are mostly bookkeeping in the same spirit. Instead of fixing a single alternative value for a potential cause, we integrate over a factual-excised alternative-value distribution; this generalises immediately to continuous variables. Instead of asking whether a particular set is an actual cause, we sample suspect sets and gauge a feature’s responsibility in terms of its participation in sets with causal impact. Instead of asking whether the outcome differs, a customisable causal impact function ci measures the differences across counterfactual worlds, including between continuous variables. The variable-selection distribution Γ , the alternative-value distribution Δ , and the impact function ci become explicit, user-facing components of the framework rather than implicit defaults.

Against SHAP-family attributions, the framework’s central commitment is to *intervention with witnesses* rather than *observational marginalisation*. The controlled comparisons of Section 4 show that this commitment is load-bearing on canonical overdetermination and continuous-mediation examples, where SHAP and Causal SHAP fail in characteristic ways that PCI does not. The synthetic benchmark of Section 5 confirms this on a model with known ground truth, and the case study of Section 11 reports the same qualitative pattern on a production-grade trained model with millions of data points. A separate comparison to gradient-based attribution, where the contrast is differentiability and locality rather than the causal/non-causal axis, is reported in Section 6. The thread connecting these results is structural: the witness mechanism that satisfies D-A-rank for Alice in

the binary OBCB walkthrough of Section 4 is the same machinery driving the upstream-versus-downstream attribution split on the AVM in Section 11.

A broader test is how PCI fares against the desiderata that the XAI literature typically asks counterfactual-explanation methods to meet, of which Verma et al. [2024] list five. *Sparsity*, the requirement that explanations change as few features as possible, follows directly from the cardinality cap on suspect sets in Γ , which biases the search toward small C . *Multiple counterfactuals* (the ability to return several alternative explanations rather than commit to one) falls out of the Monte Carlo construction, since a single sweep over Γ and Δ produces a population of (C, c', T) triples rather than a single optimum. *Causal-constraint satisfaction*, the demand that explanations respect downstream propagation rather than treat features as independently editable, is automatic once suspect interventions are routed through the causal model: changing C propagates to descendants by construction. *Model-agnosticism* (applicability to arbitrary predictors) PCI satisfies only partially: it can wrap a black-box predictor, but at the cost of the causal sensitivity that motivates the framework in the first place.

Actionability, the requirement that explanations not recommend changes to features users cannot in practice change, is where we part company with the literature. The standard recommendation is to fold ease-of-change into the cause-identification procedure itself, weighting features by mutability so that immutable ones are never returned as causes (Verma et al. [2024]; cf. Karimi et al. [2021], who shift the focus from explanations altogether to interventions). We think this conflates two distinct questions. A procedure that suppresses gender as a possible cause whenever a loan decision is at stake will help applicants game the existing system, but it will also prevent them from noticing the kind of structural problem that responsibility attribution can in principle surface. Cost-of-action layers can sit downstream of PCI for users who need actionable recommendations, but they should compose with responsibility attribution rather than replace it.

Several limitations are worth flagging. PCI presumes access to a trained probabilistic causal model from which counterfactual outcomes can be sampled, a stronger requirement than methods that work from observational data and a causal graph alone. The Monte Carlo variance of the estimator grows with model size and with the cardinalities of suspect and witness sets; Section 9 reports empirical scaling and an ablation that isolates the contribution of witness pinning, but sample-efficient estimators for high-cardinality settings remain an open direction. Finally, the AVM case study is deliberately limited in disclosure: the apples-to-apples SHAP-vs-PCI comparison rests on the synthetic benchmark of Section 5, with the production model serving as a feasibility witness rather than as quantitative evidence.

Where PCI falls short. Three caveats about PCI’s reach deserve explicit discussion. (i) *AC mismatch under noise integration.* Observation 33 shows that integrating the kernel against P_U before maximisation can drop AC causes that are subset-minimal but not cardinality-minimal: the Γ -weighted ranking is governed by cardinality, while AC’s minimality is governed by the subset relation. Proposition 34 provides the operational route to AC verdicts in such cases; Theorem 31 closes the gap only under *necessity dilution* (Definition 30), a regularity that fails in disjunctive overdetermination. (ii) *Cardinality knob versus sample efficiency.* The cardinality cap is a tunable hyperparameter rather than a hidden default, but it trades precision for variance: tight bounds shrink the search space and reduce Monte Carlo variance per iteration but exclude joint-cause sets that AC’s exhaustive enumeration would surface; loose bounds restore generality at the cost of slower convergence. Problem families with diffuse joint causation need more generous settings than the canonical AC benchmark uses, and the cap is therefore worth reporting alongside results. (iii) *Cost of the PSCM-availability assumption.* Pointwise verdicts at the configuration level require a sampleable structural model. Methods that bound PN/PS/PNS from observational data and a causal graph alone, e.g. Tian and Pearl’s bounds, apply in settings where PCI cannot, and the stronger PSCM commitment should be weighed against this when only a graph and a dataset are available.

Taken together, PCI provides a scalable, context-sensitive, causal explanation method for causal probabilistic machine-learning models. A natural probabilistic generalisation of ideas already in the literature leads to a tractable estimator that can be paired with standard inference algorithms. We benchmarked the approach against exact actual-causality computations, characterised its qualitative behaviour on synthetic models with known ground truth and on canonical SHAP failure cases, and showed its computational feasibility on a production-grade model.

Responsibility attribution has not yet been effectively mechanised in software for large, complicated models, which makes the exact notions present in the literature infeasible to apply. PCI develops approximate causal attribution workable for users of models of non-trivial size, with the design choices — which variables are suspects, what counts as an alternative value, what counts as a difference — exposed to the user rather than baked in.

References

- Sander Beckers and Joost Vennekens. Combining probabilistic, causal, and normative reasoning in CP-logic. *ArXiv*, abs/1503.01051, 2015.
- K. Butler, G. Feng, and P. M. Djurić. A differential measure of strength of causation. *IEEE Signal Processing Letters*, 29, 2022.
- Paul Cheshire and Stefano Magrini. Growing urban gdp or attracting people? different causes, different consequences. In Charlie Karlsson, Åke E. Andersson, Paul C. Cheshire, and Roger R. Stough, editors, *New Directions in Regional Economic Development*, Advances in Spatial Science, pages 291–315. Springer, Berlin, Heidelberg, 2009. ISBN 978-3-642-01016-3, 978-3-642-01017-0. doi: 10.1007/978-3-642-01017-0_16.
- H. Chockler and J. Y. Halpern. Responsibility and Blame: A Structural-Model Approach. *Journal of Artificial Intelligence Research*, 22:93–115, October 2004. ISSN 1076-9757. doi: 10.1613/jair.1391.
- Jacob Cohen. *Statistical Power Analysis for the Behavioral Sciences*. Lawrence Erlbaum Associates, Hillsdale, NJ, 2nd edition, 1988.
- Thomas Eiter and Thomas Lukasiewicz. Complexity results for structure-based causality. *Artificial Intelligence*, 142(1):53–89, November 2002. ISSN 00043702. doi: 10.1016/S0004-3702(02)00271-0.
- C. W. J. Granger. Investigating causal relations by econometric models and cross-spectral methods. *Econometrica*, 37(3):424–438, 1969. doi: 10.2307/1912791. URL <https://www.jstor.org/stable/1912791>.
- Joseph Y. Halpern. *Actual Causality*. The MIT Press, 2016. ISBN 9780262035026. URL <http://www.jstor.org/stable/j.ctt1f5g5p9>.
- Joseph Y. Halpern and Christopher Hitchcock. Graded causation and defaults. *British Journal for the Philosophy of Science*, 66(2):413–457, 2015. doi: 10.1093/bjps/axt050.
- James J. Heckman and Rodrigo Pinto. The econometric model for causal policy analysis. *Annual Review of Economics*, 14(1):893–923, 2022. doi: 10.1146/annurev-economics-051520-015456. URL <https://doi.org/10.1146/annurev-economics-051520-015456>.
- James Hensman, Nicolò Fusi, and Neil D Lawrence. Gaussian processes for big data. In *Proceedings of the Twenty-Ninth Conference on Uncertainty in Artificial Intelligence*, pages 282–290, 2013.
- Tom Heskes, Evi Sijben, Ioan Gabriel Bucur, and Tom Claassen. Causal Shapley values: Exploiting causal knowledge to explain individual predictions of complex models. In *Advances in Neural Information Processing Systems*, volume 33, pages 4778–4789, 2020.
- Thomas F. Icard, Jonathan F. Kominsky, and Joshua Knobe. Normality and actual causal strength. *Cognition*, 161:80–93, April 2017. ISSN 00100277. doi: 10.1016/j.cognition.2017.01.010.
- Dominik Janzing, Lenon Minorics, and Patrick Blöbaum. Feature relevance quantification in explainable AI: A causal problem. In *International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2020.
- Amir-Hossein Karimi, Bernhard Schölkopf, and Isabel Valera. Algorithmic recourse: from counterfactual explanations to interventions. In *Proceedings of the 2021 ACM Conference on Fairness, Accountability, and Transparency, FAccT ’21*, page 353–362. ACM, March 2021. doi: 10.1145/3442188.3445899. URL <http://dx.doi.org/10.1145/3442188.3445899>.

- Yuta Kawakami, Manabu Kuroki, and Jin Tian. Probabilities of Causation for Continuous and Vector Variables. In Negar Kiyavash and Joris M. Mooij, editors, *Proceedings of the Fortieth Conference on Uncertainty in Artificial Intelligence*, volume 244 of *Proceedings of Machine Learning Research*, pages 1901–1921. PMLR, July 2024.
- John K. Kruschke. Rejecting or accepting parameter values in Bayesian estimation. *Advances in Methods and Practices in Psychological Science*, 1(2):270–280, 2018. doi: 10.1177/2515245918771304.
- Scott M Lundberg and Su-In Lee. A unified approach to interpreting model predictions. In *Advances in Neural Information Processing Systems*, volume 30, pages 4766–4777, 2017.
- Robert Osazuwa Ness. *Causal AI*. Simon and Schuster, 2025.
- Konstantinos Papageorgiou, Theodosios Theodosiou, Aikaterini Rapti, Elpiniki I. Papageorgiou, Nikolaos Dimitriou, Dimitrios Tzovaras, and George Margetis. A systematic review on machine learning methods for root cause analysis towards zero-defect manufacturing. *Frontiers in Manufacturing Technology*, Volume 2 - 2022, 2022. ISSN 2813-0359. doi: 10.3389/fmtec.2022.972712. URL <https://www.frontiersin.org/journals/manufacturing-technology/articles/10.3389/fmtec.2022.972712>.
- Judea Pearl. *Causality: Models, Reasoning and Inference*. Cambridge University Press, USA, 2nd edition, 2009. ISBN 052189560X.
- Judea Pearl. Probabilities of causation: three counterfactual interpretations and their identification. In *Probabilistic and Causal Inference: The Works of Judea Pearl*, pages 317–372. 2022.
- Lloyd S. Shapley. A value for n -person games. In Harold W. Kuhn and Albert W. Tucker, editors, *Contributions to the Theory of Games, Volume II*, volume 28 of *Annals of Mathematics Studies*, pages 307–317. Princeton University Press, Princeton, 1953.
- Amit Sharma, Hua Li, and Jian Jiao. The counterfactual-Shapley value: Attributing change in system metrics. *arXiv preprint arXiv:2208.08399*, 2022.
- Daisy A. Shepherd, David J. Amor, and Margarita Moreno-Betancur. Statistical analysis of observational studies in disability research. *Developmental Medicine & Child Neurology*, 66(11): 1408–1418, 2024. doi: 10.1111/dmcn.15948.
- Michalis Titsias. Variational learning of inducing variables in sparse Gaussian processes. In *Artificial Intelligence and Statistics*, pages 567–574. PMLR, 2009.
- Sahil Verma, Varich Boonsanong, Minh Hoang, Keegan Hines, John Dickerson, and Chirag Shah. Counterfactual explanations and algorithmic recourses for machine learning: A review. *ACM Computing Surveys*, 56(12):1–42, 2024.

A Computations for the OBCB Running Example

This appendix collects the worked computations for the stochastic OBCB model of Example 6 that are summarised in the main text: the per-branch traces behind Alice’s individual PN and PS values in Table 1 (Section 2), and the closed-form SHAP value function behind Tables 4–5 (Section 4.2). The remaining individual and population entries of Table 1 are obtained analogously; all values are reproduced in `docs/source/obcb_computations.ipynb`.

A.1 Alice’s probability of necessity

Alice is factually $gender = \text{female}$, $credit = \text{bad}$, $loan = F$; the counterfactual intervenes to set $gender = \text{male}$ while leaving $credit = \text{bad}$. Tracing through the model:

- Under the intervention, $check \sim \text{Bern}(0.9)$: men are checked 90% of the time.
- If checked, $check\text{-}failed = 1$ (Alice’s credit is bad), and from the loan-probability table $P(loan\text{-}if\text{-}checked = 1 \mid gender = \text{male}, check\text{-}failed = 1) = 0.05$.

- If not checked, $loan = 0$ deterministically since $loan = loan\text{-if-checked} \cdot check$.

Combining the two branches:

$$\begin{aligned}
PN(gender = \text{female}, loan = F \mid \text{Alice}) \\
&= P(loan_{gender=\text{male}} = T \mid gender = \text{female}, credit = \text{bad}) \\
&= 0.9 \cdot 0.05 = 0.045.
\end{aligned}$$

A.2 Alice’s probability of sufficiency

Following the same convention as for PN , the individual-level computation additionally conditions on Alice’s credit ($credit = \text{bad}$). The PS conditional then has Alice’s credit fixed but the cause and outcome flipped to their non-factual values ($gender = \text{male}$, $loan = T$); we intervene to restore $gender = \text{female}$ and ask whether the loan is again denied:

- With $gender = \text{female}$ and $credit = \text{bad}$, $check \sim \text{Bern}(0.2)$.
- If checked, $check\text{-failed} = 1$, and from the loan-probability table $P(loan\text{-if-checked} = 1 \mid gender = \text{female}, check\text{-failed} = 1) = 0$.
- If not checked, $loan = 0$ deterministically.

Either branch gives $loan = F$, so

$$\begin{aligned}
PS(gender = \text{female}, loan = F \mid \text{Alice}) \\
&= P(loan_{gender=\text{female}} = F \mid gender = \text{male}, loan = T, credit = \text{bad}) = 1.
\end{aligned}$$

A.3 SHAP value function for Option A

For the 2-feature game $N = \{gender, credit\}$ with $f(g, c) = P(loan = 1 \mid g, c)$, by the SHAP definition (and equivalently by Definition 23 once we note that no dropped feature is a descendant of any coalition member), the value function is

$$v(\mathcal{S}) = \mathbb{E}[f(X) \mid X_{\mathcal{S}} = x_{\mathcal{S}}^*] = \sum_{x_{N \setminus \mathcal{S}} \in \{0,1\}^{|N \setminus \mathcal{S}|}} P(X_{N \setminus \mathcal{S}} = x_{N \setminus \mathcal{S}}) f(x_{\mathcal{S}}^*, x_{N \setminus \mathcal{S}}).$$

Under independent uniform marginals $P(gender) = P(credit) = \frac{1}{2}$ this specialises to

$$\begin{aligned}
v(\emptyset) &= \frac{1}{4} \sum_{g,c \in \{0,1\}} f(g, c), & [\text{both dropped: weight } \frac{1}{2} \cdot \frac{1}{2}] \\
v(\{gender\}) &= \frac{1}{2}[f(g^*, 0) + f(g^*, 1)], & [\text{only credit dropped: weight } P(c) = \frac{1}{2}] \\
v(\{credit\}) &= \frac{1}{2}[f(0, c^*) + f(1, c^*)], & [\text{only gender dropped: weight } P(g) = \frac{1}{2}] \\
v(\{gender, credit\}) &= f(g^*, c^*). & [\text{nothing dropped: no average}]
\end{aligned}$$

Plugging in the four values $f(F, \text{bad}) = 0.000$, $f(F, \text{good}) = 0.180$, $f(M, \text{bad}) = 0.045$, $f(M, \text{good}) = 0.900$ yields the coalition values of Table 4; pushing those through the closed-form Shapley $\phi_i = \frac{1}{2}[v(\{i\}) - v(\emptyset)] + \frac{1}{2}[v(N) - v(N \setminus \{i\})]$ (and flipping signs, since we attribute rejection $g = 1 - f$) gives the magnitudes in Table 5.

B Computations for the Signal-with-Mediation Example

This appendix collects the closed-form derivations for the signal-with-mediation example of Section 4.3: the Gaussian chain $X \rightarrow M \rightarrow Y$ of Eqs. (9)–(11), with $X \sim \mathcal{N}(0.5, 0.25)$, $M = X + \varepsilon_M$, $Y = M + \varepsilon_Y$, and independent noise $\varepsilon_M, \varepsilon_Y \sim \mathcal{N}(0, 0.1)$. The distributional facts and conditional expectations stated in Example 24, and the per-target SHAP and per-pair PCI computations summarised in Table 7, are derived here. Unless noted otherwise the factual instance is Instance 1, $X^* = M^* = Y^* = 1$.

B.1 Distributional facts

From the structural equations, $\text{Var}(M) = \text{Var}(X) + \text{Var}(\varepsilon_M) = 0.25 + 0.1 = 0.35$ and $\text{Var}(Y) = \text{Var}(M) + \text{Var}(\varepsilon_Y) = 0.35 + 0.1 = 0.45$. For the covariances, $\text{Cov}(X, M) = \text{Cov}(X, X + \varepsilon_M) = \text{Var}(X) = 0.25$; $\text{Cov}(X, Y) = \text{Cov}(X, M + \varepsilon_Y) = \text{Cov}(X, M) = 0.25$; and $\text{Cov}(M, Y) = \text{Cov}(M, M + \varepsilon_Y) = \text{Var}(M) = 0.35$.

B.2 Conditional expectations

$\mathbb{E}[Y \mid X, M] = M$ since $Y = M + \varepsilon_Y$ with $\varepsilon_Y \perp (X, M)$. For $\mathbb{E}[M \mid X, Y]$, the covariance matrix of (X, Y) is $\Sigma_{XY} = \begin{pmatrix} 0.25 & 0.25 \\ 0.25 & 0.45 \end{pmatrix}$ with $\det(\Sigma_{XY}) = 0.05$, so $\Sigma_{XY}^{-1} = 20 \begin{pmatrix} 0.45 & -0.25 \\ -0.25 & 0.25 \end{pmatrix}$; the regression coefficients are $[\text{Cov}(M, X), \text{Cov}(M, Y)] \Sigma_{XY}^{-1} = [0.25, 0.35] \cdot 20 \begin{pmatrix} 0.45 & -0.25 \\ -0.25 & 0.25 \end{pmatrix} = [0.5, 0.5]$, giving $\mathbb{E}[M \mid X, Y] = 0.5 + 0.5(X - 0.5) + 0.5(Y - 0.5) = 0.5X + 0.5Y$. For $\mathbb{E}[X \mid M, Y]$, by d -separation, $X \perp Y \mid M$ in the chain $X \rightarrow M \rightarrow Y$, so $\mathbb{E}[X \mid M, Y] = \mathbb{E}[X \mid M] = 0.5 + \frac{\text{Cov}(X, M)}{\text{Var}(M)}(M - 0.5) = 0.5 + \frac{0.25}{0.35}(M - 0.5) = 0.5 + \frac{5}{7}(M - 0.5)$.

B.3 Plain SHAP at Instance 1

Target Y , features $\{X, M\}$. First, compute the coalition values:

$$v(\emptyset) = \mathbb{E}[M] = 0.5 \quad (22)$$

$$v(\{X\}) = \mathbb{E}[M \mid X = 1] = 1 \quad (23)$$

$$v(\{M\}) = \mathbb{E}[M \mid M = 1] = 1 \quad (24)$$

$$v(\{X, M\}) = f_Y(1, 1) = 1 \quad (25)$$

Next, apply equation (5) with $|N| = 2$:

$$\phi_X = \frac{1}{2}(1 - 0.5) + \frac{1}{2}(1 - 1) = 0.25 \quad \phi_M = \frac{1}{2}(1 - 0.5) + \frac{1}{2}(1 - 1) = 0.25 \quad (26)$$

Target M , features $\{X, Y\}$. $v(\emptyset) = 0.5$; $v(\{X\}) = 1$; $v(\{Y\}) = 0.5 \mathbb{E}[X \mid Y=1] + 0.5 = 0.889$ (using $\mathbb{E}[X \mid Y=1] = 0.5 + \frac{0.25}{0.45}(0.5) = 0.778$); $v(\{X, Y\}) = 1$.

$$\phi_X = \frac{1}{2}(1 - 0.5) + \frac{1}{2}(1 - 0.889) = 0.306, \quad (27)$$

$$\phi_Y = \frac{1}{2}(0.889 - 0.5) + \frac{1}{2}(1 - 1) = 0.194. \quad (28)$$

Target X , features $\{M, Y\}$. $v(\emptyset) = 0.5$; $v(\{M\}) = 0.5 + \frac{5}{7}(0.5) = 0.857$; $v(\{Y\}) = \mathbb{E}[X \mid Y=1] = 0.778$ (as above); $v(\{M, Y\}) = \mathbb{E}[X \mid M=1] = 0.857$ (since $X \perp Y \mid M$).

$$\phi_M = \frac{1}{2}(0.857 - 0.5) + \frac{1}{2}(0.857 - 0.778) = 0.219, \quad (29)$$

$$\phi_Y = \frac{1}{2}(0.778 - 0.5) + \frac{1}{2}(0.857 - 0.857) = 0.139. \quad (30)$$

B.4 Causal SHAP at Instance 1

We apply Causal SHAP (Definition 23) and compare against plain SHAP.

Target Y , features $\{X, M\}$. $v_{\text{causal}}(\emptyset) = 0.5$; $v_{\text{causal}}(\{X\}) = \mathbb{E}[M \mid \text{do}(X=1)] = 1$ (M is downstream of X : $P(M \mid \text{do}(X=1)) = \mathcal{N}(1, 0.1)$); $v_{\text{causal}}(\{M\}) = 1$ ($\text{do}(M=1)$ sets $f_Y = M = 1$ directly, $P(X \mid \text{do}(M=1)) = P(X)$ since X has no causal parents); $v_{\text{causal}}(\{X, M\}) = 1$. All values match plain SHAP, so $\phi_X = \phi_M = 0.25$.

Target M , features $\{X, Y\}$. $v_{\text{causal}}(\emptyset) = 0.5$; $v_{\text{causal}}(\{X\}) = 1$ (Y is downstream of X , same as plain SHAP); $v_{\text{causal}}(\{Y\}) = \mathbb{E}_{X \sim P(X)}[0.5X + 0.5 \cdot 1] = 0.75$ (Y does not cause X , so $P(X \mid \text{do}(Y=1)) = P(X)$); differs from plain SHAP's 0.889; $v_{\text{causal}}(\{X, Y\}) = 1$.

$$\phi_X = \frac{1}{2}(1 - 0.5) + \frac{1}{2}(1 - 0.75) = 0.375, \quad (31)$$

$$\phi_Y = \frac{1}{2}(0.75 - 0.5) + \frac{1}{2}(1 - 1) = 0.125. \quad (32)$$

Target X , features $\{M, Y\}$. $v_{\text{causal}}(\emptyset) = 0.5$; $v_{\text{causal}}(\{M\}) = 0.857$ (Y is downstream of M but f_X does not depend on Y since $X \perp Y \mid M$, so $\mathbb{E}_{Y \sim P(Y \mid \text{do}(M=1))}[f_X(1, Y)] = f_X(1, \cdot) = 0.857$;

same as plain SHAP); $v_{\text{causal}}(\{Y\}) = \mathbb{E}_{M \sim P(M)}[f_X(M, 1)] = 0.5$ (M is not a descendant of Y , so $P(M \mid \text{do}(Y=1)) = P(M)$ and $\mathbb{E}[f_X(M, 1)] = 0.5 + \frac{5}{7}(\mathbb{E}[M] - 0.5) = 0.5$; differs from plain SHAP's 0.778); $v_{\text{causal}}(\{M, Y\}) = 0.857$.

$$\phi_M = \frac{1}{2}(0.857 - 0.5) + \frac{1}{2}(0.857 - 0.5) = 0.357, \quad (33)$$

$$\phi_Y = \frac{1}{2}(0.5 - 0.5) + \frac{1}{2}(0.857 - 0.857) = 0. \quad (34)$$

B.5 PCI at Instance 1

We work at the factual instance $X^* = M^* = Y^* = 1$ with the absolute-difference score (Example 21): $ci(y^s, y^n, y^*) = |y^n - y^*| - |y^s - y^*|$. Both worlds run the PSCM with fresh exogenous noise drawn from P_U : the necessity world replaces the suspect A with $A' \sim P(A)$ and propagates; the sufficiency world restores it to a^* and propagates. For target B , $\mathbf{S}$ is the two input features of B and $\mathbf{W}$ the remaining input; Γ_s is uniform over the three non-empty subsets of $\mathbf{S}$ and Γ_w uniform over $2^{\mathbf{W}}$.

$R(X \rightsquigarrow Y)$. $\mathbf{S} = \{X, M\}$, $\mathbf{W} = \{M\}$, $w^* = m^* = 1$, and $X' \sim \mathcal{N}(0.5, 0.25)$. Three of the four accepted $(\mathbf{C}, \mathbf{T})$ pairs intervene on X , each with weight $\frac{1}{4}$; the fourth, $(\mathbf{C}=\{M\}, \mathbf{T}=\emptyset)$, does not touch X and contributes nothing. The key observation is that the witness pair $(\mathbf{C}=\{X\}, \mathbf{T}=\{M\})$ pins M at m^* in both worlds, which equalizes their distributions and zeros out $\bar{c}$; the other two pairs let M propagate freely. Computing $\mathbb{E}[|Y^n - 1|]$ and $\mathbb{E}[|Y^s - 1|]$ in closed form via the folded-normal formula gives:

$(\mathbf{C}, \mathbf{T})$	$\mathbb{E}[Y^n - 1]$	$\mathbb{E}[Y^s - 1]$	$\bar{c}$
$(\{X\}, \emptyset)$	0.677	0.357	0.320
$(\{X\}, \{M\})$ [witness]	0.252	0.252	0.000
$(\{X, M\}, \emptyset)$	0.677	0.252	0.425

Combining at weight $\frac{1}{4}$ each,

$$R(X \rightsquigarrow Y \mid \mathbf{W}=\{M\}) = \frac{1}{4}(0.320 + 0 + 0.425) \approx 0.186.$$

$R(Y \rightsquigarrow X)$. X is exogenous: no *do*-intervention on $\{M, Y\}$ alters its structural equation, so necessity and sufficiency distributions coincide and $R(Y \rightsquigarrow X \mid \mathbf{W}=\{M\}) = 0$.

Other cells. The remaining entries of Table 7 (D-MY, D-XM, D-YM, D-MX, D-MXY, and the $\mathbf{W} = \emptyset$ column) follow the same closed-form machinery; full per-pair computations are in the companion notebook `signal_mediation_computations.ipynb`.

C Setup for the Desert-Traveller PCI Computation

This appendix records the structural model and the full PCI specification behind the desert-traveller comparison of Section 8. The basic-variant results (Table 12) and the weak-poison results (Table 14) are produced by running the PCI Monte Carlo search of Section 3 on the scenario-specific model below, with the component choices listed here; a parallel hand-rolled enumeration matches every value within Monte Carlo noise (companion notebook `docs/source/desert_traveler.ipynb`).

C.1 Structural model (basic variant)

To compare directly with Pearl, we run PCI under the same forensic conditioning: one structural model per scenario, with u pinned at the scenario's forensic realisation. The endogenous variables are $X, P, c, d, Y \in \{0, 1\}$ with

$$c = P(u' \vee X'), \quad d = X(u \vee P'), \quad y = c \vee d,$$

where primes denote Boolean complement ($u' = 1 - u$, etc.), c indicates the cyanide path operating (poisoned water drunk) and d the dehydration path operating (drank from empty canteen).

C.2 PCI components

The PCI components, all specified in advance:

- Suspects $\mathbf{S} = \{X, P\}$; candidate causes are the singletons $\{X\}$ and $\{P\}$.
- Witness pool $\mathbf{W} = \{c, d\}$ (the mediators); four subsets $\mathbf{T} \subseteq \mathbf{W}$, weighted by Γ cardinality-uniform (uniform on $|\mathbf{T}|$, then uniform within each cardinality class).
- Non-cause suspect (the one not in $\mathbf{C}$) is *not* intervened on or hard-coded; it follows the SCM under the noise prior on its own coin, consistent with the formal definition’s requirement that only $\mathbf{C}$ and $\mathbf{T}$ are intervened (Definition 18).
- Alternative-value distribution Δ deterministically flips $1 \rightarrow 0$.
- Noise distribution P_U is the forensic posterior, a point mass at the scenario’s u . We do *not* condition on the outcome Y itself; the forensic evidence sits above Y in the SCM, so conditioning on it is not the “fully conditional” anti-pattern of Section 3.

For the weak-poison variant (Section 8.4) the only mechanical changes are that the witness pool becomes $\mathbf{W} = \{c, d, v_C\}$ (the new “cyanide-was-fatal” mediator v_C is included) and that the Scenario A forensic posterior pins both $u = 0$ and $\xi = 1$; Scenario B’s posterior pins $u = 1$, with ξ structurally irrelevant since the cyanide path is dormant.