

the normal distribution

- normal distribution = gaussian distribution = bell curve
 - just two parameters: mean and standard deviation (or variance)
 - symmetric, unimodal, asymptotic
 - mean = median = mode
 - common distribution; central limit theorem; galton box videos
- normal distribution is a theoretical distribution; used in inferential statistics
- proportion between standard deviations (figure 4.5)
 - proportions: 34.13%, 13.59%, 2.15%
 - can find proportion of samples within a certain range
 - e.g., mean temperature 37 °C, standard deviation 0.5 °C
population has 68% between $37.0 - 0.5 = 36.5$ and $37.0 + 0.5 = 37.5$
- z score = standard score
 - $z = (x - \mu) / \sigma$, or $z = (x - \bar{x}) / s$
 - number of standard deviations above or below the mean
 - indicates how large a score is relative to the population
 - e.g., z score of body temperature 38 °C: $(38-37)/0.5 = 2$
 - e.g., z score of mean is zero
 - opposite direction: z score to raw score
 - $x = \mu + z \times \sigma$, or $x = \bar{x} + z \times s$
 - e.g., z score -3 occurs at: $x = 37 + (-3) \times 0.5 = 35.5$ °C
- percentile score
 - point that is just greater than p% of the population
 - urdan, appendix A
 - table gives area under standard normal distribution to left of z ($z \geq 0$)
 - can also use table to find proportion to the right of z
 - can also use table for negative z values
 - opposite direction: look up z value for a given percentile score
 - kelley, reference table 1; equals urdan's table minus 0.5
- so far we can
 - convert raw score to z score, and vice versa
 - convert z score to percentile (or proportion), and vice versa
- can also combine the above, e.g., convert raw score to percentile
 - e.g., what proportion of people have body temperature below 38.2 °C?
 $z = (38.2 - 37.0) / 0.5 = 2.4$, $p = 0.9918$
- can also find proportion of distribution between two z scores or two raw scores
 - what proportion of people have body temperature between 36 °C and 38 °C?