

Solving Differential Equations in R (book) - PDE examples

Karline Soetaert

Centre for Estuarine and Marine Ecology
Netherlands Institute of Ecology
The Netherlands

Abstract

This vignette contains the R-examples of chapter 10 from the book:
Soetaert, K., Cash, J.R. and Mazzia, F. (2012). Solving Differential Equations in R.
that will be published by Springer.

Chapter 10. Solving Partial Differential Equations in R.

Here the code is given without documentation. Of course, much more information
about each problem can be found in the book.

Keywords: partial differential equations, initial value problems, examples, R.

1. The heat Equation

```
N      <- 100
xgrid  <- setup.grid.1D(x.up = 0, x.down = 1, N = N)
x      <- xgrid$x.mid
D.coeff <- 0.01
Diffusion <- function (t, Y, parms){
  tran <- tran.1D(C = Y, C.up = 0, C.down = 1,
                 D = D.coeff, dx = xgrid)
  list(dY = tran$dC, flux.up = tran$flux.up,
       flux.down = tran$flux.down)
}
Yini  <- sin(pi*x)
times <- seq(from = 0, to = 5, by = 0.01)
print(system.time(
  out <- ode.1D(y = Yini, times = times, func = Diffusion,
               parms = NULL, dims = N)
))

user  system elapsed
0.39  0.00   0.40

par (mfrow=c(1, 2))
plot(out[1, 2:(N+1)], x, type = "l", lwd = 2,
```




Figure 1: The solution of the heat equation. See book for more information.

```

xlab = "Variable, Y", ylab = "Distance, x")
for (i in seq(2, length(times), by = 50))
  lines(out[i, 2:(N+1)], x)
image(out, grid = x, mfrow = NULL, ylab = "Distance, x",
      main = "Y")

```

2. The Wave Equation

```

dx      <- 0.2
xgrid <- setup.grid.1D(x.up = -100, x.down = 100, dx.1 = dx)
x      <- xgrid$x.mid
N      <- xgrid$N
lam    <- 0.05
uini   <- exp(-lam*x^2)
vini   <- rep(0, N)
yini   <- c(uini, vini)
times  <- seq(from = 0, to = 50, by = 1)
wave <- function(t, y, parms) {
  u <- y[1:N]
  v <- y[(N+1):(2*N)]

  du <- v
  dv <- tran.1D(C = u, C.up = 0, C.down = 0, D = 1,
               dx = xgrid)$dC

  return(list(c(du, dv)))
}
out <- ode.1D(func = wave, y = yini, times = times,
             parms = NULL, method = "adams",
             dims = N, names = c("u", "v"))
u <- subset(out, which = "u")
analytic <- function(t, x)
  0.5 * (exp(-lam * (x+1*t)^2) + exp(-lam * (x-1*t)^2))
OutAna <- outer(times, x, FUN = analytic)
max(abs(u - OutAna))

[1] 0.002188562

outtime <- seq(from = 0, to = 50, by = 10)
matplot.1D(out, which = "u", subset = time %in% outtime,
  grid = x, xlab = "x", ylab = "u", type = "l",
  lwd = 2, xlim = c(-50, 50),
  col = c("black", rep("darkgrey", 5)))
legend("topright", lty = 1:6, lwd = 2,
  col = c("black", rep("darkgrey", 5)),
  title = "t = ", legend = outtime)

```




Figure 2: The 1-D wave equation. See book for explanation.

3. Laplace Equation

```

Nx <- 100
Ny <- 100
xgrid <- setup.grid.1D (x.up = 0, x.down = 1, N = Nx)
ygrid <- setup.grid.1D (x.up = 0, x.down = 1, N = Ny)
x      <- xgrid$x.mid
y      <- ygrid$y.mid
laplace <- function(t, U, parms) {
  w <- matrix(nrow = Nx, ncol = Ny, data = U)
  dw <- tran.2D(C = w, C.x.up = 0, C.x.down = 0,
               flux.y.up = 0,
               flux.y.down = -1 * sin(pi*x)*pi*sinh(pi),
               D.x = 1, D.y = 1,
               dx = xgrid, dy = ygrid)$dC
  list(dw)
}
print(system.time(
  out <- steady.2D(y = runif(Nx*Ny), func = laplace,
                  parms = NULL, nspec = 1,
                  dims = c(Nx, Ny), lrw = 1e7)
))

user  system elapsed
0.36   0.03   0.39

w <- matrix(nrow = Nx, ncol = Ny, data = out$y)
analytic <- function (x, y) sin(pi*x) * cosh(pi*y)
OutAna <- outer(x, y, FUN = analytic)
max(abs(w - OutAna))

[1] 0.0006024049

image(out, grid = list(x, y), main = "elliptic Laplace",
      add.contour = TRUE)

```




Figure 3: The laplace equation. See book for explanation.

4. The Advection Equation

```
adv.func <- function(t, y, p, adv.method)
  list(advection.1D(C = y, C.up = y[N], C.down = y[1],
    v = 0.1, adv.method = adv.method,
    dx = xgrid)$dC)
xgrid <- setup.grid.1D(0.3, 1.3, N = 50)
x      <- xgrid$x.mid
N      <- length(x)
yini   <- sin(pi * x)^50
times  <- seq(0, 20, 0.01)
out1 <- ode.1D(y = yini, func = adv.func, times = times,
  parms = NULL, method = "euler", dims = N,
  adv.method = "muscl")
out2 <- ode.1D(y = yini, func = adv.func, times = times,
  parms = NULL, method = "euler", dims = N,
  adv.method = "super")
```

5. The Brusselator in One Dimension

```

N      <- 50
Grid <- setup.grid.1D(x.up = 0, x.down = 1, N = N)
x1ini <- 1 + sin(2 * pi * Grid$x.mid)
x2ini <- rep(x = 3, times = N)
yini  <- c(x1ini, x2ini)
brusselator1D <- function(t, y, parms) {

  X1 <- y[1:N]
  X2 <- y[(N+1):(2*N)]

  dX1 <- 1 + X1^2*X2 - 4*X1 +
    tran.1D (C = X1, C.up = 1, C.down = 1,
             D = 0.02, dx = Grid)$dC
  dX2 <- 3*X1 - X1^2*X2 +
    tran.1D (C = X2, C.up = 3, C.down = 3,
             D = 0.02, dx = Grid)$dC

  list(c(dX1, dX2))
}
times <- seq(from = 0, to = 10, by = 0.1)
print(system.time(
  out <- ode.1D(y = yini, func = brusselator1D,
               times = times, parms = NULL, nspec = 2,
               names = c("X1", "X2"), dims = N)
))

user  system elapsed
0.3    0.0    0.3

par(mfrow = c(2, 2))
image(out, mfrow = NULL, grid = Grid$x.mid,
      which = "X1", method = "contour")
image(out, mfrow = NULL, grid = Grid$x.mid,
      which = "X1")
par(mar = c(1, 1, 1, 1))
image(out, mfrow = NULL, grid = Grid$x.mid,
      which = "X1", method = "persp", col = NA)
image(out, mfrow = NULL, grid = Grid$x.mid,
      which = "X1", method = "persp", border = NA,
      shade = 0.3 )

```





Figure 4: The 1-D Brusselator model. See book for explanation.

6. The Brusselator in 2-D

```
brusselator2D <- function(t, y, parms) {

  X1 <- matrix(nrow = Nx, ncol = Ny,
               data = y[1:(Nx*Ny)])
  X2 <- matrix(nrow = Nx, ncol = Ny,
               data = y[(Nx*Ny+1) : (2*Nx*Ny)])

  dX1 <- 1 + X1^2*X2 - 4*X1 +
    tran.2D (C = X1, D.x = D_X1, D.y = D_X1,
             dx = Gridx, dy = Gridy)$dC
  dX2 <- 3*X1 - X1^2*X2 +
    tran.2D (C = X2, D.x = D_X2, D.y = D_X2,
             dx = Gridx, dy = Gridy)$dC

  list(c(dX1, dX2))
}

Nx <- 50
Ny <- 50
Gridx <- setup.grid.1D(x.up = 0, x.down = 1, N = Nx)
Gridy <- setup.grid.1D(x.up = 0, x.down = 1, N = Ny)
D_X1 <- 2
D_X2 <- 8*D_X1
X1ini <- matrix(nrow = Nx, ncol = Ny, data = runif(Nx*Ny))
X2ini <- matrix(nrow = Nx, ncol = Ny, data = runif(Nx*Ny))
yini <- c(X1ini, X2ini)
times <- 0:8
print(system.time(
  out <- ode.2D(y = yini, parms = NULL, func = brusselator2D,
               nspec = 2, dimsens = c(Nx, Ny), times = times,
               lrw = 2000000, names=c("X1", "X2"))
))

user  system elapsed
2.87   0.03   2.98

par(oma = c(0,0,1,0))
image(out, which = "X1", xlab = "x", ylab = "y",
      mfrow = c(3, 3), ask = FALSE,
      main = paste("t = ", times),
      grid = list(x = Gridx$x.mid, y = Gridy$x.mid))
mtext(side = 3, outer = TRUE, cex = 1.25, line = -1,
      "2-D Brusselator, species X1")
```




Figure 5: Solution of the 2-D Brusselator. See book for explanation.




Figure 6: The Laplace equation in polar coordinates. See book for explanation.

7. The Laplace Equation in Polar Coordinates

```
Nr <- 100
Np <- 100
r      <- seq(2, 4, len = Nr+1)
theta  <- seq(0, 2*pi, len = Np+1)
theta.mid <- 0.5*(theta[-1] + theta[-Np])
Model <- function(t, C, p) {
  y = matrix(nrow = Nr, ncol = Np, data = C)
  tran <- tran.polar (y, D.r = 1, r = r, theta = theta,
    C.r.up = 0, C.r.down = 4 * sin(5*theta.mid),
    cyclicBnd = 2)
  list(tran$dC)
}
STD <- steady.2D(y = runif(Nr*Np), parms = NULL,
  func = Model, dims = c(Nr, Np),
  lrw = 1e6, cyclicBnd = 2)
OUT <- polar2cart (STD, r = r, theta = theta,
  x = seq(-4, 4, len = 400),
  y = seq(-4, 4, len = 400))

image(OUT, main = "Laplace")
```

8. The Time-dependent 2-D Sine-Gordon Equation

```

Nx <- 100
Ny <- 100
xgrid <- setup.grid.1D(-7, 7, N=Nx)
ygrid <- setup.grid.1D(-7, 7, N=Ny)
x <- xgrid$x.mid
y <- ygrid$y.mid
sinegordon2D <- function(t, C, parms) {

  u <- matrix(nrow = Nx, ncol = Ny,
              data = C[1 : (Nx*Ny)])
  v <- matrix(nrow = Nx, ncol = Ny,
              data = C[(Nx*Ny+1) : (2*Nx*Ny)])

  dv <- tran.2D (C = u, C.x.up = 0, C.x.down = 0,
                 C.y.up = 0, C.y.down = 0,
                 D.x = 1, D.y = 1,
                 dx = xgrid, dy = ygrid)$dC - sin(u)
  list(c(v, dv))
}
peak <- function (x, y, x0 = 0, y0 = 0)
  exp(-((x-x0)^2 + (y-y0)^2))
uini <- outer(x, y,
  FUN = function(x, y) peak(x, y, 2,2) + peak(x, y,-2,-2)
    + peak(x, y,-2,2) + peak(x, y, 2,-2))
vini <- rep(0, Nx*Ny)
times <- 0:3
print(system.time(
out <- ode.2D (y = c(uini, vini), times = times,
               parms = NULL, func = sinegordon2D,
               names = c("u", "v"),
               dims = c(Nx, Ny), method = "ode45")
))

user  system elapsed
0.77   0.00   0.80

mr <- par(mar = c(0, 0, 1, 0))
image(out, main = paste("time =", times), which = "u",
      grid = list(x = x, y = y), method = "persp",
      border = NA, col = "grey", box = FALSE,
      shade = 0.5, theta = 30, phi = 60, mfrow = c(2, 2),
      ask = FALSE)
par(mar = mr)

```




Figure 7: The 2-D sine-gordon equation. See book for explanation.

9. The Nonlinear Schrodinger Equation

```

alf <- 0.5
gam <- 1
Schrodinger <- function(t, u, parms) {
  du <- 1i * tran.1D (C = u, D = 1, dx = xgrid)$dC +
    1i * gam * abs(u)^2 * u
  list(du)
}
N <- 300
xgrid <- setup.grid.1D(-20, 80, N = N)
x <- xgrid$x.mid
c1 <- 1
c2 <- 0.1
sech <- function(x) 2/(exp(x) + exp(-x))
soliton <- function (x, c1)
  sqrt(2*alf/gam) * exp(0.5*1i*c1*x) * sech(sqrt(alf)*x)
yini <- soliton(x, c1) + soliton(x-25, c2)
times <- seq(0, 40, by = 0.1)
print(system.time(
  out <- ode.1D(y = yini, parms = NULL, func = Schrodinger,
    times = times, dims = 300, method = "adams")
))

user  system elapsed
2.49   0.00   2.52

image(abs(out), grid = x, ylab = "x", main = "two solitons")

```

Affiliation:

Karline Soetaert

Centre for Estuarine and Marine Ecology (CEME)

Netherlands Institute of Ecology (NIOO)

4401 NT Yerseke, Netherlands E-mail: k.soetaert@nioo.knaw.nl

URL: <http://www.nioo.knaw.nl/users/ksoetaert>




Figure 8: Solution of the Schrödinger equation. See book for explanation.