


# HE Plot Examples

Michael Friendly

Using `heplots` version 1.3-2 and `candisc` version 0.7-1; Date: 2016-10-17



## Abstract

This vignette provides some worked examples of the analysis of multivariate linear models (MLMs) with graphical methods for visualizing results using the `heplots` package and the `candisc` package. The emphasis here is on using these methods in R, and understanding how they help reveal aspects of these models that might not be apparent from other graphical displays.

No attempt is made here to describe the theory of MLMs or the statistical details behind HE plots and their reduced-rank canonical cousins. For that, see Fox et al. (2009); Friendly (2007, 2006).

## Contents

<b>1</b>	<b>MANOVA Designs</b>	<b>1</b>	<b>2</b>	<b>MMRA Designs</b>	<b>17</b>
1.1	Plastic film data . . . . .	1	2.1	Rohwer data . . . . .	18
1.2	Mock jury decisions . . . . .	5	2.2	Hernior data . . . . .	23
1.3	Egyptian skulls . . . . .	11	2.3	SocGrades . . . . .	26

## 1 MANOVA Designs

### 1.1 Plastic film data

An experiment was conducted to determine the optimum conditions for extruding plastic film. Three responses, `tear` resistance, film `gloss` and film `opacity` were measured in relation to two factors, `rate` of extrusion and amount of an `additive`, both of these being set to two values, High and Low. The design is thus a  $2 \times 2$  MANOVA, with  $n = 5$  per cell. This example illustrates 2D and 3D HE plots, the difference between “effect” scaling and “evidence” (significance) scaling, and visualizing composite linear hypotheses.

We begin with an overall MANOVA for the two-way MANOVA model. Because each effect has 1 df, all of the multivariate statistics are equivalent, but we specify `test.statistic="Roy"` because Roy’s test has a natural visual interpretation in HE plots.

```
> plastic.mod <- lm(cbind(tear, gloss, opacity) ~ rate*additive, data=Plastic)
> Anova(plastic.mod, test.statistic="Roy")
```

```
Type II MANOVA Tests: Roy test statistic
      Df test stat approx F num Df den Df Pr(>F)
rate      1      1.619      7.55      3      14 0.003 **
additive   1      0.912      4.26      3      14 0.025 *
rate:additive 1      0.287      1.34      3      14 0.302
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

For the three responses jointly, the main effects of **rate** and **additive** are significant, while their interaction is not. In some approaches to testing effects in multivariate linear models (MLM), significant multivariate tests are often followed by univariate tests on each of the responses separately to determine which responses contribute to each significant effect. In R, these analyses are most conveniently performed using the `update()` method for the `mlm` object `plastic.mod`.

```
> Anova(update(plastic.mod, tear ~ .))
```

```
Anova Table (Type II tests)
Response: tear
      Sum Sq Df F value Pr(>F)
rate      1.74  1    15.8 0.0011 **
additive   0.76  1     6.9 0.0183 *
rate:additive 0.00  1     0.0 0.9471
Residuals  1.76 16
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
> Anova(update(plastic.mod, gloss ~ .))
```

```
Anova Table (Type II tests)
Response: gloss
      Sum Sq Df F value Pr(>F)
rate    1.301  1    7.92 0.012 *
additive 0.612  1    3.73 0.071 .
rate:additive 0.544  1    3.32 0.087 .
Residuals 2.628 16
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
> Anova(update(plastic.mod, opacity ~ .))
```

```
Anova Table (Type II tests)
Response: opacity
      Sum Sq Df F value Pr(>F)
rate      0.4  1    0.10 0.75
additive   4.9  1    1.21 0.29
rate:additive 4.0  1    0.98 0.34
Residuals 64.9 16
```

The results above show significant main effects for **tear**, a significant main effect of **rate** for **gloss**, and no significant effects for **opacity**, but they don't shed light on the *nature* of these effects. Traditional univariate plots of the means for each variable separately are useful, but they don't allow visualization of the *relations* among the response variables.

We can visualize these effects for pairs of variables in an HE plot, showing the “size” and orientation of hypothesis variation (**H**) in relation to error variation (**E**) as ellipsoids. When, as here, the model terms have 1 degree of freedom, the **H** ellipsoids degenerate to a line.

```

> # Compare evidence and effect scaling
> colors = c("red", "darkblue", "darkgreen", "brown")
> heplot(plastic.mod, size="evidence", col=colors, cex=1.25)
> heplot(plastic.mod, size="effect", add=TRUE, lwd=4, term.labels=FALSE, col=colors)

```

With effect scaling, both the  $H$  and  $E$  sums of squares and products matrices are both divided by the error df, giving multivariate analogs of univariate measures of effect size, e.g.,  $(\bar{y}_1 - \bar{y}_2)/s$ . With significance scaling, the  $H$  ellipse is further divided by  $\lambda_\alpha$ , the critical value of Roy's largest root statistic. This scaling has the property that an  $H$  ellipse will protrude somewhere outside the  $E$  ellipse *iff* the multivariate test is significant at level  $\alpha$ . Figure 1 shows both scalings, using a thinner line for significance scaling. Note that the (degenerate) ellipse for **additive** is significant, but does not protrude outside the  $E$  ellipse in this view. All that is guaranteed is that it will protrude somewhere in the 3D space of the responses.

By design, means for the levels of interaction terms are not shown in the HE plot, because doing so in general can lead to messy displays. We can add them here for the term **rate:additive** as follows:

```

> ## add interaction means
> intMeans <- termMeans(plastic.mod, 'rate:additive', abbrev.levels=2)
> #rownames(intMeans) <- apply(expand.grid(c('Lo', 'Hi'), c('Lo', 'Hi')), 1, paste, collapse=':')
> points(intMeans[,1], intMeans[,2], pch=18, cex=1.2, col="brown")
> text(intMeans[,1], intMeans[,2], rownames(intMeans), adj=c(0.5,1), col="brown")
> lines(intMeans[c(1,3),1], intMeans[c(1,3),2], col="brown")
> lines(intMeans[c(2,4),1], intMeans[c(2,4),2], col="brown")

```




Figure 1: HE plot for effects on **tear** and **gloss** according to the factors **rate**, **additive** and their interaction, **rate:additive**. The thicker lines show effect size scaling, the thinner lines show significance scaling.

The factor means in this plot (Figure 1) have a simple interpretation: The high **rate** level yields greater **tear** resistance but lower **gloss** than the low level. The high **additive** amount produces greater **tear** resistance and greater **gloss**.

The `rate:additive` interaction is not significant overall, though it approaches significance for `gloss`. The cell means for the combinations of `rate` and `additive` shown in this figure suggest an explanation, for tutorial purposes: with the low level of `rate`, there is little difference in `gloss` for the levels of `additive`. At the high level of `rate`, there is a larger difference in `gloss`. The *H* ellipse for the interaction of `rate:additive` therefore “points” in the direction of `gloss` indicating that this variable contributes to the interaction in the multivariate tests.

In some MANOVA models, it is of interest to test sub-hypotheses of a given main effect or interaction, or conversely to test composite hypotheses that pool together certain effects to test them jointly. All of these tests (and, indeed, the tests of terms in a given model) are carried out as tests of general linear hypotheses in the MLM.

In this example, it might be useful to test two composite hypotheses: one corresponding to both main effects jointly, and another corresponding to no difference among the means of the four groups (equivalent to a joint test for the overall model). These tests are specified in terms of subsets or linear combinations of the model parameters.

```
> plastic.mod
```

```
Call:
```

```
lm(formula = cbind(tear, gloss, opacity) ~ rate * additive, data = Plastic)
```

```
Coefficients:
```

	tear	gloss	opacity
(Intercept)	6.30	9.56	3.74
rateHigh	0.58	-0.84	-0.60
additiveHigh	0.38	0.02	0.10
rateHigh:additiveHigh	0.02	0.66	1.78

Thus, for example, the joint test of both main effects tests the parameters `rateHigh` and `additiveHigh`.

```
> print(linearHypothesis(plastic.mod, c("rateHigh", "additiveHigh"), title="Main effects"),
      SSP=FALSE)
```

```
Multivariate Tests: Main effects
```

	Df	test	stat	approx F	num Df	den Df	Pr(>F)
Pillai	2	0.71161	2.7616	6	30	0.029394	*
Wilks	2	0.37410	2.9632	6	28	0.022839	*
Hotelling-Lawley	2	1.44400	3.1287	6	26	0.019176	*
Roy	2	1.26253	6.3127	3	15	0.005542	**

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
> print(linearHypothesis(plastic.mod, c("rateHigh", "additiveHigh", "rateHigh:additiveHigh"),
      title="Groups"), SSP=FALSE)
```

```
Multivariate Tests: Groups
```

	Df	test	stat	approx F	num Df	den Df	Pr(>F)
Pillai	3	1.14560	3.2948	9	48.000	0.003350	**
Wilks	3	0.17802	3.9252	9	34.223	0.001663	**
Hotelling-Lawley	3	2.81752	3.9654	9	38.000	0.001245	**
Roy	3	1.86960	9.9712	3	16.000	0.000603	***

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Correspondingly, we can display these tests in the HE plot by specifying these tests in the `hypothesis` argument to `heplot()`, as shown in Figure 2.

Finally, a 3D HE plot can be produced with `heplot3d()`, giving Figure 3. This plot was rotated interactively to a view that shows both main effects protruding outside the error ellipsoid.

```
> colors = c("pink", "darkblue", "darkgreen", "brown")
> heplot3d(plastic.mod, col=colors)
```

```
> heplot(plastic.mod, hypotheses=list("Group" =
  c("rateHigh", "additiveHigh", "rateHigh:additiveHigh"),
  col=c(colors, "purple"),
  lwd=c(2, 3, 3, 3, 2), cex=1.25)
> heplot(plastic.mod, hypotheses=list("Main effects" =
  c("rateHigh", "additiveHigh")), add=TRUE,
  col=c(colors, "darkgreen"), cex=1.25)
```




Figure 2: HE plot for `tear` and `gloss`, supplemented with ellipses representing the joint tests of main effects and all group differences

## 1.2 Effects of physical attractiveness on mock jury decisions

In a social psychology study of influences on jury decisions by Plaster (1989), male participants (prison inmates) were shown a picture of one of three young women. Pilot work had indicated that one woman was beautiful, another of average physical attractiveness, and the third unattractive. Participants rated the woman they saw on each of twelve attributes on scales of 1–9. These measures were used to check on the manipulation of “attractiveness” by the photo.

Then the participants were told that the person in the photo had committed a Crime, and asked to rate the seriousness of the crime and recommend a prison sentence, in Years. The data are contained in the data frame `MockJury`.<sup>1</sup>

```
> str(MockJury)
```

<sup>1</sup>The data were made available courtesy of Karl Wuensch, from <http://core.ecu.edu/psyc/wuenschk/StatData/PLASTER.dat>




Figure 3: 3D HE plot for the plastic film data

```
'data.frame':      114 obs. of  17 variables:
 $ Attr      : Factor w/ 3 levels "Beautiful","Average",...: 1 1 1 1 1 1 1 1 1 1 ...
 $ Crime     : Factor w/ 2 levels "Burglary","Swindle": 1 1 1 1 1 1 1 1 1 1 ...
 $ Years     : int  10 3 5 1 7 7 3 7 2 3 ...
 $ Serious   : int  8 8 5 3 9 9 4 4 5 2 ...
 $ exciting  : int  6 9 3 3 1 1 5 4 4 6 ...
 $ calm      : int  9 5 4 6 1 5 6 9 8 8 ...
 $ independent : int  9 9 6 9 5 7 7 2 8 7 ...
 $ sincere   : int  8 3 3 8 1 5 6 9 7 5 ...
 $ warm      : int  5 5 6 8 8 8 7 6 1 7 ...
 $ phyattr   : int  9 9 7 9 8 8 8 5 9 8 ...
 $ sociable  : int  9 9 4 9 9 9 7 2 1 9 ...
 $ kind      : int  9 4 2 9 4 5 5 9 5 7 ...
 $ intelligent : int  6 9 4 9 7 8 7 9 9 9 ...
 $ strong    : int  9 5 5 9 9 9 5 2 7 5 ...
 $ sophisticated: int  9 5 4 9 9 9 6 2 7 6 ...
 $ happy     : int  5 5 5 9 8 9 5 2 6 8 ...
 $ ownPA     : int  9 7 5 9 7 9 6 5 3 6 ...
```

Sample sizes were roughly balanced for the independent variables in the three conditions of the attractiveness of the photo, and the combinations of this with **Crime**:

```
> table(MockJury$Attr)
```

```
Beautiful    Average Unattractive
      39          38          37
```

```
> table(MockJury$Attr, MockJury$Crime)
```

```
          Burglary Swindle
Beautiful      21      18
Average        18      20
Unattractive   20      17
```

The main questions of interest were: (a) Does attractiveness of the “defendent” influence the sentence or perceived seriousness of the crime? (b) Does attractiveness interact with the nature of the crime?

But first, we try to assess the ratings of the photos in relation to the presumed categories of the independent variable `Attr`. The questions here are (a) do the ratings of the photos on physical attractiveness (`phyattr`) confirm the original classification? (b) how do other ratings differentiate the photos? To keep things simple, we consider only a few of the other ratings in a one-way MANOVA.

```
> (jury.mod1 <- lm( cbind(phyattr, happy, independent, sophisticated) ~ Attr, data=MockJury))
```

```
Call:
lm(formula = cbind(phyattr, happy, independent, sophisticated) ~
    Attr, data = MockJury)
```

```
Coefficients:
              phyattr    happy    independent    sophisticated
(Intercept)    8.282    5.359    6.410    6.077
AttrAverage    -4.808    0.430    0.537    -1.340
AttrUnattractive -5.390   -1.359   -1.410   -1.753
```

```
> Anova(jury.mod1, test="Roy")
```

```
Type II MANOVA Tests: Roy test statistic
      Df test stat approx F num Df den Df Pr(>F)
Attr  2      1.77    48.2      4    109 <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Note that Beautiful is the baseline category of `Attr`, so the intercept term gives the means for this level. We see that the means are significantly different on all four variables collectively, by a joint multivariate test. A traditional analysis might follow up with univariate ANOVAs for each measure separately.

As an aid to interpretation of the MANOVA results We can examine the test of `Attr` in this model with an HE plot for pairs of variables, e.g., for `phyattr` and `happy` (Figure 4). The means in this plot show that Beautiful is rated higher on physical attractiveness than the other two photos, while Unattractive is rated less happy than the other two. Comparing the sizes of the ellipses, differences among group means on physical attractiveness contributes more to significance than do ratings on happy.

```
> heplot(jury.mod1, main="HE plot for manipulation check")
```

The HE plot for all pairs of variables (Figure 5) shows that the means for `happy` and `independent` are highly correlated, as are the means for `phyattr` and `sophisticated`. In most of these pairwise plots, the means form a triangle rather than a line, suggesting that these attributes are indeed measuring different aspects of the photos.

With 3 groups and 4 variables, the  $\mathbf{H}$  ellipsoid has only  $s = \min(df_h, p) = 2$  dimensions. `candisc()` carries out a canonical discriminant analysis for the MLM and returns an object that can be used to show an HE plot in the space of the canonical dimensions. This is plotted in Figure 6.

```
> jury.can <- candisc(jury.mod1)
> jury.can
```




Figure 4: HE plot for ratings of `phyattr` and `happy` according to the classification of photos on `Attr`

Canonical Discriminant Analysis for `Attr`:

	CanRsq	Eigenvalue	Difference	Percent	Cumulative
1	0.639	1.767	1.6	91.33	91.3
2	0.144	0.168	1.6	8.67	100.0

Test of H0: The canonical correlations in the current row and all that follow are zero

	LR test stat	approx F	num Df	den Df	Pr(> F)
1	0.309	43.9	4	220	< 2e-16 ***
2	0.856	18.6	1	111	3.5e-05 ***

---  
Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

From this we can see that 91% of the variation among group means is accounted for by the first dimension, and this is nearly completely aligned with `phyattr`. The second dimension, accounting for the remaining 9% is determined nearly entirely by ratings on `happy` and `independent`. This display gives a relatively simple account of the results of the MANOVA and the relations of each of the ratings to discrimination among the photos.

Proceeding to the main questions of interest, we carry out a two-way MANOVA of the responses `Years` and `Serious` in relation to the independent variables `Attr` and `Crime`.

```
> # influence of Attr of photo and nature of crime on Serious and Years
> jury.mod2 <- lm( cbind(Serious, Years) ~ Attr * Crime, data=MockJury)
> Anova(jury.mod2, test="Roy")
```



```
> pairs(jury.mod1)
```




Figure 5: HE plots for all pairs of ratings according to the classification of photos on **Attr**

```
Type II MANOVA Tests: Roy test statistic
              Df test stat approx F num Df den Df Pr(>F)
Attr          2    0.0756     4.08      2   108 0.020 *
Crime         1    0.0047     0.25      2   107 0.778
Attr:Crime    2    0.0501     2.71      2   108 0.071 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

We see that there is a nearly significant interaction between **Attr** and **Crime** and a strong effect of **Attr**.

The HE plot shows that the nearly significant interaction of **Attr:Crime** is mainly in terms of differences among the groups on the response of **Years** of sentence, with very little contribution of **Serious**. We explore this interaction in a bit more detail below. The main effect of **Attr** is also dominated by differences among groups on **Years**.

If we assume that **Years** of sentence is the main outcome of interest, it also makes sense to carry out a step-down test of this variable by itself, controlling for the rating of seriousness (**Serious**) of the crime. The model `jury.mod3` below is equivalent to an ANCOVA for **Years**.

```
> # stepdown test (ANCOVA), controlling for Serious
> jury.mod3 <- lm( Years ~ Serious + Attr * Crime, data=MockJury)
> t(coef(jury.mod3))

(Intercept) Serious AttrAverage AttrUnattractive CrimeSwindle
[1,]      0.011612 0.83711      0.39586      0.60285     -0.26302
```

```
> opar <- par(xpd=TRUE)
> heplot(jury.can, prefix="Canonical dimension", main="Canonical HE plot")
> par(opar)
```




Figure 6: Canonical discriminant HE plot for the MockJury data

```
[1,] AttrAverage:CrimeSwindle AttrUnattractive:CrimeSwindle
      -0.53701                2.5123
```

```
> Anova(jury.mod3)
```

Anova Table (Type II tests)

```
Response: Years
      Sum Sq Df F value Pr(>F)
Serious    379  1  41.14 3.9e-09 ***
Attr        74  2   4.02  0.021 *
Crime        4  1   0.43  0.516
Attr:Crime   49  2   2.67  0.074 .
Residuals  987 107

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Thus, even when adjusting for **Serious** rating, there is still a significant main effect of **Attr** of the photo, but also a hint of an interaction of **Attr** with **Crime**. The coefficient for **Serious** indicates that participants awarded 0.84 additional years of sentence for each 1 unit step on the scale of seriousness of crime.

A particularly useful method for visualizing the fitted effects in such univariate response models is provided by the **effects** package. By default **allEffects()** calculates the predicted

```
> heplot(jury.mod2)
```




Figure 7: HE plot for the two-way MANOVA for **Years** and **Serious**

values for all high-order terms in a given model, and the `plot` method produces plots of these values for each term. The statements below produce Figure 8.

```
> library(effects)
> jury.eff <- allEffects(jury.mod3)
> plot(jury.eff, ask=FALSE)
```

The effect plot for **Serious** shows the expected linear relation between that variable and **Years**. Of greater interest here is the nature of the possible interaction of **Attr** and **Crime** on **Years** of sentence, controlling for **Serious**. The effect plot shows that for the crime of Swindle, there is a much greater **Years** of sentence awarded to Unattractive defendants.

### 1.3 Egyptian skulls from five epochs

This example examines physical measurements of size and shape made on 150 Egyptian skulls from five epochs ranging from 4000 BC to 150 AD. The measures are: maximal breadth (mb), basibregmatic height (bh), basialveolar length (bl), and nasal height (nh) of each skull. See [http://www.redwoods.edu/instruct/agarwin/anth\\_6\\_measurements.htm](http://www.redwoods.edu/instruct/agarwin/anth_6_measurements.htm) for the definitions of these measures, and Figure 9 for a diagram. The question of interest is whether and how these measurements change over time. Systematic changes over time is of interest because this would indicate interbreeding with immigrant populations.




Figure 8: Effect plots for `Serious` and the `Attr * Crime` in the ANCOVA model `jury.mod3`.

```
> data(Skulls)
> str(Skulls)

'data.frame':      150 obs. of  5 variables:
 $ epoch: Ord.factor w/ 5 levels "c4000BC"<"c3300BC"<...: 1 1 1 1 1 1 1 1 1 ...
 $ mb   : num  131 125 131 119 136 138 139 125 131 134 ...
 $ bh   : num  138 131 132 132 143 137 130 136 134 134 ...
 $ bl   : num   89  92  99  96 100  89 108  93 102  99 ...
 $ nh   : num   49  48  50  44  54  56  48  48  51  51 ...

> table(Skulls$epoch)

c4000BC c3300BC c1850BC c200BC cAD150
      30      30      30      30      30
```

Note that `epoch` is an ordered factor, so the default contrasts will be orthogonal polynomials. This assumes that `epoch` values are equally spaced, which they are not. However, examining the linear and quadratic trends is useful to a first approximation.

For ease of labeling various outputs, it is useful to trim the `epoch` values and assign more meaningful variable labels.

```
> # make shorter labels for epochs
> Skulls$epoch <- factor(Skulls$epoch, labels=sub("c","",levels(Skulls$epoch)))
> # assign better variable labels
> vlab <- c("maxBreadth", "basibHeight", "basialLength", "nasalHeight")
```

We start with some simple displays of the means by epoch. From the numbers, the means don't seem to vary much. A `pairs` plot, Figure 10, joining points by `epoch` is somewhat more revealing for the bivariate relations among means.

```
> means <- aggregate(cbind(mb, bh, bl, nh) ~ epoch, data=Skulls, FUN=mean)[-1]
> rownames(means) <- levels(Skulls$epoch)
> means
```




Figure 9: Diagram of the skull measurements. Maximal breadth and basibregmatic height are the basic measures of “size” of a skull. Basilveolar length and nasal height are important anthropometric measures of “shape”.

	mb	bh	bl	nh
4000BC	131.37	133.60	99.167	50.533
3300BC	132.37	132.70	99.067	50.233
1850BC	134.47	133.80	96.033	50.567
200BC	135.50	132.30	94.533	51.967
AD150	136.17	130.33	93.500	51.367

```
> pairs(means, vlab,
      panel = function(x, y) {
        text(x, y, levels(Skulls$epoch))
        lines(x,y)
      })
```

Perhaps better for visualizing the trends over time is a set of boxplots, joining means over epoch. Using `bwplot()` from the `lattice` package requires reshaping the data from wide to long format. The following code produces Figure 11.

```
> library(lattice)
> library(reshape2)
> sklong <- melt(Skulls, id="epoch")
> bwplot(value ~ epoch | variable, data=sklong, scales="free",
      ylab="Variable value", xlab="Epoch",
      strip=strip.custom(factor.levels=paste(vlab, " (", levels(sklong$variable), ") ", sep="")),
      panel = function(x,y, ...) {
        panel.bwplot(x, y, ...)
        panel.linejoin(x,y, col="red", ...)
      })
```

Now, fit the MANOVA model, and test the effect of `epoch` with `Manova()`. We see that the multivariate means differ substantially.

```
> # fit manova model
> sk.mod <- lm(cbind(mb, bh, bl, nh) ~ epoch, data=Skulls)
> Manova(sk.mod)
```

```
Type II MANOVA Tests: Pillai test statistic
      Df test stat approx F num Df den Df Pr(>F)
epoch  4      0.353      3.51    16    580 4.7e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Perhaps of greater interest are the more focused tests of trends over time. These are based on tests of the coefficients in the model `sk.mod` being jointly equal to zero, for subsets of the (polynomial) contrasts in `epoch`.




Figure 10: Pairs plot of means of Skulls data, by epoch

```
> coef(sk.mod)
```

```

              mb              bh              bl              nh
(Intercept) 133.97333 132.54667 96.460000 50.93333
epoch.L      4.02663  -2.19251 -5.017481  1.07517
epoch.Q     -0.46325  -1.26504 -0.089087  0.12472
epoch.C     -0.46380  -0.78003  1.075174 -0.83273
epoch^4      0.34263   0.80479 -0.661360 -0.41833

```

We use `linearHypothesis()` for a multivariate test of the `epoch.L` linear effect. The linear trend is highly significant. It is not obvious from Figure 10 that maximal breadth and nasal are increasing over time, while the other two measurements have negative slopes.

```
> coef(sk.mod)["epoch.L",]
```

```

      mb      bh      bl      nh
4.0266 -2.1925 -5.0175  1.0752

```

```
> print(linearHypothesis(sk.mod, "epoch.L"), SSP=FALSE) # linear component
```




Figure 11: Boxplots of Skulls data, by epoch, for each variable

```
Multivariate Tests:
      Df test stat approx F num Df den Df Pr(>F)
Pillai 1 0.29138 14.597 4 142 5.195e-10 ***
Wilks 1 0.70862 14.597 4 142 5.195e-10 ***
Hotelling-Lawley 1 0.41119 14.597 4 142 5.195e-10 ***
Roy 1 0.41119 14.597 4 142 5.195e-10 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

`linearHypothesis()` can also be used to test composite hypotheses. Here we test all non-linear coefficients jointly. The result indicates that, collectively, all non-linear terms are not significantly different from zero.

```
> print(linearHypothesis(sk.mod, c("epoch.Q", "epoch.C", "epoch^4")), SSP=FALSE)
```

```
Multivariate Tests:
      Df test stat approx F num Df den Df Pr(>F)
Pillai 3 0.06819 0.83726 12 432.00 0.6119
Wilks 3 0.93296 0.83263 12 375.99 0.6167
Hotelling-Lawley 3 0.07063 0.82791 12 422.00 0.6216
Roy 3 0.04519 1.62676 4 144.00 0.1707
```

Again, HE plots can show the patterns of these tests of multivariate hypotheses. With four response variables, it is easiest to look at all pairwise HE plots with the `pairs.mlm()`

function. The statement below produces Figure 12. In this plot, we show the hypothesis ellipsoids for the overall effect of `epoch`, as well as those for the tests just shown for the linear trend component `epoch.L` as well as the joint test of all non-linear terms.

```
> pairs(sk.mod, variables = c(1, 4, 2, 3), hypotheses = list(Lin = "epoch.L",
  NonLin = c("epoch.Q", "epoch.C", "epoch^4")), var.labels = vlab[c(1,
  4, 2, 3)])
```




Figure 12: Pairs HE plot of Skulls data, showing multivariate tests of `epoch`, as well as tests of linear and nonlinear trends.

These plots have an interesting geometric interpretation: the  $\mathbf{H}$  ellipses for the overall effect of `epoch` are representations of the additive decomposition of this effect into  $\mathbf{H}$  ellipses for the linear and nonlinear linear hypothesis tests according to

$$\mathbf{H}_{\text{epoch}} = \mathbf{H}_{\text{linear}} + \mathbf{H}_{\text{nonlinear}}$$

where the linear term has rank 1 (and so plots as a line), while the nonlinear term has rank 3. In each panel, it can be seen that the large direction of the  $\mathbf{H}_{\text{epoch}}$  leading to significance of this effect corresponds essentially to the linear contrast.  $\mathbf{H}_{\text{nonlinear}}$  is the orthogonal complement of  $\mathbf{H}_{\text{linear}}$  in the space of  $\mathbf{H}_{\text{epoch}}$ , but nowhere does it protrude beyond the boundary of the  $\mathbf{E}$  ellipsoid.



These relations can be seen somewhat more easily in 3D, as produced using `heplot3d()` by the following statement. The resulting plot is better viewed interactively in R and is not reproduced here. It would be seen there that the ellipsoid for the nonlinear terms is nearly flat in one direction, corresponding to the panel for (mb, hb) in Figure 12.

```
> heplot3d(sk.mod, hypotheses=list(Lin="epoch.L", Quad="epoch.Q",
                                   NonLin=c("epoch.Q", "epoch.C", "epoch^4")),
           col=c("pink", "blue"))
```

Finally, a simpler view of these effects can be shown in the canonical space corresponding to the canonical discriminant analysis for `epoch`. The computations are performed by `candisc()` which returns a class "candisc" object.

```
> library(candisc)
> sk.can <- candisc(sk.mod)
> sk.can
```

Canonical Discriminant Analysis for epoch:

	CanRsq	Eigenvalue	Difference	Percent	Cumulative
1	0.29829	0.42510	0.386	88.227	88.2
2	0.03754	0.03900	0.386	8.094	96.3
3	0.01546	0.01570	0.386	3.259	99.6
4	0.00202	0.00202	0.386	0.419	100.0

Test of H0: The canonical correlations in the current row and all that follow are zero

	LR test stat	approx F	num Df	den Df	Pr(> F)
1	0.664	3.90	16	434	7e-07 ***
2	0.946	0.90	9	348	0.53
3	0.983	0.64	4	288	0.64
4	0.998	0.29	1	145	0.59

---  
Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

The output above shows that, although  $H_{\text{epoch}}$  is of rank 4, two dimensions account for 96% of the between-epoch variation. By the likelihood ratio test, only the canonical correlation for the first dimension can be considered non-zero.

The canonical HE plot is produced by plotting the `sk.can` object, giving Figure 13.

```
> heplot(sk.can, prefix="Canonical dimension")
```

In this plot, the first canonical dimension (88%) essentially corresponds to the ordered values of `epoch`, from right to left. The variable vectors for maximum breadth (mb), basialveolar length (bl) and nasal height (nh) are largely aligned with this dimension, indicating that they distinguish the groups of skulls according to time. The lengths of these vectors indicates their relative contribution to discrimination among the group means. Only the variable vector for basibregmatic height (bh) points in the direction of the second canonical dimension, corresponding to higher means in the middle of the range of epochs. We leave it to forensic anthropologists to determine if this has any meaning.

## 2 Multivariate Multiple Regression Designs

The ideas behind HE plots extend naturally to multivariate multiple regression (MMRA) and multivariate analysis of covariance (MANCOVA). In MMRA, the  $\mathbf{X}$  matrix contains only quantitative predictors, while in MANCOVA designs, there is a mixture of factors and quantitative predictors (covariates).




Figure 13: Canonical discriminant HE plot of the Skulls data

In the MANCOVA case, there is often a subtle difference in emphasis: true MANCOVA analyses focus on the differences among groups defined by the factors, adjusting for (or controlling for) the quantitative covariates. Analyses concerned with *homogeneity of regression* focus on quantitative predictors and attempt to test whether the regression relations are the same for all groups defined by the factors.

## 2.1 Rohwer data

To illustrate the homogeneity of regression flavor, we use data from a study by Rohwer (given in Timm, 1975: Ex. 4.3, 4.7, and 4.23) on kindergarten children, designed to determine how well a set of paired-associate (PA) tasks predicted performance on the Peabody Picture Vocabulary test (PPVT), a student achievement test (SAT), and the Raven Progressive matrices test (Raven). The PA tasks varied in how the stimuli were presented, and are called *named* (n), *still* (s), *named still* (ns), *named action* (na), and *sentence still* (ss).

Two groups were tested: a group of  $n = 37$  children from a low socioeconomic status (SES) school, and a group of  $n = 32$  high SES children from an upper-class, white residential school. The data are in the data frame `Rohwer` in the `heplots` package:

```
> some(Rohwer, n=6)
```

	group	SES	SAT	PPVT	Raven	n	s	ns	na	ss
14	1	Lo	30	55	13	2	1	12	20	17
18	1	Lo	45	54	10	0	6	6	14	16
34	1	Lo	19	55	8	0	1	14	19	12
37	1	Lo	79	54	14	0	6	6	15	14
58	2	Hi	50	82	18	4	5	16	21	24
67	2	Hi	98	96	18	4	7	10	23	19

At one extreme, we might be tempted to fit separate regression models for each of the High and Low SES groups. This approach is *not* recommended because it lacks power and does not allow hypotheses about equality of slopes and intercepts to be tested directly.

```
> rohwer.ses1 <- lm(cbind(SAT, PPVT, Raven) ~ n + s + ns + na + ss, data=Rohwer, subset=SES=="Hi")
> Anova(rohwer.ses1)
```

```
Type II MANOVA Tests: Pillai test statistic
Df test stat approx F num Df den Df Pr(>F)
n 1 0.202 2.02 3 24 0.1376
s 1 0.310 3.59 3 24 0.0284 *
ns 1 0.358 4.46 3 24 0.0126 *
na 1 0.465 6.96 3 24 0.0016 **
ss 1 0.089 0.78 3 24 0.5173
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
> rohwer.ses2 <- lm(cbind(SAT, PPVT, Raven) ~ n + s + ns + na + ss, data=Rohwer, subset=SES=="Lo")
> Anova(rohwer.ses2)
```

```
Type II MANOVA Tests: Pillai test statistic
Df test stat approx F num Df den Df Pr(>F)
n 1 0.0384 0.39 3 29 0.764
s 1 0.1118 1.22 3 29 0.321
ns 1 0.2252 2.81 3 29 0.057 .
na 1 0.2675 3.53 3 29 0.027 *
ss 1 0.1390 1.56 3 29 0.220
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

This allows separate slopes and intercepts for each of the two groups, but it is difficult to compare the coefficients numerically.

```
> coef(rohwer.ses1)
```

	SAT	PPVT	Raven
(Intercept)	-28.46747	39.697090	13.243836
n	3.25713	0.067283	0.059347
s	2.99658	0.369984	0.492444
ns	-5.85906	-0.374380	-0.164022
na	5.66622	1.523009	0.118980
ss	-0.62265	0.410157	-0.121156

```
> coef(rohwer.ses2)
```

	SAT	PPVT	Raven
(Intercept)	4.151060	33.005769	11.173378
n	-0.608872	-0.080567	0.210995
s	-0.050156	-0.721050	0.064567
ns	-1.732395	-0.298303	0.213584
na	0.494565	1.470418	-0.037318
ss	2.247721	0.323965	-0.052143

Nevertheless, we can visualize the results with HE plots, and here we make use of the fact that several HE plots can be overlaid using the option `add=TRUE` as shown in Figure 14.




Figure 14: HE plot for SAT and PPVT, showing the effects for the PA predictors for the High and Low SES groups separately

```
> heplot(rohwer.ses1, ylim=c(40,110),col=c("red", "black"), lwd=2, cex=1.2)
> heplot(rohwer.ses2, add=TRUE, col=c("blue", "black"), grand.mean=TRUE, error.ellipse=TRUE, lwd=2, cex=1.2)
> means <- aggregate(cbind(SAT,PPVT)~SES, data=Rohwer, mean)
> text(means[,2], means[,3], labels=means[,1], pos=3, cex=2, col=c("red", "blue"))
```

We can readily see the difference in means for the two SES groups (High greater on both variables) and it also appears that the slopes of the predictor ellipses are shallower for the High than the Low group, indicating greater relation with the SAT score.

Alternatively (and optimistically), we can fit a MANCOVA model that allows different means for the two SES groups on the responses, but constrains the slopes for the PA covariates to be equal.

```
> # MANCOVA, assuming equal slopes
> rohwer.mod <- lm(cbind(SAT, PPVT, Raven) ~ SES + n + s + ns + na + ss,
  data=Rohwer)
> Anova(rohwer.mod)
```

```
Type II MANOVA Tests: Pillai test statistic
Df test stat approx F num Df den Df Pr(>F)
SES 1 0.379 12.18 3 60 2.5e-06 ***
n 1 0.040 0.84 3 60 0.4773
s 1 0.093 2.04 3 60 0.1173
ns 1 0.193 4.78 3 60 0.0047 **
na 1 0.231 6.02 3 60 0.0012 **
ss 1 0.050 1.05 3 60 0.3770
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Note that, although the multivariate tests for two of the covariates (**ns** and **na**) are highly significant, univariate multiple regression tests for the separate responses [from `summary(rohwer.mod)`] are relatively weak. We can also test the global 5 df hypothesis,  $\mathbf{B} = \mathbf{0}$ , that *all* covariates have null effects for all responses as a linear hypothesis (suppressing display of the error and hypothesis SSP matrices),

```
> (covariates <- rownames(coef(rohwer.mod))[-(1:2)])

[1] "n" "s" "ns" "na" "ss"

> Reqr<-linearHypothesis(rohwer.mod, covariates)
> print(Reqr, digits=5, SSP=FALSE)

Multivariate Tests:
              Df test stat approx F num Df den Df    Pr(>F)
Pillai        5   0.66579   3.5369    15 186.00 2.309e-05 ***
Wilks         5   0.44179   3.8118    15 166.03 8.275e-06 ***
Hotelling-Lawley 5   1.03094   4.0321    15 176.00 2.787e-06 ***
Roy           5   0.75745   9.3924     5  62.00 1.062e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Then 2D views of the additive MANCOVA model `rohwer.mod` and the overall test for all covariates are produced as follows, giving the plots in Figure 15.

```
> colors <- c("red", "blue", rep("black",5), "#969696")
> heplot(rohwer.mod, col=colors,
        hypotheses=list("Regr" = c("n", "s", "ns", "na", "ss")),
        cex=1.5, lwd=c(2, rep(3,5), 4),
        main="(SAT, PPVT, Raven) ~ SES + n + s + ns + na + ss")

> heplot(rohwer.mod, col=colors, variables=c(1,3),
        hypotheses=list("Regr" = c("n", "s", "ns", "na", "ss")),
        cex=1.5, lwd=c(2, rep(3,5), 4),
        main="(SAT, PPVT, Raven) ~ SES + n + s + ns + na + ss")
```




Figure 15: HE plot for SAT and PPVT (left) and for SAT and Raven (right) using the MANCOVA model

The positive orientation of the `Regr` ellipses shows that the predicted values for all three responses are positively correlated (more so for SAT and PPVT). As well, the High SES group is higher on all responses than the Low SES group.

Alternatively, all pairwise plots among these responses could be drawn using the `pairs` function (figure not shown),

```
> pairs(rohwer.mod, col=colors,
        hypotheses=list("Regr" = c("n", "s", "ns", "na", "ss")),
        cex=1.3, lwd=c(2, rep(3,5), 4))
```

or as a 3D plot, using `heplot3d()` as shown in Figure 16.

```
> colors <- c("pink", "blue", rep("black",5), "#969696")
> heplot3d(rohwer.mod, col=colors,
           hypotheses=list("Regr" = c("n", "s", "ns", "na", "ss")))
```




Figure 16: 3D HE plot for the MANCOVA model fit to the Rohwer data

The MANCOVA model, `rohwer.mod`, has relatively simple interpretations (large effect of SES, with `ns` and `na` as the major predictors) but the test of relies on the assumption of homogeneity of slopes for the predictors. We can test this as follows, adding interactions of SES with each of the covariates:

```
> rohwer.mod2 <- lm(cbind(SAT, PPVT, Raven) ~ SES * (n + s + ns + na + ss),
                    data=Rohwer)
> Anova(rohwer.mod2)
```

```
Type II MANOVA Tests: Pillai test statistic
      Df test stat approx F num Df den Df Pr(>F)
SES    1    0.391    11.78     3    55 4.5e-06 ***
n      1    0.079     1.57     3    55 0.20638 .
s      1    0.125     2.62     3    55 0.05952 .
ns     1    0.254     6.25     3    55 0.00100 ***
na     1    0.307     8.11     3    55 0.00015 ***
ss     1    0.060     1.17     3    55 0.32813
SES:n   1    0.072     1.43     3    55 0.24417
SES:s   1    0.099     2.02     3    55 0.12117
SES:ns  1    0.118     2.44     3    55 0.07383 .
SES:na  1    0.148     3.18     3    55 0.03081 *
SES:ss  1    0.057     1.12     3    55 0.35094
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

It appears from the above that there is only weak evidence of unequal slopes from the separate `SES:` terms. The evidence for heterogeneity is stronger, however, when these terms are tested collectively using the `linearHypothesis()` function:

```

> (coefs <- rownames(coef(rohwer.mod2)))

[1] "(Intercept)" "SESLo"      "n"      "s"      "ns"
[6] "na"          "ss"       "SESLo:n" "SESLo:s" "SESLo:ns"
[11] "SESLo:na"    "SESLo:ss"

> print(linearHypothesis(rohwer.mod2, coefs[grep(":", coefs)]), SSP=FALSE)

Multivariate Tests:
              Df test stat approx F num Df den Df    Pr(>F)
Pillai        5   0.41794   1.8452    15 171.00 0.032086 *
Wilks         5   0.62358   1.8936    15 152.23 0.027695 *
Hotelling-Lawley 5   0.53865   1.9272    15 161.00 0.023962 *
Roy           5   0.38465   4.3850     5  57.00 0.001905 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

This model (`rohwer.mod2`) is similar in spirit to the two models (`rohwer.ses1` and `rohwer.ses2`) fit for the two SES groups separately and show in Figure 14, except that model `rohwer.mod2` assumes a common within-groups error covariance matrix and allows overall tests.

To illustrate model `rohwer.mod2`, we construct an HE plot for SAT and PPVT shown in Figure 17. To simplify this display, we show the hypothesis ellipses for the overall effects of the PA tests in the baseline high-SES group, and a single combined ellipse for all the `SESLo:` interaction terms that we tested previously, representing differences in slopes between the low and high-SES groups.

Because SES is “treatment-coded” in this model, the ellipse for each covariate represents the hypothesis that the slopes for that covariate are zero in the high-SES baseline category. With this parameterization, the ellipse for `Slopes` represents the joint hypothesis that slopes for the covariates do not differ in the low-SES group.

```

> colors <- c("red", "blue", rep("black",5), "#969696")
> heplot(rohwer.mod2, col=c(colors, "brown"),
        terms=c("SES", "n", "s", "ns", "na", "ss"),
        hypotheses=list("Regr" = c("n", "s", "ns", "na", "ss"),
                        "Slopes" = coefs[grep(":", coefs)]))

```

Comparing Figure 17 for the heterogeneous slopes model with Figure 15 (left) for the homogeneous slopes model, it can be seen that most of the covariates have ellipses of similar size and orientation, reflecting similar evidence against the respective null hypotheses, as does the effect of `SES`, showing the greater performance of the high-SES group on all response measures. Somewhat more subtle, the error ellipse is noticeably smaller in Figure 17, reflecting the additional variation accounted for by differences in slopes.

## 2.2 Recovery from hernia repair

This example uses the `Hernior` data, comprising data on measures of post-operative recovery of 32 patients undergoing an elective herniorrhaphy operation, in relation to pre-operative measures.

The response measures are the patient’s condition upon leaving the recovery room (`leave`, a 1-4 scale, 1=best), level of nursing required one week after operation (`nurse`, a 1-5 scale, 1=worst) and length of stay in hospital after operation (`los`, in days)




Figure 17: HE plot for SAT and PPVT, fitting the model `rohwer.mod2` that allows unequal slopes for the covariates.

The predictor variables are patient `age`, `sex`, physical status (`pstat`, a 1-5 scale, with 1=perfect health, 5=very poor health), body build (`build`, a 1-5 scale, with 1=emaciated, ..., 5=obese), and preoperative complications with (`cardiac`) heart and respiration (`resp`).

We begin with a model fitting all predictors. Note that the ordinal predictors, `pstat`, `build`, `cardiac` and `resp` could arguably be treated as factors, rather than linear, regression terms. We ignore this possibility in this example.

```
> Hern.mod <- lm(cbind(leave, nurse, los) ~ age + sex + pstat + build + cardiac + resp,
                  data=Hernior)
> Anova(Hern.mod)
```

```
Type II MANOVA Tests: Pillai test statistic
      Df test stat approx F num Df den Df Pr(>F)
age      1      0.143      1.27      3      23 0.307
sex      1      0.026      0.21      3      23 0.892
pstat    1      0.333      3.84      3      23 0.023 *
build    1      0.257      2.65      3      23 0.073 .
cardiac   1      0.228      2.26      3      23 0.108 .
resp     1      0.248      2.53      3      23 0.082 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The results of the multivariate tests above are somewhat disappointing. Only the physical status predictor (`pstat`) appears to be significant at conventional levels.

The univariate models for each response are implicit in the MLM `Hern.mod`. These can be printed using `summary()`, or we can use `summary()` to extract certain statistics for each univariate response model, as we do here.

```
> Hern.summary <- summary(Hern.mod)
> unlist(lapply(Hern.summary, function(x) x$r.squared))
```

```
Response leave Response nurse Response los
0.59176      0.24740      0.36531
```



The univariate tests for predictors in each of these models (not shown here) are hard to interpret, and largely show only significant effects for the `leave` variable. Yet, the  $R^2$  values for the other responses are slightly promising. We proceed to an overall test of  $\mathbf{B} = 0$  for all predictors.

```
> # test overall regression
> Regr <- linearHypothesis(Hern.mod, c("age", "sexm", "pstat", "build", "cardiac", "resp"))
> print(Regr, digits=5, SSP=FALSE)
```

Multivariate Tests:

	Df	test stat	approx F	num Df	den Df	Pr(>F)
Pillai	6	1.10198	2.4192	18	75.000	0.0041356 **
Wilks	6	0.21734	2.6046	18	65.539	0.0025239 **
Hotelling-Lawley	6	2.26797	2.7300	18	65.000	0.0016285 **
Roy	6	1.55434	6.4764	6	25.000	0.0003232 ***

---  
Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

```
> clr <- c("red", "darkgray", "blue", "darkgreen", "magenta", "brown", "black")
> vlab <- c("LeaveCondition\n(leave)", "NursingCare\n(nurse)", "LengthOfStay\n(los)")
> hyp <- list("Regr" = c("age", "sexm", "pstat", "build", "cardiac", "resp"))
> pairs(Hern.mod, hypotheses=hyp, col=clr, var.labels=vlab)
```

A `pairs()` plot for the MLM gives the set of plots shown in Figure 18 helps to interpret the relations among the predictors which lead to the highly significant overall test. Among the predictors, age and sex have small and insignificant effects on the outcome measures jointly. The other predictors all contribute to the overall test of  $\mathbf{B} = 0$ , though in different ways for the various responses. For example, in the panel for (`leave`, `los`) in Figure 18, it can be seen that while only `pstat` individually is outside the  $\mathbf{E}$  ellipse, `build` and `resp` contribute to the overall test in an opposite direction.

In this multivariate regression example, all of the terms in the model `Hern.mod` have 1 df, and so plot as lines in HE plots. An alternative view of these effects can be seen in canonical discriminant space, which, for each predictor shows the scores on the linear combination of the responses that contributes most to the multivariate test of that effect, together with the weights for the responses. We use `candiscList()` to calculate the canonical analyses for all terms in `Hern.mod`.

```
> Hern.canL <- candiscList(Hern.mod)
```

1D canonical discriminant plots for all terms can be obtained interactively with a menu, simply by plotting the `Hern.canL` object.

```
> plot(Hern.canL)
```

Plots for separate terms are produced by the lines below, and shown in Figure 19 and Figure 20.

```
> plot(Hern.canL, term="pstat")
> plot(Hern.canL, term="build")
> plot(Hern.canL, term="age")
> plot(Hern.canL, term="cardiac")
```

In these plots, the canonical scores panel shows the linear combinations of the response variables which have the largest possible  $R^2$ . Better outcomes correspond to numerically smaller canonical scores. The arrows in the structure panel are proportional to the canonical weights.

These plots provide simple interpretations of the results for the canonical combinations of the responses. Better physical status, smaller body build, lower age and absence of cardiac complications are all positively related to better outcomes.




Figure 18: HE pairs plot for Hernior data

## 2.3 Grades in a Sociology Course

The data set `SocGrades` contains four outcome measures on student performance in an introductory sociology course together with six potential predictors. These data were used by Marascuilo and Levin (1983) for an example of canonical correlation analysis, but are also suitable as examples of multivariate multiple regression, MANOVA, MANCOVA and step-down analysis in multivariate linear models.

The outcome measures used here are three test scores during the course, `midterm1`, `midterm2`, `final`, and a course evaluation (`eval`).<sup>2</sup> Predictor variables are student's social class (`class`, an ordered factor with levels 1 > 2 > 3) `sex`, grade point average (`gpa`), College Board test scores (`boards`), previous high school unit in sociology? (`hssoc`: `no`, `yes`), and score on a course pretest (`pretest`).

The basic MLM is fit below as `grades.mod`.

```
> data(SocGrades)
> grades.mod <- lm(cbind(midterm1, midterm2, final, eval) ~
```

<sup>2</sup>It is arguable that the students' course evaluation should not be considered a response variable here. It could be used as a predictor in a follow-up, step-down analysis, which would address the separate question of whether the effects on exam grades remain, when `eval` is controlled for.




Figure 19: 1D Canonical discriminant plots for `pstat` and `build`. The canonical scores are such that better outcomes are associated with smaller scores.




Figure 20: 1D Canonical discriminant plots for `age` and `cardiac`.

```
class + sex + gpa + boards + hssoc + pretest, data=SocGrades)
> Anova(grades.mod, test="Roy")
```

```
Type II MANOVA Tests: Roy test statistic
Df test stat approx F num Df den Df Pr(>F)
class    2    1.567    11.75     4    30 7.3e-06 ***
sex       1    0.553     4.01     4    29 0.0104 *
gpa       1    1.208     8.76     4    29 9.2e-05 ***
boards    1    0.731     5.30     4    29 0.0025 **
hssoc     1    0.035     0.25     4    29 0.9052 .
pretest   1    0.313     2.27     4    29 0.0859 .
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

In both univariate and multivariate response models, it is often useful to screen for higher-order terms (interactions, non-linear predictors). This can most easily be done using `update()`, as we do below. First, try the extended model with all pairwise interactions of the predictors.

```
> grades.mod2 <- update(grades.mod, . ~ .^2)
> Anova(grades.mod2, test="Roy")
```

```
Type II MANOVA Tests: Roy test statistic
      Df test stat approx F num Df den Df Pr(>F)
class      2      2.817      7.04      4      10 0.0058 **
sex         1      0.487      1.09      4       9 0.4152
gpa         1      1.998      4.49      4       9 0.0286 *
boards      1      2.338      5.26      4       9 0.0183 *
hssoc       1      0.281      0.63      4       9 0.6522
pretest     1      0.510      1.15      4       9 0.3946
class:sex    2      2.039      5.10      4      10 0.0168 *
class:gpa    2      0.982      2.45      4      10 0.1137
class:boards 2      0.522      1.31      4      10 0.3321
class:hssoc   2      0.356      0.89      4      10 0.5041
class:pretest 2      1.005      2.51      4      10 0.1082
sex:gpa       1      0.269      0.60      4       9 0.6694
sex:boards    1      0.184      0.41      4       9 0.7944
sex:hssoc     1      0.909      2.04      4       9 0.1714
sex:pretest   1      0.885      1.99      4       9 0.1795
gpa:boards    1      0.447      1.00      4       9 0.4537
gpa:hssoc     1      0.596      1.34      4       9 0.3269
gpa:pretest   1      0.472      1.06      4       9 0.4291
boards:hssoc  1      0.353      0.80      4       9 0.5573
boards:pretest 1      0.705      1.59      4       9 0.2593
hssoc:pretest 1      1.464      3.29      4       9 0.0635 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

In the results above, only the interaction of `class:sex` is significant, and the main effects of `hssoc` and `pretest` remain insignificant. A revised model to explore is `grades.mod3`,

```
> grades.mod3 <- update(grades.mod, . ~ . + class:sex - hssoc - pretest)
> Anova(grades.mod3, test="Roy")
```

```
Type II MANOVA Tests: Roy test statistic
      Df test stat approx F num Df den Df Pr(>F)
class      2      1.588     11.91      4      30 6.5e-06 ***
sex         1      0.575      4.17      4      29 0.00864 **
gpa         1      1.434     10.40      4      29 2.4e-05 ***
boards      1      0.895      6.49      4      29 0.00074 ***
class:sex    2      0.450      3.38      4      30 0.02143 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

A pairwise HE plot for all responses (Figure 21) shows that nearly all effects are in the expected directions: higher `gpa`, `boards`, `class` leads to better performance on most outcomes. The interaction of `class:sex` seems to be confined largely to `midterm1`.

```
> pairs(grades.mod3)
```

These effects are easier to appreciate for the three exam grades jointly in a 3D HE plot. A snapshot is shown in Figure 22.

```
> heplot3d(grades.mod3, wire=FALSE)
```




Figure 21: HE pairs plot for SocGrades

Interactive rotation of this plot shows that the effect of `class` is only two dimensional, and of these, one is very small. The major axis of the `class` ellipsoid is aligned with increasing performance on all three grades, with the expected ordering of the three social classes.

The representation of these effects in canonical space is particularly useful here. Again, use `candiscList()` to compute the canonical decompositions for all terms in the model, and extract the canonical  $R^2$  from the terms in the result.

```
> # calculate canonical results for all terms
> grades.can <- candiscList(grades.mod3)
> # extract canonical R^2s
> unlist(lapply(grades.can, function(x) x$canrsq))
```

```
class1    class2    sex    gpa    boards class:sex1 class:sex2
0.613620  0.024186  0.365269  0.589145  0.472268  0.310461  0.132931
```

We use `heplot()` on the "candiscList" object to show the effects of `class` in canonical space, giving Figure 23.

```
> # plot class effect in canonical space
> op <- par(xpd=TRUE)
> heplot(grades.can, term="class", scale=4, fill=TRUE, var.col="black", var.lwd=2)
> par(op)
```




Figure 22: 3D HE plot for SocGrades, model `grades.mod3`

It can be seen in Figure 23 that nearly all variation in exam performance due to class is aligned with the first canonical dimension. The three tests and course evaluation all have similar weights on this dimension, but the course evaluation differs from the rest along a second, very small dimension.

1D plots of the canonical scores for other effects in the model are also of interest, and provide simple interpretations of these effects on the response variables. The statements below produce the plots shown in Figure 24.

```
> plot(grades.can, term="sex")
> plot(grades.can, term="gpa")
```

It is readily seen that males perform better overall, but the effect of `sex` is strongest for the `midterm2`. As well, increasing course performance on tests is strongly associated with `gpa`.

## References

- J. Fox, M. Friendly, and G. Monette. Visualizing hypothesis tests in multivariate linear models: The *heplots* package for R. *Computational Statistics*, 24(2):233–246, 2009. (Published online: 15 May 2008).
- M. Friendly. Data ellipses, HE plots and reduced-rank displays for multivariate linear models: SAS software and examples. *Journal of Statistical Software*, 17(6):1–42, 2006.
- M. Friendly. HE plots for multivariate general linear models. *Journal of Computational and Graphical Statistics*, 16(2):421–444, 2007.




Figure 23: Canonical HE plot for `class` effect in `grades.mod3`




Figure 24: 1D Canonical discriminant plots for `sex` and `gpa`. Higher canonical scores reflect better course performance.

- M. E. Plaster. *The Effect of Defendent Physical Attractiveness on Juridic Decisions Using Felon Inmates as Mock Jurors*. Unpublished master's thesis, East Carolina University, Greenville, NC, 1989.
- N. H. Timm. *Multivariate Analysis with Applications in Education and Psychology*. Wadsworth (Brooks/Cole), Belmont, CA, 1975.