

Analyse avancée

Distribution theory

Question 1/15

$\text{supp}(\alpha)$ for $\alpha \in \mathcal{D}'(O)$

Réponse 1/15

The complementary of the greatest set of
 $\mathcal{Z} = \{U \subseteq O \text{ open}, \forall u \in \mathcal{D}(U), \langle \alpha, u \rangle = 0\}$

It is the complementary of the biggest set on
which $\alpha = 0$

Question 2/15

Order of a distribution with compact support

Réponse 2/15

The order is finite

Question 3/15

\mathcal{C}^∞ Urysohn lemma

Réponse 3/15

If K is compact and O is open with $K \subseteq O \subseteq \mathbb{R}^d$ then there exists a smooth function χ such that $0 \leq \chi \leq 1$, $\chi \equiv 1$ on K and $\text{supp}(\chi) \subseteq O$

It is denoted $K \prec \chi \prec O$

Question 4/15

$\alpha \in \mathcal{D}'(O)$ is of order at most j on K

$\alpha \in \mathcal{D}'(O)$ is of order at most j

$\alpha \in \mathcal{D}'(O)$ is of order j on K

$\alpha \in \mathcal{D}'(O)$ is of order j

Réponse 4/15

There exists $C > 0$ such that, for all

$$u \in \mathcal{D}(O), |\langle \alpha, u \rangle| \leq C p_{K,j}(u)$$

α is of order at most j on all compact $K \subseteq O$

α is of order at most j but not of order at most

$$j - 1 \text{ on } K$$

The order of α is the minimal j for which α is
of order at most j

Question 5/15

Properties of the topological space $\mathcal{D}(O)$

Réponse 5/15

$\mathcal{D}(O)$ is an LCTVS

Question 6/15

$f\alpha$ for $f \in \mathcal{C}^\infty(O)$ and $\alpha \in \mathcal{D}'(O)$

Réponse 6/15

$\beta \in \mathcal{D}'(O)$ such that, for all $u \in \mathcal{D}(O)$,

$$\langle \beta, u \rangle = \langle \alpha, fu \rangle$$

If α is of order at most j then β is of order at most j and $\text{supp}(\beta) \subseteq \text{supp}(f) \cap \text{supp}(\alpha)$

Question 7/15

Topology on $\mathcal{D}_K(O) = \mathcal{C}_K^\infty(O)$, $K \subseteq O$
compact

Réponse 7/15

Topology generated by the semi-norms

$$p_{K,j} = \sup(\partial_x^m u(x), |m| \leq j, x \in K)$$

Question 8/15

Class of test functions on $O \subseteq \mathbb{R}^d$ open

Réponse 8/15

$$\mathcal{D}(O) = \mathcal{C}_c^\infty(O)$$

Question 9/15

Taylor expansion in \mathbb{R}^d

Réponse 9/15

If $u \in \mathcal{D}(O)$ and $\overline{\mathcal{B}}(0, r) \subseteq O$, there exists $C > 0$ such that for all $x \in \overline{\mathcal{B}}(0, r)$,

$$\left| u(x) - \sum_{|m| \leq j} \frac{x^m}{m!} \partial_x^m u(0) \right| \leq C|x|^{j+1}$$

Question 10/15

$$\partial_x^m \alpha \text{ for } \alpha \in \mathcal{D}'(O)$$

Réponse 10/15

$\beta \in \mathcal{D}'(O)$ such that, for all $u \in \mathcal{D}(O)$,

$$\langle \beta, u \rangle = (-1)^m \langle \alpha, \partial_x^m u \rangle$$

Question 11/15

$\tau_z \alpha$ for $z \in \mathbb{R}^d$ and $\alpha \in \mathcal{D}'(\mathbb{R}^d)$

Réponse 11/15

$\beta \in \mathcal{D}'(\mathbb{R}^d)$ such that, for all $u \in \mathcal{D}(\mathbb{R}^d)$,
 $\langle \beta, u \rangle = \langle \alpha, \tau_{-z}u \rangle$ where $\tau_y u(x) = u(x - y)$

Question 12/15

$\alpha * f(x)$ for $f \in \mathcal{C}^\infty(\mathbb{R}^d)$ and $\alpha \in \mathcal{D}'(\mathbb{R}^d)$

Réponse 12/15

$\langle \alpha, \tau_x \check{f} \rangle$ where $\tau_y u(x) = u(x - y)$ and
 $\check{u}(x) = u(-x)$

Question 13/15

Leibniz formula in \mathbb{R}^d

Réponse 13/15

If $\chi, u \in \mathcal{D}(\mathbb{R}^d)$, $\partial_x^m(\chi u) = \sum_{p+q=m} \binom{m}{p} \partial_x^p \chi \partial_x^q u$

where $\binom{m}{p} = \frac{m!}{p!q!}$ where $p + q = m$ and

$$(m_1, \dots, m_d)! = m_1! \cdots m_d!$$

Question 14/15

CNS for a distribution having a punctual support

Réponse 14/15

$\text{supp}(\alpha) = \{z\}$ and α is of order at most j iff
for all $u \in \mathcal{D}(O)$ such that $|m| \leq j$, if
 $\partial_x^m u(z) = 0$ then $\langle \alpha, u \rangle = 0$, iff there exists
 $(C_m)_{|m| \leq j}$ such that $\alpha = \sum_{|m| \leq j} C_m \partial_x^m \delta_z$

Question 15/15

Space of distributions

Réponse 15/15

Topological dual $\mathcal{D}'(O)$ of $\mathcal{D}(O)$ endowed with
the weak-star topology