# Crash course on entanglement theory

Ray Ganardi

2023.11.24

CeNT, University of Warsaw, Poland

State	Expectation value
$ \psi angle$	$ra{\psi}X\ket{\psi}$

State	Expectation value
$ \psi angle$	$\langle \psi   X   \psi \rangle$
$\{(p_i, \psi_i\rangle)\}$	$\sum_{i} p_{i} \langle \psi_{i}   X   \psi_{i} \rangle = \operatorname{Tr} \left( \left( \sum_{i} p_{i}   \psi_{i} \rangle \langle \psi_{i}   \right) X \right)$


State	Expectation value
$ \psi\rangle$	$\langle \psi   X   \psi \rangle$
$\{(p_i, \psi_i\rangle)\}$	$\sum_{i} p_{i} \langle \psi_{i}   X   \psi_{i} \rangle = \operatorname{Tr} \left( \left( \sum_{i} p_{i}   \psi_{i} \rangle \langle \psi_{i}   \right) X \right)$
$\rho = \sum_{i} p_{i}  \psi_{i}\rangle\langle\psi_{i} $	$\operatorname{Tr}\left( ho X ight)$

State	Expectation value
$ \psi angle$	$\langle \psi   X   \psi \rangle$
	$\sum_{i} p_{i} \langle \psi_{i}   X   \psi_{i} \rangle = \operatorname{Tr} \left( \left( \sum_{i} p_{i}   \psi_{i} \rangle \langle \psi_{i}   \right) X \right)$
$\rho = \sum_{i} p_i  \psi_i\rangle\langle\psi_i $	$\operatorname{Tr}\left( ho X ight)$


A (mixed) state is represented by a density matrix  $\rho$  s.t.

- 1.  $\rho \ge 0$
- 2. Tr  $\rho = 1$

 $\mathcal{D} \subset \mathcal{M}$  the set of all density matrices




We could say  $\rho_{AB}=\rho_A\otimes\rho_B$ 



We could say  $\rho_{AB}=\rho_A\otimes\rho_B$ 

Only makes sense if the systems are uncorrelated




We could say  $\rho_{AB} = \rho_A \otimes \rho_B$ 

Only makes sense if the systems are uncorrelated

#### Example

$$\rho_{AB} = \frac{1}{2} |00\rangle\langle00| + \frac{1}{2} |11\rangle\langle11|, \text{ while } \rho_A \otimes \rho_B = \frac{1}{2} \otimes \frac{1}{2}$$

3



We could say  $\rho_{AB} = \rho_A \otimes \rho_B$ 

Only makes sense if the systems are uncorrelated

#### Example

$$\rho_{AB}=\frac{1}{2}\left|00\right\rangle\!\!\left\langle 00\right|+\frac{1}{2}\left|11\right\rangle\!\!\left\langle 11\right|$$
 , while  $\rho_{A}\otimes\rho_{B}=\frac{1}{2}\otimes\frac{1}{2}$ 

In general,  $\rho_{AB}=\sum_{i}p_{i}\left|\psi_{i}\right\rangle\!\langle\psi_{i}|_{AB}$ , where  $\left|\psi_{i}\right\rangle_{AB}\in\mathcal{H}_{A}\otimes\mathcal{H}_{B}$ 

# What is entanglement?

# What is entanglement?

non-classical correlation

What is entanglement?


non-classical correlation

What is classical correlation?

LOCC: local operations and classical communication




LOCC: local operations and classical communication



Separable states:  $ho_{AB} = \sum_i p_i 
ho_A^i \otimes 
ho_B^i$ 


LOCC: local operations and classical communication



Separable states:  $ho_{AB} = \sum_i p_i 
ho_A^i \otimes 
ho_B^i$ 

Entangled states: everything else

LOCC: local operations and classical communication



Separable states:  $ho_{AB} = \sum_i p_i 
ho_A^i \otimes 
ho_B^i$ 

Entangled states: everything else

Proposition

A pure state  $\psi_{AB}$  is separable iff.  $\psi_{AB}=\psi_{A}\otimes\psi_{B}$ 

# Entanglement theory

Transformations between entangled states under LOCC

#### Bipartite pure states

Proposition (Schmidt decomposition)

Given a pure state  $\psi_{AB}$ , there exist bases  $\{|e_i\rangle_A\}, \{|f_j\rangle_B\}$  such that

$$|\psi_{AB}\rangle = \sum_{i} \sqrt{p_{i}} |e_{i}\rangle_{A} \otimes |f_{i}\rangle_{B}$$

#### Bipartite pure states

Proposition (Schmidt decomposition)

Given a pure state  $\psi_{AB}$ , there exist bases  $\{|e_i\rangle_A\},\{|f_j\rangle_B\}$  such that

$$|\psi_{AB}\rangle = \sum_{i} \sqrt{p_i} |e_i\rangle_A \otimes |f_i\rangle_B$$

Theorem (Nielsen's majorization)

$$\begin{array}{l} (|\psi\rangle = \sum_{i} \sqrt{p_{i}} \, |ii\rangle) \rightarrow \left(|\phi\rangle = \sum_{j} \sqrt{q_{j}} \, |jj\rangle\right) \text{ iff.} \\ \sum_{i < k} p_{i} \leq \sum_{j < k} q_{j} \text{ for all } k \text{ (also written } \vec{p} \preceq \vec{q}\text{)} \end{array}$$

7

## Bipartite pure states

Proposition (Schmidt decomposition)

Given a pure state  $\psi_{AB}$ , there exist bases  $\{|e_i\rangle_A\},\{|f_j\rangle_B\}$  such that

$$|\psi_{AB}\rangle = \sum_{i} \sqrt{p_i} |e_i\rangle_A \otimes |f_i\rangle_B$$

Theorem (Nielsen's majorization)

$$\begin{array}{l} (|\psi\rangle = \sum_{i} \sqrt{p_{i}} \, |ii\rangle) \rightarrow \left(|\phi\rangle = \sum_{j} \sqrt{q_{j}} \, |jj\rangle\right) \text{ iff.} \\ \sum_{i < k} p_{i} \leq \sum_{j < k} q_{j} \text{ for all } k \text{ (also written } \vec{p} \preceq \vec{q}\text{)} \end{array}$$

#### Example

$$|\Phi\rangle=\frac{1}{\sqrt{2}}\,|00\rangle+\frac{1}{\sqrt{2}}\,|11\rangle$$
 can be transformed to any pure two qubit state  $|\phi\rangle=\sqrt{q_0}\,|00\rangle+\sqrt{q_1}\,|11\rangle$ 

$$\psi \not\to \Phi \text{ since} \\ (0.6,0.4) \not\preceq (0.5,0.5)$$

$$\psi \not\rightarrow \Phi$$
 since  $(0.6, 0.4) \not \leq (0.5, 0.5)$ 

$$\psi^{\otimes 2} \to \Phi \text{ since} \\ (0.36, 0.24, 0.24, 0.16) \preceq (0.5, 0.5)$$

$$\psi \not\rightarrow \Phi$$
 since  $(0.6, 0.4) \not \leq (0.5, 0.5)$ 

$$\psi^{\otimes 2} \to \Phi \text{ since} \\ (0.36, 0.24, 0.24, 0.16) \preceq (0.5, 0.5)$$

$$\begin{array}{l} \psi^{\otimes 3} \rightarrow \Phi^{\otimes 2} \text{ since} \\ (0.216, 0.144, 0.144, 0.144, 0.096, 0.096, 0.096, 0.064) \preceq \\ (0.25, 0.25, 0.25, 0.25) \end{array}$$

What's the maximum rate of distillation?

What's the maximum rate of distillation?

Distillable entanglement

$$E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$$

What's the maximum rate of distillation?

Distillable entanglement

$$E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$$

Example

$$E_D\!\left(\sqrt{0.6}\left|00\right\rangle+\sqrt{0.4}\left|11\right\rangle\right)\geq 2/3$$

What's the maximum rate of distillation?

Distillable entanglement

$$E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$$

Example

$$E_D\!\left(\sqrt{0.6}\left|00\right\rangle+\sqrt{0.4}\left|11\right\rangle\right)\geq 2/3$$

Distillable state:  $E_D(\rho) > 0$ 

Bound entangled state:  $\rho$  entangled but  $E_D(\rho) = 0$ 

# **Asymptotic transformations**

$$R(\rho \to \sigma) = \sup \left\{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \sigma^{\otimes m} \right\}$$

# **Asymptotic transformations**

$$R(\rho \to \sigma) = \sup \left\{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \sigma^{\otimes m} \right\}$$

Theorem
For pure states,

$$R(\psi \to \phi) = \frac{S(\psi_A)}{S(\phi_A)}$$

# Summary

entanglement and LOCC

End of part 1

# **Summary**

- entanglement and LOCC
- Nielsen's majorization

End of part 1

# **Summary**

- entanglement and LOCC
- Nielsen's majorization
- entanglement distillation

End of part 1

# Catalytic and asymptotic equivalence for quantum entanglement


Can your bank help you to distill more entanglement? arxiv:2305.03488

Ray Ganardi

2023.11.24

CeNT, University of Warsaw, Poland Joint work with Tulja Varun Kondra and Alexander Streltsov

# Entanglement




# **Entanglement**



Separable states:  $\rho = \sum_i p_i \rho_A^i \otimes \rho_B^i$ 


# **Entanglement**



Separable states:  $\rho = \sum_i p_i \rho_A^i \otimes \rho_B^i$ 

Entangled states: not separable

# **Entanglement**



Separable states:  $ho = \sum_i p_i 
ho_A^i \otimes 
ho_B^i$ 

Entangled states: not separable

Entanglement is useful

Real-life source produces noisy entanglement

Real-life source produces noisy entanglement

Many copies: purify with entanglement distillation



Real-life source produces noisy entanglement

Many copies: purify with entanglement distillation



Distillable entanglement: maximal distillation rate under LOCC

$$E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$$

Real-life source produces noisy entanglement

Many copies: purify with entanglement distillation



Distillable entanglement: maximal distillation rate under LOCC

$$E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$$

Distillable state:  $E_D(\rho) > 0$ 

Bound entangled state:  $\rho$  entangled but  $E_D(\rho)=0$ 

Why limit ourselves to LOCC?

Why limit ourselves to LOCC?

What if we can "borrow" entanglement?

## **Catalysis**

definition (exact)

$$\rho \xrightarrow{ec} \sigma$$
 if there exists a catalyst state  $\tau$  and an LOCC protocol  $\Lambda$  s.t.  $\Lambda(\rho \otimes \tau) = \sigma \otimes \tau$ 

### **Catalysis**

definition (exact)

$$\rho \xrightarrow{ec} \sigma$$
 if there exists a catalyst state  $\tau$  and an LOCC protocol  $\Lambda$  s.t.  $\Lambda(\rho \otimes \tau) = \sigma \otimes \tau$ 

How powerful is this?

### **Catalysis**

definition (exact)

$$\rho \xrightarrow{ec} \sigma$$
 if there exists a catalyst state  $\tau$  and an LOCC protocol  $\Lambda$  s.t.  $\Lambda(\rho \otimes \tau) = \sigma \otimes \tau$ 

How powerful is this?

Can we find  $\rho, \sigma$  such that  $\rho \not\to \sigma$  but  $\rho \xrightarrow{ec} \sigma$ ?

Jonathan, Plenio (PRL 1999): yes

Jonathan, Plenio (PRL 1999): yes

Single-shot bipartite pure state transformation is governed by majorization (Nielsen, PRL 1999)

$$\sum_i \sqrt{p_i} \, |ii\rangle \to \sum_j \sqrt{q_j} \, |jj\rangle$$
 iff.  $\sum_{i < k} p_i \le \sum_{j < k} q_j$  for all  $k$ 

Jonathan, Plenio (PRL 1999): yes

Single-shot bipartite pure state transformation is governed by majorization (Nielsen, PRL 1999)

$$\sum_i \sqrt{p_i} \ket{ii} \to \sum_j \sqrt{q_j} \ket{jj}$$
 iff.  $\sum_{i < k} p_i \le \sum_{j < k} q_j$  for all  $k$ 

Choose

$$s(\psi) = (0.4, 0.4, 0.1, 0.1)$$
  
$$s(\phi) = (0.5, 0.25, 0.25)$$

Jonathan, Plenio (PRL 1999): yes

Single-shot bipartite pure state transformation is governed by majorization (Nielsen, PRL 1999)

$$\sum_i \sqrt{p_i} |ii\rangle \to \sum_j \sqrt{q_j} |jj\rangle$$
 iff.  $\sum_{i < k} p_i \le \sum_{j < k} q_j$  for all  $k$  Choose

$$s(\psi) = (0.4, 0.4, 0.1, 0.1)$$
  

$$s(\phi) = (0.5, 0.25, 0.25)$$
  

$$s(\tau) = (0.6, 0.4)$$

Jonathan, Plenio (PRL 1999): yes

Single-shot bipartite pure state transformation is governed by majorization (Nielsen, PRL 1999)

$$\sum_i \sqrt{p_i} \, |ii\rangle \to \sum_j \sqrt{q_j} \, |jj\rangle$$
 iff.  $\sum_{i < k} p_i \le \sum_{j < k} q_j$  for all  $k$ 

Choose

$$s(\psi) = (0.4, 0.4, 0.1, 0.1)$$
  

$$s(\phi) = (0.5, 0.25, 0.25)$$
  

$$s(\tau) = (0.6, 0.4)$$

Then

$$s(\psi \otimes \tau) = (0.24, 0.24, 0.16, 0.16, 0.06, 0.06, 0.04, 0.04)$$
  
$$s(\phi \otimes \tau) = (0.30, 0.20, 0.15, 0.15, 0.10, 0.10, 0.00, 0.00)$$

• Duan (PRA 2005): connection with asymptotics

■ Duan (PRA 2005): connection with asymptotics If  $\psi^{\otimes n} \to \phi^{\otimes n}$  for some n, then  $\psi \xrightarrow{ec} \phi$ . catalyst:  $|\tau\rangle = \sum_{i < n-1} |\psi\rangle^{\otimes i} \otimes |\phi\rangle^{\otimes n-1-i} \otimes |i\rangle / \sqrt{n}$ 

- Duan (PRA 2005): connection with asymptotics If  $\psi^{\otimes n} \to \phi^{\otimes n}$  for some n, then  $\psi \xrightarrow{ec} \phi$ . catalyst:  $|\tau\rangle = \sum_{i < n-1} |\psi\rangle^{\otimes i} \otimes |\phi\rangle^{\otimes n-1-i} \otimes |i\rangle / \sqrt{n}$
- definition (correlated)  $\rho \xrightarrow{cc} \sigma \text{ if for any } \epsilon > 0 \text{, there exists a catalyst } \tau \text{ and an LOCC}$  protocol  $\Lambda$  s.t.

$$\mu_{SC} = \Lambda(\rho_S \otimes \tau_C),$$
 
$$\|\mu_S - \sigma\|_1 \le \epsilon, \ \mu_C = \tau.$$

- Duan (PRA 2005): connection with asymptotics If  $\psi^{\otimes n} \to \phi^{\otimes n}$  for some n, then  $\psi \xrightarrow{ec} \phi$ . catalyst:  $|\tau\rangle = \sum_{i < n-1} |\psi\rangle^{\otimes i} \otimes |\phi\rangle^{\otimes n-1-i} \otimes |i\rangle / \sqrt{n}$
- definition (correlated)  $\rho \xrightarrow{cc} \sigma \text{ if for any } \epsilon > 0 \text{, there exists a catalyst } \tau \text{ and an LOCC}$  protocol  $\Lambda$  s.t.

• Kondra (PRL 2021):  $\psi \xrightarrow{cc} \phi$  iff.  $S(\psi_A) \geq S(\phi_A)$ .

- Duan (PRA 2005): connection with asymptotics If  $\psi^{\otimes n} \to \phi^{\otimes n}$  for some n, then  $\psi \xrightarrow{ec} \phi$ . catalyst:  $|\tau\rangle = \sum_{i < n-1} |\psi\rangle^{\otimes i} \otimes |\phi\rangle^{\otimes n-1-i} \otimes |i\rangle / \sqrt{n}$
- definition (correlated)  $\rho \xrightarrow{cc} \sigma \text{ if for any } \epsilon > 0 \text{, there exists a catalyst } \tau \text{ and an LOCC}$  protocol  $\Lambda$  s.t.

• Kondra (PRL 2021):  $\psi \xrightarrow{cc} \phi$  iff.  $S(\psi_A) \geq S(\phi_A)$ . Catalytic-asymptotic equivalence (for bipartite pure states)!

- Duan (PRA 2005): connection with asymptotics If  $\psi^{\otimes n} \to \phi^{\otimes n}$  for some n, then  $\psi \xrightarrow{ec} \phi$ . catalyst:  $|\tau\rangle = \sum_{i < n-1} |\psi\rangle^{\otimes i} \otimes |\phi\rangle^{\otimes n-1-i} \otimes |i\rangle / \sqrt{n}$
- definition (correlated)  $\rho \xrightarrow{cc} \sigma \text{ if for any } \epsilon > 0 \text{, there exists a catalyst } \tau \text{ and an LOCC}$  protocol  $\Lambda$  s.t.

$$\mu_{SC} = \Lambda(\rho_S \otimes \tau_C),$$
 
$$\|\mu_S - \sigma\|_1 \le \epsilon, \ \mu_C = \tau.$$

- Kondra (PRL 2021):  $\psi \xrightarrow{cc} \phi$  iff.  $S(\psi_A) \geq S(\phi_A)$ . Catalytic-asymptotic equivalence (for bipartite pure states)!
- Duan construction can be extended to mixed states

How powerful is catalysis?


How powerful is catalysis?


Literature: at least as powerful as asymptotics


How powerful is catalysis?

Literature: at least as powerful as asymptotics


This work: exactly as powerful as asymptotics\*\*







For bipartite pure states,  $\psi \xrightarrow{a} \phi$  iff.  $\psi \xrightarrow{ma} \phi$ 



#### **Theorem**

When the initial state is distillable, correlated catalysis is equivalent to marginal asymptotics

#### **Theorem**

When the initial state is distillable, correlated catalysis is equivalent to marginal asymptotics

Proof sketch:

Catalysis  $\leftarrow$  marginal asymptotics: Duan construction

Catalysis  $\Rightarrow$  marginal asymptotics: use distillability to create the catalyst from the initial state

#### **Theorem**

When the initial state is distillable, correlated catalysis is equivalent to marginal asymptotics

Proof sketch:

Catalysis  $\leftarrow$  marginal asymptotics: Duan construction

Catalysis  $\Rightarrow$  marginal asymptotics: use distillability to create the catalyst from the initial state

For two-qubit states, this shows general equivalence!

#### Theorem

When the initial state is distillable, correlated catalysis is equivalent to marginal asymptotics

Proof sketch:

Catalysis  $\leftarrow$  marginal asymptotics: Duan construction

Catalysis  $\Rightarrow$  marginal asymptotics: use distillability to create the catalyst from the initial state

For two-qubit states, this shows general equivalence!

Also works in the multipartite setting

Distillable entanglement:  $E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$ 

Distillable entanglement:  $E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$ 

#### **Theorem**

Distillable entanglement for a distillable state cannot increase under correlated catalysis

Distillable entanglement:  $E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$ 

#### **Theorem**

Distillable entanglement for a distillable state cannot increase under correlated catalysis

#### Proof sketch:

ullet standard rate  $\leq$  marginal rate = marginal catalytic rate

Distillable entanglement:  $E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$ 

#### Theorem

Distillable entanglement for a distillable state cannot increase under correlated catalysis

#### Proof sketch:

- standard rate ≤ marginal rate = marginal catalytic rate
- ullet standard rate  $\leq$  catalytic rate  $\leq$  marginal catalytic rate

Distillable entanglement:  $E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$ 

#### Theorem

Distillable entanglement for a distillable state cannot increase under correlated catalysis

#### Proof sketch:

- standard rate ≤ marginal rate = marginal catalytic rate
- ullet standard rate  $\leq$  catalytic rate  $\leq$  marginal catalytic rate
- marginal rate = standard rate (for pure target state)

Distillable entanglement:  $E_D(\rho) = \sup \{ m/n \, | \, \Lambda(\rho^{\otimes n}) \approx \Phi^{\otimes m} \}$ 

#### Theorem

Distillable entanglement for a distillable state cannot increase under correlated catalysis

#### Proof sketch:

- standard rate ≤ marginal rate = marginal catalytic rate
- standard rate  $\leq$  catalytic rate  $\leq$  marginal catalytic rate
- marginal rate = standard rate (for pure target state)

Lami (arxiv 2023): There exist bound entangled states that cannot be distilled under catalysis

 Catalysis is exactly as powerful as marginal asymptotics for distillable states

- Catalysis is exactly as powerful as marginal asymptotics for distillable states
- Catalysis cannot increase distillable entanglement

- Catalysis is exactly as powerful as marginal asymptotics for distillable states
- Catalysis cannot increase distillable entanglement

### Open questions

Equivalence for bound entangled state

- Catalysis is exactly as powerful as marginal asymptotics for distillable states
- Catalysis cannot increase distillable entanglement

#### Open questions

- Equivalence for bound entangled state
- Gap between marginal asymptotic and standard asymptotic

- Catalysis is exactly as powerful as marginal asymptotics for distillable states
- Catalysis cannot increase distillable entanglement

#### Open questions

- Equivalence for bound entangled state
- Gap between marginal asymptotic and standard asymptotic

#### Thanks!

arxiv:2305.03488