

Coherence manipulation in asymmetry and thermodynamics

arxiv:2308.12814


Ray Ganardi

2023.12.14

CeNT, University of Warsaw, Poland

Joint work with Tulja Varun Kondra and Alexander Streltsov

What is thermodynamics?



study of state transformations

states: (ρ, H)

Resource theory of thermodynamics

states: (ρ, H)

free state: Gibbs state

free operations: thermal operations

$$\Lambda(\rho_S) = \text{Tr}_E \left[U(\rho_S \otimes \gamma_E) U^\dagger \right]$$

with $[U, H_S + H_E] = 0$

Resource theory of thermodynamics

states: (ρ, H)

free state: Gibbs state

free operations: thermal operations

$$\Lambda(\rho_S) = \text{Tr}_E \left[U(\rho_S \otimes \gamma_E) U^\dagger \right]$$

with $[U, H_S + H_E] = 0$

state transformations under thermal operations? hard

Catalytic transformations

definition (exact catalysis)


$$\rho \xrightarrow[ec]{} \sigma \text{ if } \exists \tau, \Lambda \text{ s.t. } \Lambda(\rho \otimes \tau) = \sigma \otimes \tau$$

$$\begin{array}{cc} \rho \bullet & \bullet \sigma \\ \tau \bullet & \bullet \tau \end{array} \rightarrow$$

Catalytic transformations

definition (exact catalysis)

$$\rho \xrightarrow[ec]{} \sigma \text{ if } \exists \tau, \Lambda \text{ s.t. } \Lambda(\rho \otimes \tau) = \sigma \otimes \tau$$



(Brandão, PNAS 2015) characterization

For ρ, σ incoherent, $\rho \xrightarrow[ec]{} \sigma$ iff. $S_\alpha(\rho \parallel \gamma) \geq S_\alpha(\sigma \parallel \gamma)$

Correlated catalysis

definition (correlated catalysis)

$$\rho \xrightarrow{cc} \sigma \text{ if } \forall \epsilon > 0, \exists \tau, \Lambda \text{ s.t.}$$

$$\mu_{SC} = \Lambda(\rho_S \otimes \tau_C)$$

$$\|\mu_S - \sigma\|_1 \leq \epsilon, \mu_C = \tau$$

Correlated catalysis

definition (correlated catalysis)

$$\rho \xrightarrow{cc} \sigma \text{ if } \forall \epsilon > 0, \exists \tau, \Lambda \text{ s.t.}$$

$$\mu_{SC} = \Lambda(\rho_S \otimes \tau_C)$$

$$\|\mu_S - \sigma\|_1 \leq \epsilon, \mu_C = \tau$$

(Müller, PRX 2018) characterization

For ρ, σ incoherent states, $\rho \xrightarrow{cc} \sigma$ iff. $S(\rho||\gamma) \geq S(\sigma||\gamma)$

What about transformations with coherence?

Thermal operations

$$\Lambda(\rho_S) = \text{Tr}_E \left[U(\rho_S \otimes \gamma_E) U^\dagger \right]$$

with $[U, H_S + H_E] = 0$

Covariant operations

Thermal operations

$$\Lambda(\rho_S) = \text{Tr}_E \left[U(\rho_S \otimes \gamma_E) U^\dagger \right]$$

with $[U, H_S + H_E] = 0$

(Lostaglio, Nat Comms 2015): they are covariant!

$$\Lambda\left(e^{-iH_S t} \rho e^{iH_S t}\right) = e^{-iH_S t} \Lambda(\rho) e^{iH_S t}$$

Covariant operations

Thermal operations

$$\Lambda(\rho_S) = \text{Tr}_E \left[U(\rho_S \otimes \gamma_E) U^\dagger \right]$$

with $[U, H_S + H_E] = 0$

(Lostaglio, Nat Comms 2015): they are covariant!

$$\Lambda\left(e^{-iH_S t} \rho e^{iH_S t}\right) = e^{-iH_S t} \Lambda(\rho) e^{iH_S t}$$

Covariant operations

$$\Lambda(\rho) = \text{Tr}_E \left[U(\rho_S \otimes \sigma_E) U^\dagger \right]$$

with $[U, H_S + H_E] = 0$ and σ incoherent

symmetry: $\{U_g \mid g \in \mathcal{G}\}$

Asymmetry

symmetry: $\{U_g \mid g \in \mathcal{G}\}$

free states

incoherent states

$$\sigma = U_g \sigma U_g^\dagger$$

Asymmetry

symmetry: $\{U_g \mid g \in \mathcal{G}\}$

free states

incoherent states

$$\sigma = U_g \sigma U_g^\dagger$$

free operations

covariant operations

$$\Lambda(U_g \rho U_g^\dagger) = U_g \Lambda(\rho) U_g^\dagger$$

Asymmetry

symmetry: e^{-iHt}

free states

incoherent states

$$\sigma = \sum_i p_i |i\rangle\langle i|$$

free operations

covariant operations

$$\Lambda(\rho) = \text{Tr}_E \left[U(\rho_S \otimes \sigma_E) U^\dagger \right]$$

with $[U, H_S + H_E] = 0$ and σ incoherent

Asymmetry

symmetry: e^{-iHt}

free states

incoherent states

$$\sigma = \sum_i p_i |i\rangle\langle i|$$

free operations

covariant operations

$$\Lambda(\rho) = \text{Tr}_E \left[U(\rho_S \otimes \sigma_E) U^\dagger \right]$$

with $[U, H_S + H_E] = 0$ and σ incoherent

captures coherence limitations in thermodynamics

Result

definition (available/reachable coherence)

$$\mathcal{I}(\rho) = \{\Delta_{ij} = E_i - E_j \mid \langle i | \rho | j \rangle\}$$

$$\mathcal{J}(\rho) = \bigcup_n \mathcal{I}(\rho^{\otimes n}) = \left\{ \sum_{\Delta_{ij} \in \mathcal{I}(\rho)} m_{ij} \Delta_{ij} \right\}$$

Result

definition (available/reachable coherence)

$$\mathcal{I}(\rho) = \{\Delta_{ij} = E_i - E_j \mid \langle i | \rho | j \rangle\}$$

$$\mathcal{J}(\rho) = \bigcup_n \mathcal{I}(\rho^{\otimes n}) = \left\{ \sum_{\Delta_{ij} \in \mathcal{I}(\rho)} m_{ij} \Delta_{ij} \right\}$$

theorem

$$\rho \xrightarrow{cc} \sigma \text{ if } \mathcal{J}(\rho) \supseteq \mathcal{J}(\sigma)$$

definition (available/reachable coherence)

$$\mathcal{I}(\rho) = \{\Delta_{ij} = E_i - E_j \mid \langle i | \rho | j \rangle\}$$
$$\mathcal{J}(\rho) = \bigcup_n \mathcal{I}(\rho^{\otimes n}) = \left\{ \sum_{\Delta_{ij} \in \mathcal{I}(\rho)} m_{ij} \Delta_{ij} \right\}$$

theorem

$$\rho \xrightarrow{cc} \sigma \text{ if } \mathcal{J}(\rho) \supseteq \mathcal{J}(\sigma)$$

consequences

- coherence amplification
- zero or infinite distillation rate

What is known (Takagi, PRL 2022)

1. Given enough copies of $|+_{ij}\rangle$ for all $\Delta_{ij} \in \mathcal{I}(\rho)$, we can create ρ by covariant operations

What is known (Takagi, PRL 2022)

1. Given enough copies of $|+_{ij}\rangle$ for all $\Delta_{ij} \in \mathcal{I}(\rho)$, we can create ρ by covariant operations
2. There exists sequence of correlated catalytic transformations that create arbitrarily many copies of $|+_{ij}\rangle$ if $\Delta_{ij} \in \mathcal{J}(\rho)$

What is known (Takagi, PRL 2022)

1. Given enough copies of $|+_{ij}\rangle$ for all $\Delta_{ij} \in \mathcal{I}(\rho)$, we can create ρ by covariant operations
2. There exists sequence of correlated catalytic transformations that create arbitrarily many copies of $|+_{ij}\rangle$ if $\Delta_{ij} \in \mathcal{J}(\rho)$

Just need to show that we can merge this sequence into one

Proof sketch

What we showed


We can merge the sequence into one big catalytic transformation

Proof sketch

What we showed

We can merge the sequence into one big catalytic transformation

tool: marginal reducibility



talk more in entanglement session

- idea: use catalytic-asymptotic equivalence for asymmetry (Ganardi, 2023)

- idea: use catalytic-asymptotic equivalence for asymmetry (Ganardi, 2023)
- does not work because of bound coherence states

- idea: use catalytic-asymptotic equivalence for asymmetry (Ganardi, 2023)
- does not work because of bound coherence states
- marginal reducibility \Rightarrow catalysis is good (Duan state)

- idea: use catalytic-asymptotic equivalence for asymmetry (Ganardi, 2023)
- does not work because of bound coherence states
- marginal reducibility \Rightarrow catalysis is good (Duan state)
- catalysis \Rightarrow marginal reducibility for two-level systems

- idea: use catalytic-asymptotic equivalence for asymmetry (Ganardi, 2023)
- does not work because of bound coherence states
- marginal reducibility \Rightarrow catalysis is good (Duan state)
- catalysis \Rightarrow marginal reducibility for two-level systems
- marginal reducibility is transitive

What is known (Takagi, PRL 2022)

1. Given enough copies of $|+_{ij}\rangle$ for all $\Delta_{ij} \in \mathcal{I}(\rho)$, we can create ρ by covariant operations
2. There exists sequence of correlated catalytic transformations that create arbitrarily many copies of $|+_{ij}\rangle$ if $\Delta_{ij} \in \mathcal{J}(\rho)$

We can merge this sequence into one big catalytic transformation!

conjecture

$$\rho \xrightarrow{cc} \sigma \text{ iff. } S(\rho||\gamma) \geq S(\sigma||\gamma) \text{ and } \mathcal{J}(\rho) \supseteq \mathcal{J}(\sigma)$$

conjecture

$$\rho \xrightarrow{cc} \sigma \text{ iff. } S(\rho\|\gamma) \geq S(\sigma\|\gamma) \text{ and } \mathcal{J}(\rho) \supseteq \mathcal{J}(\sigma)$$

why/how?

- For Gibbs-preserving operations, $\rho \xrightarrow{cc} \sigma$ iff. $S(\rho\|\gamma) \geq S(\sigma\|\gamma)$
(Shiraishi, PRL 2021)

conjecture

$$\rho \xrightarrow{cc} \sigma \text{ iff. } S(\rho\|\gamma) \geq S(\sigma\|\gamma) \text{ and } \mathcal{I}(\rho) \supseteq \mathcal{I}(\sigma)$$

why/how?

- For Gibbs-preserving operations, $\rho \xrightarrow{cc} \sigma$ iff. $S(\rho\|\gamma) \geq S(\sigma\|\gamma)$ (Shiraishi, PRL 2021)
- For covariant operations, $\rho \xrightarrow{cc} \sigma$ iff. $\mathcal{I}(\rho) \supseteq \mathcal{I}(\sigma)$

conjecture

$$\rho \xrightarrow{cc} \sigma \text{ iff. } S(\rho||\gamma) \geq S(\sigma||\gamma) \text{ and } \mathcal{J}(\rho) \supseteq \mathcal{J}(\sigma)$$

why/how?

- For Gibbs-preserving operations, $\rho \xrightarrow{cc} \sigma$ iff. $S(\rho||\gamma) \geq S(\sigma||\gamma)$ (Shiraishi, PRL 2021)
- For covariant operations, $\rho \xrightarrow{cc} \sigma$ iff. $\mathcal{J}(\rho) \supseteq \mathcal{J}(\sigma)$
- Thermal operations = Gibbs-preserving operations \cap covariant operations

- catalytic coherence manipulation

Summary

- catalytic coherence manipulation
- marginal reducibility/correlated catalysis is transitive

Summary

- catalytic coherence manipulation
- marginal reducibility/correlated catalysis is transitive

open questions

- are $\mathcal{J}(\rho)$ monotones?

Summary

- catalytic coherence manipulation
- marginal reducibility/correlated catalysis is transitive

open questions

- are $\mathcal{J}(\rho)$ monotones?
- coherence amplification in thermodynamics

Summary

- catalytic coherence manipulation
- marginal reducibility/correlated catalysis is transitive

open questions

- are $\mathcal{J}(\rho)$ monotones?
- coherence amplification in thermodynamics
- monotones for thermal operations

Summary

- catalytic coherence manipulation
- marginal reducibility/correlated catalysis is transitive

open questions

- are $\mathcal{J}(\rho)$ monotones?
- coherence amplification in thermodynamics
- monotones for thermal operations

Thanks!

arxiv:2308.12814