Introduction to Complex Numbers

Ray Ganardi

May 20, 2021

IFTiA

University of Gdańsk, Poland

What is it?

$$i^2 = -1$$

(en): complex number, imaginary number

(pl): liczba zespolona, liczba urojona

What is it?

$$i^{2} = -1$$

$$\mathbb{C} = \{(a+bi) \mid a, b \in \mathbb{R}\}$$

(en): complex number, imaginary number

(pl): liczba zespolona, liczba urojona

Rules

$$(a+bi) + (c+di) = (a+c) + (b+d)i$$

Rules

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

 $(a + bi)(c + di) = (ac - bd) + (bc + ad)i$

History

Conventional:

- 1545 Cardano: Introduction, "Ars Magna"
- 1572 Bombelli: Calculation, "L'Algebra"
- 1770 **Euler**: $\sqrt{-2}\sqrt{-3} = \sqrt{6}$
- end of 18th century Wessel, Argand, Gauss: Geometric interpretation
- 1814–1851: Cauchy, Riemann, etc.: Complex analysis

4

Consider the quadratic equation

$$x^2 = mx + c$$

Consider the quadratic equation

$$x^2 = mx + c$$



2000 BC: solutions are

$$x = \frac{1}{2} \left(m \pm \sqrt{m^2 + 4c} \right)$$

Consider the quadratic equation

$$x^2 = mx + c$$


2000 BC: solutions are

$$x = \frac{1}{2} \left(m \pm \sqrt{m^2 + 4c} \right)$$


Cardano: If $m^2 + 4c < 0$, then x must be imaginary.




Problem is geometrical




Problem is geometrical



if $m^2 + 4c < 0$, then there is no intersection




Consider the cubic equation $x^3 = 3px + 2q$.



7

Consider the cubic equation $x^3 = 3px + 2q$.



Cardano: Solution is


$$x = \sqrt[3]{q + \sqrt{q^2 - p^3}} + \sqrt[3]{q - \sqrt{q^2 - p^3}}$$

$$x^{3} = 3px + 2q$$

$$x = \sqrt[3]{q + \sqrt{q^{2} - p^{3}}} + \sqrt[3]{q - \sqrt{q^{2} - p^{3}}}$$

Suppose $p^3 > q^2$.

For example $x^3 = 15x + 4$.



$$x^3 = 15x + 4$$

Cardano: solution is

$$x = \sqrt[3]{2 + 11i} + \sqrt[3]{2 - 11i}$$



$$x^3 = 15x + 4$$

Cardano: solution is

$$x = \sqrt[3]{2 + 11i} + \sqrt[3]{2 - 11i}$$

Inspection shows that x=4 is a solution.

$$x^3 = 15x + 4$$

Cardano: solution is

$$x = \sqrt[3]{2 + 11i} + \sqrt[3]{2 - 11i}$$

Inspection shows that x=4 is a solution.

Bombelli: What if $\sqrt[3]{2 \pm 11i} = 2 \pm ni$ for some n?



Assuming the multiplication rule,

$$(2 \pm ni)^3 = 2 \pm 11i$$

Assuming the multiplication rule,

$$(2 \pm ni)^3 = 2 \pm 11i$$

But it's easy to show that

$$(2 \pm i)^3 = 2 \pm 11i$$

Assuming the multiplication rule,

$$(2 \pm ni)^3 = 2 \pm 11i$$

But it's easy to show that

$$(2 \pm i)^3 = 2 \pm 11i$$


The formula $x=\sqrt[3]{q+\sqrt{q^2-p^3}}+\sqrt[3]{q-\sqrt{q^2-p^3}}$ works if we interpret $\sqrt{q^2-p^3}=\left(\sqrt{p^3-q^2}\right)i$.

Geometric interpretation (addition)

$$(a+bi) + (c+di) = (a+c) + (b+d)i$$

Geometric interpretation (addition)

$$(a+bi) + (c+di) = (a+c) + (b+d)i$$



$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$


Notice that multiplication by i is a "rotation".

$$(a+bi)\,i = -b + ai$$

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

Notice that multiplication by i is a "rotation".

$$(a+bi)\,i = -b + ai$$



$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$


Multiplication by a real number is a "scaling".

$$(a+bi) r = ra + rbi$$

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

Multiplication by a real number is a "scaling".

$$(a+bi) r = ra + rbi$$



$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

$$|a+bi| = \sqrt{a^2 + b^2}$$

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

$$|a+bi| = \sqrt{a^2 + b^2}$$

$$|(a+bi)(c+di)| = \sqrt{(ac-bd)^2 + (bc+ad)^2}$$

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

$$|a+bi| = \sqrt{a^2 + b^2}$$

$$|(a+bi)(c+di)| = \sqrt{(ac-bd)^2 + (bc+ad)^2}$$

= $\sqrt{(ac)^2 + (bd)^2 + (bc)^2 + (ad)^2}$

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

$$|a+bi| = \sqrt{a^2 + b^2}$$

$$|(a+bi)(c+di)| = \sqrt{(ac-bd)^2 + (bc+ad)^2}$$

$$= \sqrt{(ac)^2 + (bd)^2 + (bc)^2 + (ad)^2}$$

$$= \sqrt{(a^2 + b^2)(c^2 + d^2)}$$

$$(a+bi)(c+di) = (ac-bd) + (bc+ad)i$$

$$|a+bi| = \sqrt{a^2 + b^2}$$

$$|(a+bi)(c+di)| = \sqrt{(ac-bd)^2 + (bc+ad)^2}$$

$$= \sqrt{(ac)^2 + (bd)^2 + (bc)^2 + (ad)^2}$$

$$= \sqrt{(a^2 + b^2)(c^2 + d^2)}$$

$$= |a+bi||c+di|$$

Perhaps for every complex number z = a + bi,

$$z = abs(z) arg(z)$$
scaling rotation

Perhaps for every complex number z = a + bi,

$$z = abs(z) arg(z)$$

Multiplication by a real number gives a "pure" scaling.

Perhaps for every complex number z = a + bi,

$$z = abs(z) arg(z)$$
 scaling rotation

Multiplication by a real number gives a "pure" scaling.

Is there a "pure" rotation?


Consider multiplication by $z = \cos \theta + i \sin \theta$.

Consider multiplication by $z = \cos \theta + i \sin \theta$.

$$(a+bi)(\cos\theta+i\sin\theta)=(a\cos\theta-b\sin\theta)+i(b\cos\theta+a\sin\theta)$$

Consider multiplication by $z = \cos \theta + i \sin \theta$.

$$(a+bi)(\cos\theta + i\sin\theta) = (a\cos\theta - b\sin\theta) + i(b\cos\theta + a\sin\theta)$$



Geometric interpretation

For every $z=a+bi\in\mathbb{C}$, we have

$$z = |z| (\cos \theta + i \sin \theta),$$

where

$$|z| = \sqrt{a^2 + b^2}$$
$$\theta = \arctan \frac{b}{a}$$

Happily ever after

Thanks!

References:

- Needham, Visual Complex Analysis
- Stillwell, Mathematics and Its History

Questions?