

Quaternary Golay Sequence Pairs

Richard Gibson

Department of Mathematics
Simon Fraser University

Masters Thesis Defence
November 6, 2008

① Background

Outline

- ① Background
- ② Classifying Quaternary Golay Sequence Pairs

- ① Background
- ② Classifying Quaternary Golay Sequence Pairs
- ③ Constructing a Binary Barker Sequence from a Quaternary Golay Sequence

- ① Background
- ② Classifying Quaternary Golay Sequence Pairs
- ③ Constructing a Binary Barker Sequence from a Quaternary Golay Sequence
- ④ Summary and Open Problems

Background

Sequences

Example

0	1	2	0	3
---	---	---	---	---

quaternary sequence

Sequences

Example

0	1	2	0	3
---	---	---	---	---

quaternary sequence

$A = (a_0, \dots, a_{n-1})$ is a **quaternary sequence** if $a_j \in \mathbb{Z}_4$ for all $0 \leq j < n$.

Sequences

Example

0	1	2	0	3
---	---	---	---	---

quaternary sequence

$A = (a_0, \dots, a_{n-1})$ is a **quaternary sequence** if $a_j \in \mathbb{Z}_4$ for all $0 \leq j < n$.

Example

0	0	0	0	0	2	2	0	0	2	0	2	0
---	---	---	---	---	---	---	---	---	---	---	---	---

binary sequence

Sequences

Example

0	1	2	0	3
---	---	---	---	---

quaternary sequence

$A = (a_0, \dots, a_{n-1})$ is a **quaternary sequence** if $a_j \in \mathbb{Z}_4$ for all $0 \leq j < n$.

Example

0	0	0	0	0	2	2	0	0	2	0	2	0
---	---	---	---	---	---	---	---	---	---	---	---	---

binary sequence

$A = (a_0, \dots, a_{n-1})$ is a **binary sequence** if $a_j \in \{0, 2\}$ for all $0 \leq j < n$.

Aperiodic autocorrelation function

Define $C_A(u)$, the **aperiodic autocorrelation function** of the quaternary sequence A at shift u , by:

Aperiodic autocorrelation function

Define $C_A(u)$, the **aperiodic autocorrelation function** of the quaternary sequence A at shift u , by:

$$A = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

Aperiodic autocorrelation function

Define $C_A(u)$, the **aperiodic autocorrelation function** of the quaternary sequence A at shift u , by:

$$A = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

$$A = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

Aperiodic autocorrelation function

Define $C_A(u)$, the **aperiodic autocorrelation function** of the quaternary sequence A at shift u , by:

$$\begin{array}{l} A = \\ A = \end{array} \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

$$\begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

Diagram illustrating the aperiodic autocorrelation function $C_A(u)$ for a quaternary sequence A at shift $u=1$. The sequence A is shown as a row of five boxes containing the values 0, 1, 2, 0, 3. Below it, the sequence A is shown again, shifted by one position to the right, resulting in the values 0, 1, 2, 0, 3. Vertical double-headed arrows connect the corresponding elements of the original sequence and the shifted sequence, indicating the shift.

$$C_A(1) = i^{0-1} + i^{1-2} + i^{2-0} + i^{0-3} = -1 - i \text{ (where } i = \sqrt{-1}\text{)}$$

Aperiodic autocorrelation function

Define $C_A(u)$, the **aperiodic autocorrelation function** of the quaternary sequence A at shift u , by:

$$A = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

$$A = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

Aperiodic autocorrelation function

Define $C_A(u)$, the **aperiodic autocorrelation function** of the quaternary sequence A at shift u , by:

$$\begin{array}{rcccl} A = & & \boxed{0} & \boxed{1} & \boxed{2} & \boxed{0} & \boxed{3} \\ & & \updownarrow & \updownarrow & \updownarrow & & \\ A = & \boxed{0} & \boxed{1} & \boxed{2} & \boxed{0} & \boxed{3} \end{array}$$

$$C_A(2) = i^{0-2} + i^{1-0} + i^{2-3} = -1$$

Aperiodic autocorrelation function

Define $C_A(u)$, the **aperiodic autocorrelation function** of the quaternary sequence A at shift u , by:

$$\begin{array}{rccccc} A = & & \boxed{0} & \boxed{1} & \boxed{2} & \boxed{0} & \boxed{3} \\ & & \updownarrow & \updownarrow & \updownarrow & & \\ A = & \boxed{0} & \boxed{1} & \boxed{2} & \boxed{0} & \boxed{3} \end{array}$$

$$C_A(2) = i^{0-2} + i^{1-0} + i^{2-3} = -1$$

$$C_A(u) := \sum_{j=0}^{n-u-1} i^{a_j - a_{j+u}} \text{ for all } 0 \leq u < n$$

What is a Barker sequence?

A quaternary (binary) sequence A of length n is a **quaternary (binary) Barker sequence** if

$$|C_A(u)| \in \{0, 1\} \text{ for all } 1 \leq u < n.$$

What is a Barker sequence?

A quaternary (binary) sequence A of length n is a **quaternary (binary) Barker sequence** if

$$|C_A(u)| \in \{0, 1\} \text{ for all } 1 \leq u < n.$$

Example

$A =$

0	0	0	0	0	2	2	0	0	2	0	2	0
---	---	---	---	---	---	---	---	---	---	---	---	---

u	0	1	2	3	4	5	6	7	8	9	10	11	12
$ C_A(u) $	13	0	1	0	1	0	1	0	1	0	1	0	1

What is a Barker sequence?

A quaternary (binary) sequence A of length n is a **quaternary (binary) Barker sequence** if

$$|C_A(u)| \in \{0, 1\} \text{ for all } 1 \leq u < n.$$

Example

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 2 & 2 & 0 & 0 & 2 & 0 & 2 & 0 \end{bmatrix}$$

u	0	1	2	3	4	5	6	7	8	9	10	11	12
$ C_A(u) $	13	0	1	0	1	0	1	0	1	0	1	0	1

Applications include:

- radar
- pulse compression

History of Barker sequences

- Binary Barker sequences exist for lengths 2, 3, 4, 5, 7, 11, and 13.

History of Barker sequences

- Binary Barker sequences exist for lengths 2, 3, 4, 5, 7, 11, and 13.
- There are no binary Barker sequences of odd length > 13 (Turyn and Storer, 1961).

History of Barker sequences

- Binary Barker sequences exist for lengths 2, 3, 4, 5, 7, 11, and 13.
- There are no binary Barker sequences of odd length > 13 (Turyn and Storer, 1961).
- **Barker Sequence Conjecture**: There are no binary Barker sequences of length > 13 .
 - smallest open case is for length $> 10^{22}$ (Leung and Schmidt, 2005).

What is a Golay sequence?

Let A and B be quaternary (binary) sequences of length n .

What is a Golay sequence?

Let A and B be quaternary (binary) sequences of length n .

A sequence pair (A, B) is a **quaternary (binary) Golay sequence pair** if

$$C_A(u) + C_B(u) = 0 \text{ for all } 1 \leq u < n$$

(we say A and B are **quaternary (binary) Golay sequences**).

What is a Golay sequence?

Let A and B be quaternary (binary) sequences of length n .

A sequence pair (A, B) is a **quaternary (binary) Golay sequence pair** if

$$C_A(u) + C_B(u) = 0 \text{ for all } 1 \leq u < n$$

(we say A and B are **quaternary (binary) Golay sequences**).

Example

$$\begin{array}{l} A = \\ B = \end{array} \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array}$$

u	0	1	2	3	4
$C_A(u)$	5	$-1 - i$	-1	0	i
$C_B(u)$	5	$1 + i$	1	0	$-i$
$C_A(u) + C_B(u)$	10	0	0	0	0

What is a Golay sequence?

Let A and B be quaternary (binary) sequences of length n .

A sequence pair (A, B) is a **quaternary (binary) Golay sequence pair** if

$$C_A(u) + C_B(u) = 0 \text{ for all } 1 \leq u < n$$

(we say A and B are **quaternary (binary) Golay sequences**).

Example

$$\begin{array}{l} A = \\ B = \end{array} \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array}$$

u	0	1	2	3	4
$C_A(u)$	5	$-1 - i$	-1	0	i
$C_B(u)$	5	$1 + i$	1	0	$-i$
$C_A(u) + C_B(u)$	10	0	0	0	0

Applications include:

- medical ultrasound, etc.

History of Golay sequences

- There exist binary Golay pairs of length $2^a 10^b 26^c$ for all integers $a, b, c \geq 0$ (Turyn, 1974).

History of Golay sequences

- There exist binary Golay pairs of length $2^a 10^b 26^c$ for all integers $a, b, c \geq 0$ (Turyn, 1974).
- All binary pairs of length less than 100 have been classified (Borwein and Ferguson, 2003).
 - i.e. each pair either derivable from some general construction method or identified as one of five “seed” pair

History of Golay sequences

- There exist binary Golay pairs of length $2^a 10^b 26^c$ for all integers $a, b, c \geq 0$ (Turyn, 1974).
- All binary pairs of length less than 100 have been classified (Borwein and Ferguson, 2003).
 - i.e. each pair either derivable from some general construction method or identified as one of five “seed” pair
- **Multi-dimensional construction process**: Can obtain Golay pairs from a series of shorter Golay pairs (Fiedler, Jedwab and Parker, 2008).
 - Explains all known binary and quaternary Golay sequences of length 2^m .

History of Golay sequences

- There exist binary Golay pairs of length $2^a 10^b 26^c$ for all integers $a, b, c \geq 0$ (Turyn, 1974).
- All binary pairs of length less than 100 have been classified (Borwein and Ferguson, 2003).
 - i.e. each pair either derivable from some general construction method or identified as one of five “seed” pair
- **Multi-dimensional construction process**: Can obtain Golay pairs from a series of shorter Golay pairs (Fiedler, Jedwab and Parker, 2008).
 - Explains all known binary and quaternary Golay sequences of length 2^m .

There are only two nonexistence results for Golay sequences:

History of Golay sequences

- There exist binary Golay pairs of length $2^a 10^b 26^c$ for all integers $a, b, c \geq 0$ (Turyn, 1974).
- All binary pairs of length less than 100 have been classified (Borwein and Ferguson, 2003).
 - i.e. each pair either derivable from some general construction method or identified as one of five “seed” pair
- **Multi-dimensional construction process**: Can obtain Golay pairs from a series of shorter Golay pairs (Fiedler, Jedwab and Parker, 2008).
 - Explains all known binary and quaternary Golay sequences of length 2^m .

There are only two nonexistence results for Golay sequences:

- Binary Golay sequences must have even length (Marcel Golay, 1961).

History of Golay sequences

- There exist binary Golay pairs of length $2^a 10^b 26^c$ for all integers $a, b, c \geq 0$ (Turyn, 1974).
- All binary pairs of length less than 100 have been classified (Borwein and Ferguson, 2003).
 - i.e. each pair either derivable from some general construction method or identified as one of five “seed” pair
- **Multi-dimensional construction process:** Can obtain Golay pairs from a series of shorter Golay pairs (Fiedler, Jedwab and Parker, 2008).
 - Explains all known binary and quaternary Golay sequences of length 2^m .

There are only two nonexistence results for Golay sequences:

- Binary Golay sequences must have even length (Marcel Golay, 1961).
- The length of a binary Golay pair has no prime factor congruent to 3 modulo 4 (Eliahou, Kervaire and Saffari, 1991).

History of Golay sequences

- There exist binary Golay pairs of length $2^a 10^b 26^c$ for all integers $a, b, c \geq 0$ (Turyn, 1974).
- All binary pairs of length less than 100 have been classified (Borwein and Ferguson, 2003).
 - i.e. each pair either derivable from some general construction method or identified as one of five “seed” pair
- **Multi-dimensional construction process:** Can obtain Golay pairs from a series of shorter Golay pairs (Fiedler, Jedwab and Parker, 2008).
 - Explains all known binary and quaternary Golay sequences of length 2^m .

There are only two nonexistence results for Golay sequences:

- Binary Golay sequences must have even length (Marcel Golay, 1961).
- The length of a binary Golay pair has no prime factor congruent to 3 modulo 4 (Eliahou, Kervaire and Saffari, 1991).

No nonexistence results for quaternary Golay sequences.

Classifying Quaternary Golay Sequence Pairs

Ordered quaternary Golay sequence pair counts

In 2002, Craigen, Holzmann, and Kharaghani exhaustively found all ordered quaternary Golay sequence pairs of small length n :

n	# pairs	n	# pairs	n	# pairs
1	16	8	6656	15	0
2	64	9	0	16	106496
3	128	10	12288	17	0
4	512	11	512	18	24576
5	512	12	36864	19	0
6	2048	13	512	20	215040*
7	0	14	0	21	0

* Frank Fiedler, personal communication.

Ordered quaternary Golay sequence pair counts

In 2002, Craigen, Holzmann, and Kharaghani exhaustively found all ordered quaternary Golay sequence pairs of small length n :

n	# pairs	n	# pairs	n	# pairs
1	16	8	6656	15	0
2	64	9	0	16	106496
3	128	10	12288	17	0
4	512	11	512	18	24576
5	512	12	36864	19	0
6	2048	13	512	20	215040*
7	0	14	0	21	0

- Using small “seed” pairs, how many of the pairs in the table above can be explained using the multi-dimensional construction process and other constructions?

* Frank Fiedler, personal communication.

Ordered quaternary Golay sequence pair counts

In 2002, Craigen, Holzmann, and Kharaghani exhaustively found all ordered quaternary Golay sequence pairs of small length n :

n	# pairs	n	# pairs	n	# pairs
1	16	8	6656	15	0
2	64	9	0	16	106496
3	128	10	12288	17	0
4	512	11	512	18	24576
5	512	12	36864	19	0
6	2048	13	512	20	215040*
7	0	14	0	21	0

- Using small “seed” pairs, how many of the pairs in the table above can be explained using the multi-dimensional construction process and other constructions?
- How can we explain the existence of the seed pairs?

* Frank Fiedler, personal communication.

Multi-dimensional construction process

How do we use the multi-dimensional construction process?

Multi-dimensional construction process

How do we use the multi-dimensional construction process?

- **Input:** $m + 1$ Golay sequence pairs of length n_0, n_1, \dots, n_m , where $m \geq 1$, to create a multi-dimensional object.

Multi-dimensional construction process

How do we use the multi-dimensional construction process?

- **Input:** $m + 1$ Golay sequence pairs of length n_0, n_1, \dots, n_m , where $m \geq 1$, to create a multi-dimensional object.
- Process multi-dimensional object...

Multi-dimensional construction process

How do we use the multi-dimensional construction process?

- **Input:** $m + 1$ Golay sequence pairs of length n_0, n_1, \dots, n_m , where $m \geq 1$, to create a multi-dimensional object.
- Process multi-dimensional object...
- **Output:** A collection of Golay sequence pairs, all of length $n_0 \cdot n_1 \cdot \dots \cdot n_m \cdot 2^m$.

Let's use the multi-dimensional construction process

We can construct ordered quaternary Golay sequence pairs using “seed” Golay pairs of length 1, 3 and 5 as follows:

Let's use the multi-dimensional construction process

We can construct ordered quaternary Golay sequence pairs using “seed” Golay pairs of length 1, 3 and 5 as follows:

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \quad \left. \begin{array}{l} \\ \\ \end{array} \right\} \quad \mapsto \quad \begin{array}{l} \text{64 ordered pairs of} \\ \text{length } 1 \cdot 1 \cdot 2 = 2 \end{array}$$

Ordered quaternary Golay sequence pair counts

In 2002, Craigen, Holzmann, and Kharaghani exhaustively found all ordered quaternary Golay sequence pairs of small length n :

n	# pairs	n	# pairs	n	# pairs
1	16	8	6656	15	0
2	64	9	0	16	106496
3	128	10	12288	17	0
4	512	11	512	18	24576
5	512	12	36864	19	0
6	2048	13	512	20	215040*
7	0	14	0	21	0

* Frank Fiedler, personal communication.

Let's use the multi-dimensional construction process

We can construct ordered quaternary Golay sequence pairs using “seed” Golay pairs of length 1, 3 and 5 as follows:

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \quad \left. \begin{array}{l} \\ \\ \end{array} \right\} \quad \mapsto \quad \begin{array}{l} \text{64 ordered pairs of} \\ \text{length } 1 \cdot 1 \cdot 2 = 2 \end{array}$$

Let's use the multi-dimensional construction process

We can construct ordered quaternary Golay sequence pairs using “seed” Golay pairs of length 1, 3 and 5 as follows:

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \\ \end{array} \right\} \mapsto \begin{array}{l} 64 \text{ ordered pairs of} \\ \text{length } 1 \cdot 1 \cdot 2 = 2 \end{array}$$

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \\ \\ \end{array} \right\} \mapsto \begin{array}{l} 512 \text{ ordered pairs of} \\ \text{length } 1 \cdot 1 \cdot 1 \cdot 2^2 = 4 \end{array}$$

Ordered quaternary Golay sequence pair counts

In 2002, Craigen, Holzmann, and Kharaghani exhaustively found all ordered quaternary Golay sequence pairs of small length n :

n	# pairs	n	# pairs	n	# pairs
1	16	8	6656	15	0
2	64	9	0	16	106496
3	128	10	12288	17	0
4	512	11	512	18	24576
5	512	12	36864	19	0
6	2048	13	512	20	215040*
7	0	14	0	21	0

* Frank Fiedler, personal communication.

Let's use the multi-dimensional construction process

We can construct ordered quaternary Golay sequence pairs using “seed” Golay pairs of length 1, 3 and 5 as follows:

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \\ \end{array} \right\} \mapsto 64 \text{ ordered pairs of} \\ \left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \text{ length } 1 \cdot 1 \cdot 2 = 2$$

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \\ \\ \end{array} \right\} \mapsto 512 \text{ ordered pairs of} \\ \left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \text{ length } 1 \cdot 1 \cdot 1 \cdot 2^2 = 4 \\ \left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right)$$

Let's use the multi-dimensional construction process

We can construct ordered quaternary Golay sequence pairs using “seed” Golay pairs of length 1, 3 and 5 as follows:

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array}, \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \\ \end{array} \right\} \mapsto 64 \text{ ordered pairs of} \\ \left(\begin{array}{|c|} \hline 0 \\ \hline \end{array}, \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \text{ length } 1 \cdot 1 \cdot 2 = 2$$

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array}, \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \\ \\ \end{array} \right\} \mapsto 512 \text{ ordered pairs of} \\ \left(\begin{array}{|c|} \hline 0 \\ \hline \end{array}, \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \text{ length } 1 \cdot 1 \cdot 1 \cdot 2^2 = 4$$

$$\left(\begin{array}{|c|c|c|} \hline 2 & 0 & 0 \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 0 & 1 & 0 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \\ \end{array} \right\} \mapsto 2048 \text{ ordered pairs of} \\ \left(\begin{array}{|c|} \hline 0 \\ \hline \end{array}, \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \text{ length } 3 \cdot 1 \cdot 2 = 6$$

Let's use the multi-dimensional construction process

We can construct ordered quaternary Golay sequence pairs using “seed” Golay pairs of length 1, 3 and 5 as follows:

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \vphantom{\begin{array}{|c|} \hline 0 \\ \hline \end{array}} \right\} \mapsto 64 \text{ ordered pairs of length } 1 \cdot 1 \cdot 2 = 2$$

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \vphantom{\begin{array}{|c|} \hline 0 \\ \hline \end{array}} \right\} \mapsto 512 \text{ ordered pairs of length } 1 \cdot 1 \cdot 1 \cdot 2^2 = 4$$

$$\left(\begin{array}{|c|c|c|} \hline 2 & 0 & 0 \\ \hline \end{array} , \begin{array}{|c|c|c|} \hline 0 & 1 & 0 \\ \hline \end{array} \right) \left. \vphantom{\begin{array}{|c|c|c|} \hline 2 & 0 & 0 \\ \hline \end{array}} \right\} \mapsto 2048 \text{ ordered pairs of length } 3 \cdot 1 \cdot 2 = 6$$
$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right)$$

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left. \vphantom{\begin{array}{|c|} \hline 0 \\ \hline \end{array}} \right\} \mapsto 6144 \text{ ordered pairs of length } 1 \cdot 1 \cdot 1 \cdot 1 \cdot 2^3 = 8$$
$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right)$$

Let's use the multi-dimensional construction process

Continue as follows:

Let's use the multi-dimensional construction process

Continue as follows:

$$\left(\begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array} , \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array} \right) \left\{ \begin{array}{l} (\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array}) \end{array} \right\} \mapsto 8192 \text{ pairs of length 10}$$

$$\left(\begin{array}{|c|c|c|} \hline 2 & 0 & 0 \\ \hline \end{array} , \begin{array}{|c|c|c|} \hline 0 & 1 & 0 \\ \hline \end{array} \right) \left\{ \begin{array}{l} (\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array}) \\ (\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array}) \end{array} \right\} \mapsto 36864 \text{ pairs of length 12}$$

$$\left(\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \left\{ \begin{array}{l} (\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array}) \\ (\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array}) \\ (\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array}) \\ (\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array}) \\ (\begin{array}{|c|} \hline 0 \\ \hline \end{array} , \begin{array}{|c|} \hline 0 \\ \hline \end{array}) \end{array} \right\} \mapsto 98304 \text{ pairs of length 16}$$

Let's use the multi-dimensional construction process

Continue as follows:

$$\left(\begin{array}{|c|c|c|} \hline 2 & 0 & 0 \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 0 & 1 & 0 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \left(\begin{array}{|c|c|c|} \hline 2 & 0 & 0 \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 0 & 1 & 0 \\ \hline \end{array} \right) \end{array} \right\} \mapsto 24576 \text{ pairs of length 18}$$

$$\left(\begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array} \right) \left. \begin{array}{l} \\ \left(\begin{array}{|c|} \hline 0 \\ \hline \end{array}, \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \\ \left(\begin{array}{|c|} \hline 0 \\ \hline \end{array}, \begin{array}{|c|} \hline 0 \\ \hline \end{array} \right) \end{array} \right\} \mapsto 147456 \text{ pairs of length 20}$$

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	512	15	0	0
2	64	0	9	0	0	16	106496	8192
3	128	128	10	12288	4096	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	67584
7	0	0	14	0	0	21	0	0

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	512	15	0	0
2	64	0	9	0	0	16	106496	8192
3	128	128	10	12288	4096	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	67584
7	0	0	14	0	0	21	0	0

- “Shared autocorrelation property”

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	512	15	0	0
2	64	0	9	0	0	16	106496	8192
3	128	128	10	12288	4096	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	67584
7	0	0	14	0	0	21	0	0

- “Shared autocorrelation property”
- Multi-dimensional construction process with special length 8 pairs and a trivial length 1 pair

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	4096	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	67584
7	0	0	14	0	0	21	0	0

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	4096	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	67584
7	0	0	14	0	0	21	0	0

- Symmetry Lemma: (A, B) are a Golay pair $\Leftrightarrow (A + B, A - B)$ are a Golay pair (where A and B are in “multiplicative” notation).

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	4096	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	67584
7	0	0	14	0	0	21	0	0

- Symmetry Lemma: (A, B) are a Golay pair $\Leftrightarrow (A + B, A - B)$ are a Golay pair (where A and B are in “multiplicative” notation).
- Explains all remaining length 10 pairs and 2048 of the remaining length 20 pairs from binary Golay seed pairs.

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	0	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	65536
7	0	0	14	0	0	21	0	0

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	0	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	65536
7	0	0	14	0	0	21	0	0

- Multi-dimensional construction process with special length 10 pairs and a trivial length 1 pair

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	0	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	0
7	0	0	14	0	0	21	0	0

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	0	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	0
7	0	0	14	0	0	21	0	0

Let's look at lengths 5 and 13...

Lengths 5 and 13

$$G_{5,1} = \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array}$$

$$G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

quaternary Golay pair of length 5

$$G_{13,1} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 0 & 0 & 1 & 2 & 0 & 0 & 3 & 0 & 2 & 0 & 3 & 1 \\ \hline \end{array}$$

$$G_{13,2} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 1 & 2 & 2 & 2 & 1 & 2 & 0 & 0 & 3 & 2 & 0 & 3 \\ \hline \end{array}$$

quaternary Golay pair of length 13

Lengths 5 and 13

$$G_{5,1} = \begin{bmatrix} 0 & 0 & 0 & 3 & 1 \end{bmatrix}$$

$$G_{5,2} = \begin{bmatrix} 0 & 1 & 2 & 0 & 3 \end{bmatrix}$$

quaternary Golay pair of length 5

$$G_{13,1} = \begin{bmatrix} 0 & 0 & 0 & 1 & 2 & 0 & 0 & 3 & 0 & 2 & 0 & 3 & 1 \end{bmatrix}$$

$$G_{13,2} = \begin{bmatrix} 0 & 1 & 2 & 2 & 2 & 1 & 2 & 0 & 0 & 3 & 2 & 0 & 3 \end{bmatrix}$$

quaternary Golay pair of length 13

$$B_5 = \begin{bmatrix} 0 & 0 & 0 & 2 & 0 \end{bmatrix}$$

binary Barker sequence of length 5

$$B_{13} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 2 & 2 & 0 & 0 & 2 & 0 & 2 & 0 \end{bmatrix}$$

binary Barker sequence of length 13

Lengths 5 and 13 (cont.)

$$B_5 + G_{5,1} = \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 1 & 1 \\ \hline \end{array}$$

$$B_5 + G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 2 & 3 \\ \hline \end{array}$$

Lengths 5 and 13 (cont.)

$$B_5 + G_{5,1} =$$

0	0	0	1	1
---	---	---	---	---

$$B_5 + G_{5,2} =$$

0	1	2	2	3
---	---	---	---	---

$$B_{13} + G_{13,1} =$$

0	0	0	1	2	2	2	3	0	0	0	1	1
---	---	---	---	---	---	---	---	---	---	---	---	---

$$B_{13} + G_{13,2} =$$

0	1	2	2	2	3	0	0	0	1	2	2	3
---	---	---	---	---	---	---	---	---	---	---	---	---

Lengths 5 and 13 (cont.)

$$B_5 + G_{5,1} =$$

0	0	0	1	1
---	---	---	---	---

$$B_5 + G_{5,2} =$$

0	1	2	2	3
---	---	---	---	---

$$B_{13} + G_{13,1} =$$

0	0	0	1	2	2	2	3	0	0	0	1	1
---	---	---	---	---	---	---	---	---	---	---	---	---

$$B_{13} + G_{13,2} =$$

0	1	2	2	2	3	0	0	0	1	2	2	3
---	---	---	---	---	---	---	---	---	---	---	---	---

Lengths 5 and 13 (cont.)

$$B_5 + G_{5,1} = \begin{array}{|c|c|c|c|c|}\hline 0 & 0 & 0 & 1 & 1 \\ \hline\end{array}$$

$$B_5 + G_{5,2} = \begin{array}{|c|c|c|c|c|}\hline 0 & 1 & 2 & 2 & 3 \\ \hline\end{array}$$

$$B_{13} + G_{13,1} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|}\hline 0 & 0 & 0 & 1 & 2 & 2 & 2 & 3 & 0 & 0 & 0 & 1 & 1 \\ \hline\end{array}$$

$$B_{13} + G_{13,2} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|}\hline 0 & 1 & 2 & 2 & 2 & 3 & 0 & 0 & 0 & 1 & 2 & 2 & 3 \\ \hline\end{array}$$

$$X_m := ((0 \ 1 \ 2 \ 3)^m \ 0 \ 1)$$

$$Z_m := ((1 \ 2 \ 3 \ 0)^m \ 1 \ 2)$$

Lengths 5 and 13 (cont.)

$$B_5 + G_{5,1} = \begin{array}{|c|c|c|c|c|}\hline 0 & 0 & 0 & 1 & 1 \\ \hline\end{array}$$

$$B_5 + G_{5,2} = \begin{array}{|c|c|c|c|c|}\hline 0 & 1 & 2 & 2 & 3 \\ \hline\end{array}$$

$$B_{13} + G_{13,1} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|}\hline 0 & 0 & 0 & 1 & 2 & 2 & 2 & 3 & 0 & 0 & 0 & 1 & 1 \\ \hline\end{array}$$

$$B_{13} + G_{13,2} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|}\hline 0 & 1 & 2 & 2 & 2 & 3 & 0 & 0 & 0 & 1 & 2 & 2 & 3 \\ \hline\end{array}$$

$$X_m := ((0\ 1\ 2\ 3)^m\ 0\ 1)$$

$$Z_m := ((1\ 2\ 3\ 0)^m\ 1\ 2)$$

$$W_m := ((0\ 0\ 2\ 2)^m\ 0\ 0\ 1)$$

$$Y_m := ((0\ 2\ 2\ 0)^m\ 0\ 2\ 3)$$

Lengths 5 and 13 (cont.)

$$B_5 + G_{5,1} = \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 1 & 1 \\ \hline \end{array} = \text{int}(W_0, X_0)$$

$$B_5 + G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 2 & 3 \\ \hline \end{array} = \text{int}(Y_0, Z_0)$$

$$B_{13} + G_{13,1} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 0 & 0 & 1 & 2 & 2 & 2 & 3 & 0 & 0 & 0 & 1 & 1 \\ \hline \end{array} = \text{int}(W_1, X_1)$$

$$B_{13} + G_{13,2} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 1 & 2 & 2 & 2 & 3 & 0 & 0 & 0 & 1 & 2 & 2 & 3 \\ \hline \end{array} = \text{int}(Y_1, Z_1)$$

$$X_m := ((0\ 1\ 2\ 3)^m\ 0\ 1)$$

$$Z_m := ((1\ 2\ 3\ 0)^m\ 1\ 2)$$

$$W_m := ((0\ 0\ 2\ 2)^m\ 0\ 0\ 1)$$

$$Y_m := ((0\ 2\ 2\ 0)^m\ 0\ 2\ 3)$$

Binary Barker to quaternary Golay

Theorem

Let $m \in \mathbf{N}$. Suppose $\text{int}(A, B)$ is a binary Barker sequence of length $8m + 5$ where $A = ((0\ 0\ 0\ 2)^m\ 0\ 0\ 0)$. Then the sequences

$$E := \text{int}(A + W_m, B + X_m),$$

$$F := \text{int}(A + Y_m, B + Z_m)$$

form a quaternary Golay pair of length $8m + 5$.

Binary Barker to quaternary Golay

Theorem

Let $m \in \mathbf{N}$. Suppose $\text{int}(A, B)$ is a binary Barker sequence of length $8m + 5$ where $A = ((0\ 0\ 0\ 2)^m\ 0\ 0\ 0)$. Then the sequences

$$E := \text{int}(A + W_m, B + X_m),$$

$$F := \text{int}(A + Y_m, B + Z_m)$$

form a quaternary Golay pair of length $8m + 5$.

- Explains all length 5 and 13 quaternary Golay pairs!

Binary Barker to quaternary Golay

Theorem

Let $m \in \mathbf{N}$. Suppose $\text{int}(A, B)$ is a binary Barker sequence of length $8m + 5$ where $A = ((0 \ 0 \ 0 \ 2)^m \ 0 \ 0 \ 0)$. Then the sequences

$$E := \text{int}(A + W_m, B + X_m),$$

$$F := \text{int}(A + Y_m, B + Z_m)$$

form a quaternary Golay pair of length $8m + 5$.

- Explains all length 5 and 13 quaternary Golay pairs!
- Unfortunately, this result does not give rise to any new quaternary Golay pairs.

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	0	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	512	12	36864	0	19	0	0
6	2048	0	13	512	512	20	215040	0
7	0	0	14	0	0	21	0	0

- binary Barker to quaternary Golay theorem

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	0	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	0	12	36864	0	19	0	0
6	2048	0	13	512	0	20	215040	0
7	0	0	14	0	0	21	0	0

Ordered quaternary Golay pairs left to explain

n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain	n	# pairs	# pairs left to explain
1	16	0	8	6656	0	15	0	0
2	64	0	9	0	0	16	106496	0
3	128	128	10	12288	0	17	0	0
4	512	0	11	512	512	18	24576	0
5	512	0	12	36864	0	19	0	0
6	2048	0	13	512	0	20	215040	0
7	0	0	14	0	0	21	0	0

• ??

Constructing a Binary Barker Sequence from a Quaternary Golay Sequence

Goal

- We have seen that
particular binary Barker of length $\equiv 5 \pmod{8}$ \Rightarrow quaternary Golay of length $\equiv 5 \pmod{8}$

Goal

- We have seen that
particular binary Barker of length $\equiv 5 \pmod{8}$ \Rightarrow quaternary Golay of length $\equiv 5 \pmod{8}$
- Our (optimistic) objective:
particular binary Barker of length $\equiv 5 \pmod{8}$ \Leftarrow quaternary Golay of length $\equiv 5 \pmod{8}$

Goal

- We have seen that
particular binary Barker of length $\equiv 5 \pmod{8}$ \Rightarrow quaternary Golay of length $\equiv 5 \pmod{8}$
- Our (optimistic) objective:
particular binary Barker of length $\equiv 5 \pmod{8}$ \Leftarrow quaternary Golay of length $\equiv 5 \pmod{8}$
- Since there are no binary Barker sequences of odd length greater than 13, this would prove that there are no more quaternary Golay sequences for these lengths.

“Good” sequences

$$G_{5,1} = \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array}$$

$$G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

“Good” sequences

$$G_{5,1} = \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array}$$

$$G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

$$G_{5,1} + G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 3 & 0 \\ \hline \end{array}$$

“Good” sequences

$$G_{5,1} = \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array}$$

$$G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

$$G_{5,1} + G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 3 & 0 \\ \hline \end{array}$$

$$G_{13,1} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 0 & 0 & 1 & 2 & 0 & 0 & 3 & 0 & 2 & 0 & 3 & 1 \\ \hline \end{array}$$

$$G_{13,2} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 1 & 2 & 2 & 2 & 1 & 2 & 0 & 0 & 3 & 2 & 0 & 3 \\ \hline \end{array}$$

“Good” sequences

$$G_{5,1} = \begin{array}{|c|c|c|c|c|} \hline 0 & 0 & 0 & 3 & 1 \\ \hline \end{array}$$

$$G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 0 & 3 \\ \hline \end{array}$$

$$G_{5,1} + G_{5,2} = \begin{array}{|c|c|c|c|c|} \hline 0 & 1 & 2 & 3 & 0 \\ \hline \end{array}$$

$$G_{13,1} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 0 & 0 & 1 & 2 & 0 & 0 & 3 & 0 & 2 & 0 & 3 & 1 \\ \hline \end{array}$$

$$G_{13,2} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 1 & 2 & 2 & 2 & 1 & 2 & 0 & 0 & 3 & 2 & 0 & 3 \\ \hline \end{array}$$

$$G_{13,1} + G_{13,2} = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline 0 & 1 & 2 & 3 & 0 & 1 & 2 & 3 & 0 & 1 & 2 & 3 & 0 \\ \hline \end{array}$$

“Good sequences” (cont.)

Lemma

If A and B are sequences of length n and $A + B = (0 \ 1 \ 2 \ 3 \ \dots)$, then

$$C_B(u) = i^{-u} \cdot \overline{C_A(u)}$$

for all integers $0 < u < n$.

“Good sequences” (cont.)

Lemma

If A and B are sequences of length n and $A + B = (0 \ 1 \ 2 \ 3 \ \dots)$, then

$$C_B(u) = i^{-u} \cdot \overline{C_A(u)}$$

for all integers $0 < u < n$.

Call a sequence A of length n **good** if

$$C_A(u) = -i^{-u} \overline{C_A(u)} \text{ for all integers } 0 < u < n.$$

“Good sequences” (cont.)

Lemma

If A and B are sequences of length n and $A + B = (0 \ 1 \ 2 \ 3 \ \dots)$, then

$$C_B(u) = i^{-u} \cdot \overline{C_A(u)}$$

for all integers $0 < u < n$.

Call a sequence A of length n **good** if

$$C_A(u) = -i^{-u} \overline{C_A(u)} \text{ for all integers } 0 < u < n.$$

- A good sequence is necessarily a Golay sequence.

“Good sequences” (cont.)

Lemma

If A and B are sequences of length n and $A + B = (0 \ 1 \ 2 \ 3 \ \dots)$, then

$$C_B(u) = i^{-u} \cdot \overline{C_A(u)}$$

for all integers $0 < u < n$.

Call a sequence A of length n **good** if

$$C_A(u) = -i^{-u} \overline{C_A(u)} \text{ for all integers } 0 < u < n.$$

- A good sequence is necessarily a Golay sequence.
- $G_{5,1}$, $G_{5,2}$, $G_{13,1}$, and $G_{13,2}$ are all good.

“Good sequences” (cont.)

Lemma

If A and B are sequences of length n and $A + B = (0 \ 1 \ 2 \ 3 \ \dots)$, then

$$C_B(u) = i^{-u} \cdot \overline{C_A(u)}$$

for all integers $0 < u < n$.

Call a sequence A of length n **good** if

$$C_A(u) = -i^{-u} \overline{C_A(u)} \text{ for all integers } 0 < u < n.$$

- A good sequence is necessarily a Golay sequence.
- $G_{5,1}$, $G_{5,2}$, $G_{13,1}$, and $G_{13,2}$ are all good.
- Output sequences of Barker-to-Golay theorem are good.

A partial Barker-to-Golay converse

Theorem

Let $A = (a_0, \dots, a_{n-1})$ be a good sequence of length $n = 8m + 5$. Assume that

- (1) $a_{2u-1} + a_{2u+1} \equiv 1 \pmod{2}$, for all $1 \leq 2u - 1 \leq \frac{n-7}{2}$, and
- (2) $a_{4u} \equiv 0 \pmod{2}$, for all $4 \leq 4u \leq \frac{n-5}{2}$.

Then there exists a binary Barker sequence of length n , and so $m \in \{0, 1\}$.

A partial Barker-to-Golay converse

Theorem

Let $A = (a_0, \dots, a_{n-1})$ be a good sequence of length $n = 8m + 5$. Assume that

- (1) $a_{2u-1} + a_{2u+1} \equiv 1 \pmod{2}$, for all $1 \leq 2u - 1 \leq \frac{n-7}{2}$, and
- (2) $a_{4u} \equiv 0 \pmod{2}$, for all $4 \leq 4u \leq \frac{n-5}{2}$.

Then there exists a binary Barker sequence of length n , and so $m \in \{0, 1\}$.

Proof: About 20 pages of lemmas. \square

Summary and Open Problems

Summary and open problems

Summary of results:

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.
- Found a general construction of a quaternary Golay pair from a particular type of binary Barker sequence.

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.
- Found a general construction of a quaternary Golay pair from a particular type of binary Barker sequence.
- Established a partial converse to the Barker-to-Golay construction.

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.
- Found a general construction of a quaternary Golay pair from a particular type of binary Barker sequence.
- Established a partial converse to the Barker-to-Golay construction.

Open problems:

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.
- Found a general construction of a quaternary Golay pair from a particular type of binary Barker sequence.
- Established a partial converse to the Barker-to-Golay construction.

Open problems:

- Explain the existence of the length 3 and 11 quaternary Golay pairs.

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.
- Found a general construction of a quaternary Golay pair from a particular type of binary Barker sequence.
- Established a partial converse to the Barker-to-Golay construction.

Open problems:

- Explain the existence of the length 3 and 11 quaternary Golay pairs.
- Are there any quaternary Golay sequences of odd length greater than 13?

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.
- Found a general construction of a quaternary Golay pair from a particular type of binary Barker sequence.
- Established a partial converse to the Barker-to-Golay construction.

Open problems:

- Explain the existence of the length 3 and 11 quaternary Golay pairs.
- Are there any quaternary Golay sequences of odd length greater than 13?
 - Odd length \Rightarrow good?

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.
- Found a general construction of a quaternary Golay pair from a particular type of binary Barker sequence.
- Established a partial converse to the Barker-to-Golay construction.

Open problems:

- Explain the existence of the length 3 and 11 quaternary Golay pairs.
- Are there any quaternary Golay sequences of odd length greater than 13?
 - Odd length \Rightarrow good?
 - Can we overcome conditions (1) and (2) of Barker-to-Golay converse?

Summary and open problems

Summary of results:

- Classified and explained (almost) all ordered quaternary Golay pairs of length less than 22.
- Found a general construction of a quaternary Golay pair from a particular type of binary Barker sequence.
- Established a partial converse to the Barker-to-Golay construction.

Open problems:

- Explain the existence of the length 3 and 11 quaternary Golay pairs.
- Are there any quaternary Golay sequences of odd length greater than 13?
 - Odd length \Rightarrow good?
 - Can we overcome conditions (1) and (2) of Barker-to-Golay converse?
 - Alternative approach?

Thanks for listening!