

# Locally Equivalent Weights for Bayesian MrP

---

Ryan Giordano, Alice Cima, Erin Hartman, Jared Murray, Avi Feller

**UT Austin Statistics Seminar**

**September 2025**




# The basic problem


We have a survey population, for whom we observe:

- Covariates  $\mathbf{x}$  (e.g. race, gender, zip code, age, education level)
- Responses  $y$  (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe  $(\mathbf{x}_i, y_i)$  for  $i = 1, \dots, N_S$




Observe  $\mathbf{x}_j$  for  $j = 1, \dots, N_T$

# The basic problem


We have a survey population, for whom we observe:

- Covariates  $\mathbf{x}$  (e.g. race, gender, zip code, age, education level)
- Responses  $y$  (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe  $(\mathbf{x}_i, y_i)$  for  $i = 1, \dots, N_S$



Observe  $\mathbf{x}_j$  for  $j = 1, \dots, N_T$


**The problem is that the populations may be very different.**

# The basic problem


We have a survey population, for whom we observe:

- Covariates  $\mathbf{x}$  (e.g. race, gender, zip code, age, education level)
- Responses  $y$  (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe  $(\mathbf{x}_i, y_i)$  for  $i = 1, \dots, N_S$



Observe  $\mathbf{x}_j$  for  $j = 1, \dots, N_T$

**The problem is that the populations may be very different.**

Our survey results may be biased.

How can we use the covariates to say something about the target responses?

## Two approaches

We want  $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$ , but don't observe target population  $y_j$ .

- Assume  $p(y|\mathbf{x})$  is the same in both populations,
- But the distribution of  $\mathbf{x}$  may be different in the survey and target.

## Two approaches

We want  $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$ , but don't observe target population  $y_j$ .

- Assume  $p(y|\mathbf{x})$  is the same in both populations,
- But the distribution of  $\mathbf{x}$  may be different in the survey and target.

### Calibration weighting (CW)

- Choose “calibration weights”  $w_i$   
using only the regressors  $\mathbf{x}$   
(e.g. raking weights)

### Bayesian hierarchical modeling (MrP)

- Choose  $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$ ,  
choose prior  $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$   
(e.g. Hierarchical logistic regression)

## Two approaches

We want  $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$ , but don't observe target population  $y_j$ .

- Assume  $p(y|\mathbf{x})$  is the same in both populations,
- But the distribution of  $\mathbf{x}$  may be different in the survey and target.

### Calibration weighting (CW)

- Choose “calibration weights”  $w_i$   
using only the regressors  $\mathbf{x}$   
(e.g. raking weights)
- Take  $\hat{\mu}_{\text{CW}} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$

### Bayesian hierarchical modeling (MrP)

- Choose  $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$ ,  
choose prior  $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$   
(e.g. Hierarchical logistic regression)
- Take  $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$  and  
 $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$

## Two approaches

We want  $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$ , but don't observe target population  $y_j$ .

- Assume  $p(y|\mathbf{x})$  is the same in both populations,
- But the distribution of  $\mathbf{x}$  may be different in the survey and target.

### Calibration weighting (CW)

- Choose “calibration weights”  $w_i$  using only the regressors  $\mathbf{x}$  (e.g. raking weights)
- Take  $\hat{\mu}_{\text{CW}} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$
- Dependence on  $y_i$  is clear

### Bayesian hierarchical modeling (MrP)

- Choose  $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$ , choose prior  $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$  (e.g. Hierarchical logistic regression)
- Take  $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$  and  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$
- Dependence on  $y_i$  very complicated (Typically via MCMC draws from  $\mathcal{P}(\theta|\text{Survey data})$ )



# Two approaches

We want  $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$ , but don't observe target population  $y_j$ .

- Assume  $p(y|\mathbf{x})$  is the same in both populations,
- But the distribution of  $\mathbf{x}$  may be different in the survey and target.

## Calibration weighting (CW)

- ▶ Choose “calibration weights”  $w_i$  using only the regressors  $\mathbf{x}$  (e.g. raking weights)

- ▶ Take  $\hat{\mu}_{CW} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$

- ▶ Dependence on  $y_i$  is clear

- ▶ Weights give interpretable diagnostics:

- Frequentist variability
- Partial pooling
- Regressor balance

## Bayesian hierarchical modeling (MrP)

- ▶ Choose  $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$ , choose prior  $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$  (e.g. Hierarchical logistic regression)

- ▶ Take  $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$  and  $\hat{\mu}_{MrP} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$

- ▶ Dependence on  $y_i$  very complicated (Typically via MCMC draws from  $\mathcal{P}(\theta|\text{Survey data})$ )

- ▶ **Black box**

# Two approaches

We want  $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$ , but don't observe target population  $y_j$ .

- Assume  $p(y|\mathbf{x})$  is the same in both populations,
- But the distribution of  $\mathbf{x}$  may be different in the survey and target.

## Calibration weighting (CW)

- ▶ Choose “calibration weights”  $w_i$  using only the regressors  $\mathbf{x}$  (e.g. raking weights)

- ▶ Take  $\hat{\mu}_{CW} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$

- ▶ Dependence on  $y_i$  is clear

- ▶ Weights give interpretable diagnostics:

- Frequentist variability
- Partial pooling
- Regressor balance

## Bayesian hierarchical modeling (MrP)

- ▶ Choose  $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$ , choose prior  $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$  (e.g. Hierarchical logistic regression)

- ▶ Take  $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$  and  $\hat{\mu}_{MrP} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$

- ▶ Dependence on  $y_i$  very complicated (Typically via MCMC draws from  $\mathcal{P}(\theta|\text{Survey data})$ )

## ▶ Black box

- ← We open this box, providing analogues of all these diagnostics

## Prior work: Equivalent weights for linear models

Gelman (2007b) observes that MrP is a CW estimator when one uses linear regression to form  $\hat{y}$ :

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j = \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\mathbf{x}_j^\top \hat{\theta}}_{\text{Linear in } y_i}$$

Most existing literature on comparing CW and MrP focus on such linear models.<sup>1</sup>

---

<sup>1</sup>For example, Gelman (2007b), B., F., and H. (2021), and Chattopadhyay and Zubizarreta (2023).

## Prior work: Equivalent weights for linear models

Gelman (2007b) observes that MrP is a CW estimator when one uses linear regression to form  $\hat{y}$ :

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j = \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\mathbf{x}_j^\top \hat{\boldsymbol{\theta}}}_{\text{Linear in } y_i}$$

Most existing literature on comparing CW and MrP focus on such linear models.<sup>1</sup>

But what if you use a non-linear link function? Or a hierarchical model?

*“It would also be desirable to use nonlinear methods ... but then it would seem difficult to construct even approximately equivalent weights. Weighting and fully nonlinear models would seem to be completely incompatible methods.” — (Gelman 2007a)*

---

<sup>1</sup>For example, Gelman (2007b), B., F., and H. (2021), and Chattopadhyay and Zubizarreta (2023).

## Equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

The map from  $Y_S = y_1, \dots, y_{N_S} \mapsto m(\mathbf{x}_j^\top \hat{\theta})$  is *inherently nonlinear*.

But *some sample averages* of  $m(\mathbf{x}_j^\top \hat{\theta})$  can be approximately linear.

## Equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

The map from  $Y_S = y_1, \dots, y_{N_S} \mapsto m(\mathbf{x}_j^\top \hat{\theta})$  is *inherently nonlinear*.

But *some sample averages* of  $m(\mathbf{x}_j^\top \hat{\theta})$  can be approximately linear.

### Example #1

Additionally suppose  $\mathbf{x} \in \mathcal{X}$  is discrete and saturated. **Then MrP is a CW estimator.**

## Equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

The map from  $Y_S = y_1, \dots, y_{N_S} \mapsto m(\mathbf{x}_j^\top \hat{\theta})$  is *inherently nonlinear*.

But *some sample averages* of  $m(\mathbf{x}_j^\top \hat{\theta})$  can be approximately linear.

### Example #1

Additionally suppose  $\mathbf{x} \in \mathcal{X}$  is discrete and saturated. **Then MrP is a CW estimator.**

- Let  $\bar{y}_S^c$  denote the survey average among  $\mathbf{x} = c$  for  $c \in \mathcal{X}$

## Equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

The map from  $Y_S = y_1, \dots, y_{N_S} \mapsto m(\mathbf{x}_j^\top \hat{\theta})$  is *inherently nonlinear*.

But *some sample averages* of  $m(\mathbf{x}_j^\top \hat{\theta})$  can be approximately linear.

### Example #1

Additionally suppose  $\mathbf{x} \in \mathcal{X}$  is discrete and saturated. **Then MrP is a CW estimator.**

- Let  $\bar{y}_S^c$  denote the survey average among  $\mathbf{x} = c$  for  $c \in \mathcal{X}$
- For  $\mathbf{x} = c$ , the MLE satisfies  $m(\hat{\theta}^\top \mathbf{x}) = \bar{y}_S^c$



## Equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

The map from  $Y_S = y_1, \dots, y_{N_S} \mapsto m(\mathbf{x}_j^\top \hat{\theta})$  is *inherently nonlinear*.

But *some sample averages* of  $m(\mathbf{x}_j^\top \hat{\theta})$  can be approximately linear.

### Example #1

Additionally suppose  $\mathbf{x} \in \mathcal{X}$  is discrete and saturated. **Then MrP is a CW estimator.**

- Let  $\bar{y}_S^c$  denote the survey average among  $\mathbf{x} = c$  for  $c \in \mathcal{X}$
- For  $\mathbf{x} = c$ , the MLE satisfies  $m(\hat{\theta}^\top \mathbf{x}) = \bar{y}_S^c$
- Let  $N_S^c$  (or  $N_S^c$ ) denote the # of survey (or target) observations with  $\mathbf{x}_n = c$ .

## Equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

The map from  $Y_S = y_1, \dots, y_{N_S} \mapsto m(\mathbf{x}_j^\top \hat{\theta})$  is *inherently nonlinear*.

But *some sample averages* of  $m(\mathbf{x}_j^\top \hat{\theta})$  can be approximately linear.

### Example #1

Additionally suppose  $\mathbf{x} \in \mathcal{X}$  is discrete and saturated. **Then MrP is a CW estimator.**

- Let  $\bar{y}_S^c$  denote the survey average among  $\mathbf{x} = c$  for  $c \in \mathcal{X}$
- For  $\mathbf{x} = c$ , the MLE satisfies  $m(\hat{\theta}^\top \mathbf{x}) = \bar{y}_S^c$
- Let  $N_S^c$  (or  $N_S^c$ ) denote the # of survey (or target) observations with  $\mathbf{x}_n = c$ .

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_T} \sum_{c \in \mathcal{X}} \underbrace{N_T^c \bar{y}_S^c}_{\text{Linear in } y_i} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i^{\text{MrP}} y_i$$

$$\text{For } w_i^{\text{MrP}} = \frac{N_T^c / N_T}{N_S^c / N_S} \text{ when } \mathbf{x}_i = c.$$

## Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . **Then MrP is a *approximately* a CW estimator.**

## Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . **Then MrP is a *approximately* a CW estimator.**

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$$

## Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . **Then MrP is a *approximately* a CW estimator.**

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} \quad (\text{Law of large numbers})\end{aligned}$$

## Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . **Then MrP is a *approximately* a CW estimator.**

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)}\end{aligned}$$

## Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . **Then MrP is a *approximately* a CW estimator.**

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)}\end{aligned}$$

## Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . Then MrP is a *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)} \\ &\approx \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i m(\mathbf{x}_i^\top \hat{\theta}) && \text{(Law of large numbers)}\end{aligned}$$



## Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . Then MrP is *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)} \\ &\approx \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i m(\mathbf{x}_i^\top \hat{\theta}) && \text{(Law of large numbers)} \\ &= \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i y_i && \text{(Property of exponential family MLEs)}\end{aligned}$$

## Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . **Then MrP is approximately a CW estimator.**

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \underbrace{w_i^{\text{MrP}}}_{\alpha^\top \mathbf{x}_i} y_i + \text{Small error}$$

But what are the weights? We don't observe  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$ , so can't estimate  $\alpha$  directly.

---

<sup>2</sup>Krantz and Parks 2012; G., Stephenson, et al. 2019.

## Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . **Then MrP is approximately a CW estimator.**

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \underbrace{w_i^{\text{MrP}}}_{\alpha^\top \mathbf{x}_i} y_i + \text{Small error}$$

But what are the weights? We don't observe  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$ , so can't estimate  $\alpha$  directly.

### Key idea (informal)

If  $\hat{\mu}_{\text{MrP}}$  is approximately linear, then  $w_i^{\text{MrP}} \approx \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$ .

---

<sup>2</sup>Krantz and Parks 2012; G., Stephenson, et al. 2019.

## Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ , with MLE  $\hat{\theta}$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$ .

### Example #2

Suppose  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$  for some  $\alpha$ . Then MrP is *approximately* a CW estimator.

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \underbrace{w_i^{\text{MrP}}}_{\alpha^\top \mathbf{x}_i} y_i + \text{Small error}$$

But what are the weights? We don't observe  $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$ , so can't estimate  $\alpha$  directly.

### Key idea (informal)

If  $\hat{\mu}_{\text{MrP}}$  is approximately linear, then  $w_i^{\text{MrP}} \approx \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$ .

For logistic regression, could compute and analyze  $\frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$  using the implicit function theorem.<sup>2</sup>

---

<sup>2</sup>Krantz and Parks 2012; G., Stephenson, et al. 2019.

## Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ .
- Set a hierarchical prior  $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$ , use MCMC to draw from  $\mathcal{P}(\theta|\text{Survey data})$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[ m(\mathbf{x}_j^\top \theta) \right]$ .

No reason to think  $Y_{\mathcal{S}} \mapsto \hat{\mu}_{\text{MrP}}(Y_{\mathcal{S}})$  is even approximately linear.

Butg we can still compute and analyze  $w_i^{\text{MrP}} := \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$  using Bayesian sensitivity analysis!<sup>3</sup>

---

<sup>3</sup>Gustafson 1996; G., Broderick, and Jordan 2018.

# Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ .
- Set a hierarchical prior  $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$ , use MCMC to draw from  $\mathcal{P}(\theta|\text{Survey data})$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[ m(\mathbf{x}_j^\top \theta) \right]$ .

No reason to think  $Y_S \mapsto \hat{\mu}_{\text{MrP}}(Y_S)$  is even approximately linear.

Butg we can still compute and analyze  $w_i^{\text{MrP}} := \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$  using Bayesian sensitivity analysis!<sup>3</sup>

## MrP locally equivalent weights (MrPlew)

For new data  $\tilde{Y}_S$ , form a series expansion

$$\hat{\mu}_{\text{MrP}}(\tilde{Y}_S) \approx \hat{\mu}_{\text{MrP}}(Y_S) + \sum_{i=1}^{N_S} w_i^{\text{MrP}} (\tilde{y}_i - y_i) \quad \text{where} \quad w_i^{\text{MrP}} := \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}.$$

---

<sup>3</sup>Gustafson 1996; G., Broderick, and Jordan 2018.

# Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is  $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$ .
- Set a hierarchical prior  $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$ , use MCMC to draw from  $\mathcal{P}(\theta|\text{Survey data})$ .
- MrP is  $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[ m(\mathbf{x}_j^\top \theta) \right]$ .

No reason to think  $Y_S \mapsto \hat{\mu}_{\text{MrP}}(Y_S)$  is even approximately linear.

Butg we can still compute and analyze  $w_i^{\text{MrP}} := \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$  using Bayesian sensitivity analysis!<sup>3</sup>

## MrP locally equivalent weights (MrPlew)

For new data  $\tilde{Y}_S$ , form a series expansion

$$\hat{\mu}_{\text{MrP}}(\tilde{Y}_S) \approx \hat{\mu}_{\text{MrP}}(Y_S) + \sum_{i=1}^{N_S} w_i^{\text{MrP}} (\tilde{y}_i - y_i) \quad \text{where} \quad w_i^{\text{MrP}} := \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}.$$


**Our task is to rigorously show that even such local weights can be used diagnostically.**

---

<sup>3</sup>Gustafson 1996; G., Broderick, and Jordan 2018.

# The weights can look very different!

Does this mean anything? Are the differences important?



**Figure 1:** Comparison between raking and MrPlew weights for a particular example



Note that there was no talk of correct specification for the data you have.

That was a foregone conclusion when we started looking at equivalent weights!

How do you perform model checking with sensitivity analysis?

Existing methods evaluate whether the analysis changes “a lot” when you:

- Parametrically perturb the model (e.g. fit a richer model class)
- Non-parametrically perturb the data (e.g. produce gross outliers)

The problem is:

- How much is “a lot”?
- Non-parametric data perturbations are hard to reason about
- It’s hard to say whether parametric model changes are enough

Instead, we

- Parametrically perturb the data
- Observe whether our model could detect the change
- Know exactly the expected change (don’t have to decide on what “a lot” means)
- Easy to reason about whether the data perturbation is reasonable
- Don’t need to propose an alternative model, instead study the model you have

### Student contributions and future work:

- **Alice Cima** contributed significantly to this work
- **Vladimir Palmin** is working on extending MrPlew to lme4
- **Sequoia Andrade** is working on generalizing to other local sensitivity checks
- **Lucas Schwengber** is working on novel flow-based techniques for local sensitivity




Alice Cima

No picture!  
Vladimir Palmin











Sequoia Andrade



Lucas Schwengber

# References

- 
- B., Eli, Avi F., and Erin H. (2021).
- Multilevel calibration weighting for survey data*
- . arXiv: 2102.09052 [stat.ME].
- 
- 
- Chattopadhyay, A. and J. Zubizarreta (2023). “On the implied weights of linear regression for causal inference”. In:
- Biometrika*
- 110.3, pp. 615–629.
- 
- 
- G., T. Broderick, and M. I. Jordan (2018). “Covariances, robustness and variational bayes”. In:
- Journal of machine learning research*
- 19.51.
- 
- 
- G., W. Stephenson, et al. (2019). “A swiss army infinitesimal jackknife”. In:
- The 22nd International Conference on Artificial Intelligence and Statistics*
- . PMLR, pp. 1139–1147.
- 
- 
- Gelman, A. (2007a). “Rejoinder: Struggles with survey weighting and regression modelling”. In:
- Statistical Science*
- 22.2, pp. 184–188.
- 
- 
- (2007b). “Struggles with survey weighting and regression modeling”. In.
- 
- 
- Gustafson, P. (1996). “Local sensitivity of posterior expectations”. In:
- The Annals of Statistics*
- 24.1, pp. 174–195.
- 
- 
- Krantz, S. and H. Parks (2012).
- The Implicit Function Theorem: History, Theory, and Applications*
- . Springer Science & Business Media.