

An Automatic Finite-Sample Robustness Metric: Can Dropping a Little Data Make a Big Difference?



Ryan Giordano
MIT



Rachael Meager
LSE



Tamara Broderick
MIT

Job talk 2021

Dropping data: Motivation

You're a data analyst, and you've

- Gathered some exchangeable data,
- Cleaned up / removed outliers,
- Checked for correct specification, and
- Drawn a conclusion from your statistical analysis
(e.g., based the sign / significance of some estimated parameter).

Dropping data: Motivation

You're a data analyst, and you've

- Gathered some exchangeable data,
- Cleaned up / removed outliers,
- Checked for correct specification, and
- Drawn a conclusion from your statistical analysis
(e.g., based the sign / significance of some estimated parameter).

Well done!

Would you be concerned if you could **reverse your conclusion** by removing a **small proportion** (say, 0.1%) of your data?

Dropping data: Mexico Microcredit

Consider β , a randomized controlled trial study of the efficacy of microcredit in Mexico based on 16,560 data points.

The variable “Beta” estimates the effect of microcredit in US dollars.

| | Left out points | Beta (SE) |
|----------|-----------------|--------------|
| Original | 0 | -4.55 (5.88) |

Dropping data: Mexico Microcredit

Consider β , a randomized controlled trial study of the efficacy of microcredit in Mexico based on 16,560 data points.

The variable “Beta” estimates the effect of microcredit in US dollars.

| | Left out points | Beta (SE) |
|-------------|-----------------|--------------|
| Original | 0 | -4.55 (5.88) |
| Change sign | 1 | 0.4 (3.19) |

Dropping data: Mexico Microcredit

Consider β , a randomized controlled trial study of the efficacy of microcredit in Mexico based on 16,560 data points.

The variable “Beta” estimates the effect of microcredit in US dollars.

| | Left out points | Beta (SE) |
|---------------------|-----------------|---------------|
| Original | 0 | -4.55 (5.88) |
| Change sign | 1 | 0.4 (3.19) |
| Change significance | 14 | -10.96 (5.57) |

Dropping data: Mexico Microcredit

Consider β , a randomized controlled trial study of the efficacy of microcredit in Mexico based on 16,560 data points.

The variable “Beta” estimates the effect of microcredit in US dollars.

| | Left out points | Beta (SE) |
|---------------------|-----------------|---------------|
| Original | 0 | -4.55 (5.88) |
| Change sign | 1 | 0.4 (3.19) |
| Change significance | 14 | -10.96 (5.57) |
| Change both | 15 | 7.03 (2.55) |

Dropping data: Mexico Microcredit

Consider β , a randomized controlled trial study of the efficacy of microcredit in Mexico based on 16,560 data points.

The variable “Beta” estimates the effect of microcredit in US dollars.

| | Left out points | Beta (SE) |
|---------------------|-----------------|---------------|
| Original | 0 | -4.55 (5.88) |
| Change sign | 1 | 0.4 (3.19) |
| Change significance | 14 | -10.96 (5.57) |
| Change both | 15 | 7.03 (2.55) |

By removing very few data points ($15/16560 \approx 0.1\%$), we can reverse the qualitative conclusions of the original study!

Dropping data: Mexico Microcredit

Consider β , a randomized controlled trial study of the efficacy of microcredit in Mexico based on 16,560 data points.

The variable “Beta” estimates the effect of microcredit in US dollars.

| | Left out points | Beta (SE) |
|---------------------|-----------------|---------------|
| Original | 0 | -4.55 (5.88) |
| Change sign | 1 | 0.4 (3.19) |
| Change significance | 14 | -10.96 (5.57) |
| Change both | 15 | 7.03 (2.55) |

By removing very few data points ($15/16560 \approx 0.1\%$), we can reverse the qualitative conclusions of the original study!

Question: Is the reported interval $-4.55 \pm (5.88)$ a reasonable description of the uncertainty in the estimated efficacy of microcredit?

Dropping data: Motivation

Would you be concerned if you could **reverse your conclusion** by removing a **small proportion** (say, 0.1%) of your data?

Dropping data: Motivation

Would you be concerned if you could **reverse your conclusion** by removing a **small proportion** (say, 0.1%) of your data?

Not always!

Dropping data: Motivation

Would you be concerned if you could **reverse your conclusion** by removing a **small proportion** (say, 0.1%) of your data?

Not always!

...but sometimes, surely yes.

For example, often in economics:

- Small fractions of data are missing not-at-random,
- Policy population is different from analyzed population,
- We report a convenient summary (e.g. mean) of a complex effect,
- Models are stylized proxies of reality.

Objective

Estimate the effect of leaving out $\lfloor \alpha N \rfloor$ datapoints, where α is small.

Objective

Estimate the effect of leaving out $\lfloor \alpha N \rfloor$ datapoints, where α is small.

Question 1: How do we find influential datapoints?

The number of subsets $\binom{N}{\lfloor \alpha N \rfloor}$ can be very large even when α is very small.

In the MX microcredit study, $\binom{16560}{15} \approx 1.4 \cdot 10^{51}$ sets to check for $\alpha = 0.0009$.

We provide a fast, automatic approximation based on the **influence function**.

Objective

Estimate the effect of leaving out $\lfloor \alpha N \rfloor$ datapoints, where α is small.

Question 1: How do we find influential datapoints?

Question 2: What makes an estimator non-robust?

Objective

Estimate the effect of leaving out $\lfloor \alpha N \rfloor$ datapoints, where α is small.

Question 1: How do we find influential datapoints?

Question 2: What makes an estimator non-robust?

Non-robustness to removal of $\lfloor \alpha N \rfloor$ points is:

- Not (necessarily) caused by misspecification.
- Not (necessarily) caused by outliers.
- Not captured by standard errors.
- Not mitigated by large N .
- Primarily determined by the **signal to noise** ratio
... in a sense which we will define.

Objective

Estimate the effect of leaving out $\lfloor \alpha N \rfloor$ datapoints, where α is small.

Question 1: How do we find influential datapoints?

Question 2: What makes an estimator non-robust?

Question 3: When is our approximation accurate?

Objective

Estimate the effect of leaving out $\lfloor \alpha N \rfloor$ datapoints, where α is small.

Question 1: How do we find influential datapoints?

Question 2: What makes an estimator non-robust?

Question 3: When is our approximation accurate?

- We provide deterministic error bounds for small α .
- We show the accuracy in simple experiments.
- We show the accuracy in a number of real-world experiments.

Objective

Estimate the effect of leaving out $\lfloor \alpha N \rfloor$ datapoints, where α is small.

Question 1: How do we find influential datapoints?

Question 2: What makes an estimator non-robust?

Question 3: When is our approximation accurate?

Conclusion: Related work and future directions

Question 1:

How do we find influential datapoints?

Which estimators do we study?

We study “Z-estimators,” i.e., roots of estimating equations.

Suppose we have N data points d_1, \dots, d_N . Then:

$$\hat{\theta} := \vec{\theta} \text{ such that } \sum_{n=1}^N G(\vec{\theta}, d_n) = 0_P.$$

Examples: all minimizers of empirical loss (OLS, MLE, VB), and more.

Which estimators do we study?

We study “Z-estimators,” i.e., roots of estimating equations.

Suppose we have N data points d_1, \dots, d_N . Then:

$$\hat{\theta}(\vec{w}) := \vec{\theta} \text{ such that } \sum_{n=1}^N \vec{w}_n G(\vec{\theta}, d_n) = 0_P.$$

Examples: all minimizers of empirical loss (OLS, MLE, VB), and more.

Leave points out by setting their elements of \vec{w} to zero.

Which estimators do we study?

Suppose we have N data points d_1, \dots, d_N . Then:

$$\hat{\theta}(\vec{w}) := \vec{\theta} \text{ such that } \sum_{n=1}^N \vec{w}_n G(\vec{\theta}, d_n) = 0_P.$$

Leave points out by setting their elements of \vec{w} to zero.

Which estimators do we study?

Suppose we have N data points d_1, \dots, d_N . Then:

$$\hat{\theta}(\vec{w}) := \vec{\theta} \text{ such that } \sum_{n=1}^N \vec{w}_n G(\vec{\theta}, d_n) = 0_P.$$

Leave points out by setting their elements of \vec{w} to zero.

Fix a quantity of interest, $\phi(\vec{\theta})$:

$$\phi(\vec{\theta}) = \vec{\theta}_p$$

$$\phi(\vec{\theta}) = \vec{\theta}_p + \frac{1.96}{\sqrt{N}} \hat{\sigma}_{\phi}(\vec{\theta})$$

Which estimators do we study?

Suppose we have N data points d_1, \dots, d_N . Then:

$$\hat{\theta}(\vec{w}) := \vec{\theta} \text{ such that } \sum_{n=1}^N \vec{w}_n G(\vec{\theta}, d_n) = 0_P.$$

Leave points out by setting their elements of \vec{w} to zero.

Fix a quantity of interest, $\phi(\vec{\theta})$:

$$\phi(\vec{\theta}) = \vec{\theta}_p$$

$$\phi(\vec{\theta}) = \vec{\theta}_p + \frac{1.96}{\sqrt{N}} \hat{\sigma}_{\phi}(\vec{\theta})$$

Let the “**signal**”, Δ , be a
“large” change in ϕ .

Which estimators do we study?

Suppose we have N data points d_1, \dots, d_N . Then:

$$\hat{\theta}(\vec{w}) := \vec{\theta} \text{ such that } \sum_{n=1}^N \vec{w}_n G(\vec{\theta}, d_n) = 0_P.$$


Leave points out by setting their elements of \vec{w} to zero.

Fix a quantity of interest, $\phi(\vec{\theta})$:

$$\phi(\vec{\theta}) = \vec{\theta}_p$$

$$\phi(\vec{\theta}) = \vec{\theta}_p + \frac{1.96}{\sqrt{N}} \hat{\sigma}_{\phi}(\vec{\theta})$$

Let the “**signal**”, Δ , be a “large” change in ϕ .



Which estimators do we study?

Suppose we have N data points d_1, \dots, d_N . Then:

$$\hat{\theta}(\vec{w}) := \vec{\theta} \text{ such that } \sum_{n=1}^N \vec{w}_n G(\vec{\theta}, d_n) = 0_P.$$

Leave points out by setting their elements of \vec{w} to zero.

Fix a quantity of interest, $\phi(\vec{\theta})$.

Let the “**signal**”, Δ , be a “large” change in ϕ .

Can we reverse our conclusion by dropping $\lfloor \alpha N \rfloor$ datapoints?

Which estimators do we study?

Suppose we have N data points d_1, \dots, d_N . Then:

$$\hat{\theta}(\vec{w}) := \vec{\theta} \text{ such that } \sum_{n=1}^N \vec{w}_n G(\vec{\theta}, d_n) = 0_P.$$

Leave points out by setting their elements of \vec{w} to zero.

Fix a quantity of interest, $\phi(\vec{\theta})$.

Let the “**signal**”, Δ , be a “large” change in ϕ .

Can we reverse our conclusion by dropping $\lfloor \alpha N \rfloor$ datapoints? \Leftrightarrow

Is there a \vec{w} , with $\lfloor \alpha N \rfloor$ zeros, such that $\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \geq \Delta$?

Hard! Evaluating $\hat{\theta}(\vec{w})$ is costly and lots of \vec{w} have $\lfloor \alpha N \rfloor$ zeros.

Taylor series approximation.

Is there a \vec{w} , with $\lfloor \alpha N \rfloor$ zeros, such that $\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \geq \Delta$?

To simplify the search over \vec{w} , we form the Taylor series approximation:

$$\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \approx \phi^{\text{lin}}(\vec{w}) - \phi(\hat{\theta}) := - \sum_{n: \vec{w}_n=0} \psi_n, \text{ where } \psi_n := \left. \frac{\partial \phi(\hat{\theta}(\vec{w}))}{\partial \vec{w}_n} \right|_{\vec{1}}.$$

Taylor series approximation.

Is there a \vec{w} , with $\lfloor \alpha N \rfloor$ zeros, such that $\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \geq \Delta$?

To simplify the search over \vec{w} , we form the Taylor series approximation:

$$\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \approx \phi^{\text{lin}}(\vec{w}) - \phi(\hat{\theta}) := - \sum_{n: \vec{w}_n=0} \psi_n, \text{ where } \psi_n := \left. \frac{\partial \phi(\hat{\theta}(\vec{w}))}{\partial \vec{w}_n} \right|_{\vec{1}}.$$

The values ψ_n are the “**empirical influence function.**” (?)

The ψ_n can be **easily and automatically** computed from $\hat{\theta}$.

The approximation is **typically accurate** for small α . (?)

Taylor series approximation.

Is there a \vec{w} , with $\lfloor \alpha N \rfloor$ zeros, such that $\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \geq \Delta$?

To simplify the search over \vec{w} , we form the Taylor series approximation:

$$\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \approx \phi^{\text{lin}}(\vec{w}) - \phi(\hat{\theta}) := - \sum_{n: \vec{w}_n=0} \psi_n, \text{ where } \psi_n := \left. \frac{\partial \phi(\hat{\theta}(\vec{w}))}{\partial \vec{w}_n} \right|_{\vec{1}}.$$

The values ψ_n are the “**empirical influence function.**” (?)

The ψ_n can be **easily and automatically** computed from $\hat{\theta}$.

The approximation is **typically accurate** for small α . (?)

Is there a \vec{w} , with $\lfloor \alpha N \rfloor$ zeros, such that $\phi^{\text{lin}}(\vec{w}) - \phi(\hat{\theta}) \geq \Delta$?

Taylor series approximation.

Is there a \vec{w} , with $\lfloor \alpha N \rfloor$ zeros, such that $\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \geq \Delta$?

To simplify the search over \vec{w} , we form the Taylor series approximation:

$$\phi(\hat{\theta}(\vec{w})) - \phi(\hat{\theta}) \approx \phi^{\text{lin}}(\vec{w}) - \phi(\hat{\theta}) := - \sum_{n: \vec{w}_n=0} \psi_n, \text{ where } \psi_n := \left. \frac{\partial \phi(\hat{\theta}(\vec{w}))}{\partial \vec{w}_n} \right|_{\vec{1}}.$$

The values ψ_n are the “**empirical influence function.**” (?)

The ψ_n can be **easily and automatically** computed from $\hat{\theta}$.

The approximation is **typically accurate** for small α . (?)

Is there a \vec{w} , with $\lfloor \alpha N \rfloor$ zeros, such that $\phi^{\text{lin}}(\vec{w}) - \phi(\hat{\theta}) \geq \Delta$?

Easy! The most influential points for $\phi^{\text{lin}}(\vec{w})$ have the most negative ψ_n .

Taylor series approximation.

Procedure:

Taylor series approximation.

Procedure:

1. Compute the “original” estimator, $\hat{\theta}$ and $\phi(\hat{\theta})$.

Taylor series approximation.

Procedure:

1. Compute the “original” estimator, $\hat{\theta}$ and $\phi(\hat{\theta})$.
2. Compute and sort the influence scores, $\psi_{(1)} \leq \psi_{(2)} \leq \dots \leq \psi_{(N)}$.

Taylor series approximation.

Procedure:

1. Compute the “original” estimator, $\hat{\theta}$ and $\phi(\hat{\theta})$.
2. Compute and sort the influence scores, $\psi_{(1)} \leq \psi_{(2)} \leq \dots \leq \psi_{(N)}$.
3. Let \vec{w}^* leave out the data corresponding to $\psi_{(1)}, \dots, \psi_{(\lfloor \alpha N \rfloor)}$.

Taylor series approximation.

Procedure:

1. Compute the “original” estimator, $\hat{\theta}$ and $\phi(\hat{\theta})$.
2. Compute and sort the influence scores, $\psi_{(1)} \leq \psi_{(2)} \leq \dots \leq \psi_{(N)}$.
3. Let \vec{w}^* leave out the data corresponding to $\psi_{(1)}, \dots, \psi_{(\lfloor \alpha N \rfloor)}$.
4. Report non-robustness if $\Delta \leq \phi^{\text{lin}}(\vec{w}^*) - \phi(\hat{\theta}) = -\sum_{n=1}^{\lfloor \alpha N \rfloor} \psi_{(n)}$.

Taylor series approximation.

Procedure:

1. Compute the “original” estimator, $\hat{\theta}$ and $\phi(\hat{\theta})$.
2. Compute and sort the influence scores, $\psi_{(1)} \leq \psi_{(2)} \leq \dots \leq \psi_{(N)}$.
3. Let \vec{w}^* leave out the data corresponding to $\psi_{(1)}, \dots, \psi_{(\lfloor \alpha N \rfloor)}$.
4. Report non-robustness if $\Delta \leq \phi^{\text{lin}}(\vec{w}^*) - \phi(\hat{\theta}) = -\sum_{n=1}^{\lfloor \alpha N \rfloor} \psi_{(n)}$.
- 5 **Optional:** Compute $\hat{\theta}(\vec{w}^*)$, and verify that $\Delta \leq \phi(\hat{\theta}(\vec{w}^*)) - \phi(\hat{\theta})$.

Computing the influence function.

How to compute $\psi_n := \left. \frac{\partial \phi(\hat{\theta}(\vec{w}))}{\partial \vec{w}_n} \right|_{\vec{1}}$? Recall $\sum_{n=1}^N \vec{w}_n G(\hat{\theta}(\vec{w}), d_n) = 0_P$.

Step zero: Implement software to compute $G(\theta, d_n)$ and $\phi(\theta)$. Find $\hat{\theta}$.

Step one: By the chain rule, $\psi_n = \left. \frac{\partial \phi(\hat{\theta}(\vec{w}))}{\partial \vec{w}_n} \right|_{\vec{1}} = \left. \frac{\partial \phi(\theta)}{\partial \theta^T} \right|_{\hat{\theta}} \left. \frac{\partial \hat{\theta}(\vec{w})}{\partial \vec{w}_n} \right|_{\vec{1}}$.

Step two: By the implicit function theorem:

$$\left. \frac{\partial \hat{\theta}(\vec{w})}{\partial \vec{w}_n} \right|_{\vec{1}} = \frac{1}{N} \left(\frac{1}{N} \sum_{n'=1}^N \left. \frac{\partial}{\partial \theta^T} G(\vec{\theta}, d_{n'}) \right|_{\hat{\theta}} \right)^{-1} G(\hat{\theta}, d_n).$$

Step three: Use *automatic differentiation* on $\phi(\theta)$ and $G(\theta, d_n)$ from step zero to compute $\left. \frac{\partial \phi(\theta)}{\partial \theta^T} \right|_{\hat{\theta}}$ and $\left. \frac{\partial}{\partial \theta^T} G(\vec{\theta}, d_n) \right|_{\hat{\theta}}$.

-
- The user does step zero. The rest is automatic.
 - The primary computational expense is the Hessian inverse.
 - Automatic differentiation is the chain rule applied to a program.
 - Typically $\psi_n = O(N^{-1})$.

Question 2:

What makes an estimator non-robust?

What makes an estimator non-robust? A tail sum.

$$\Delta \leq \phi^{\text{lin}}(\vec{w}^*) - \phi(\hat{\theta})$$

Report non-robustness

$$= - \sum_{n=1}^{\lfloor \alpha N \rfloor} \psi_{(n)}$$

(By definition)

$$= - \frac{1}{N} \sum_{n=1}^{\lfloor \alpha N \rfloor} N \psi_{(n)}$$

(Recall $\psi_n = O_p(N^{-1})$)

$$\leq \underbrace{\left(\frac{1}{N} \sum_{n=1}^N N^2 \psi_{(n)}^2 \right)^{1/2}}_{=:\hat{\sigma}_\phi} \underbrace{\left(\frac{1}{N} \sum_{n=1}^N \mathbb{I}(n \leq \lfloor \alpha N \rfloor) \right)^{1/2}}_{=:\mathcal{S}_\alpha \leq \sqrt{\alpha}}$$

(Cauchy-Schwartz)

Suppose that $\hat{\theta} \xrightarrow{P} \theta_0$ and $\phi(\hat{\theta}) \rightsquigarrow \mathcal{N}(\phi(\theta_0), \sigma^2)$.

Typically, $\hat{\sigma}_\phi \xrightarrow{P} \sigma$ (?).


A slightly more careful analysis gives $\mathcal{S}_\alpha \leq \sqrt{\alpha(1-\alpha)}$.

What makes an estimator non-robust? A tail sum.

Report non-robustness if the “**signal to noise ratio**” $\frac{\Delta}{\hat{\sigma}_\phi} \leq \mathcal{S}_\alpha$ where

- The “noise” $\hat{\sigma}_\phi^2 \rightarrow \text{Var}(\sqrt{N}\phi)$ (?)
- The “shape” $\mathcal{S}_\alpha \leq \sqrt{\alpha(1-\alpha)}$ and converges to a nonzero constant

Influence score histogram (N = 10000, $\alpha = 0.05$)



Corollaries.

Corollaries.

Report non-robustness if the “**signal to noise ratio**” $\frac{\Delta}{\hat{\sigma}_\phi} \leq \mathcal{S}_\alpha$.

Corollaries.

Report non-robustness if the “**signal to noise ratio**” $\frac{\Delta}{\hat{\sigma}_\phi} \leq \mathcal{S}_\alpha$.

Corollary: Non-robustness possible even with correct specification.

Corollaries.

Report non-robustness if the “**signal to noise ratio**” $\frac{\Delta}{\hat{\sigma}_\phi} \leq S_\alpha$.

Corollary: Non-robustness possible even with correct specification.

Corollary: Leave- $\lfloor \alpha N \rfloor$ -out robustness does not vanish as $N \rightarrow \infty$.

Corollaries.

Report non-robustness if the “**signal to noise ratio**” $\frac{\Delta}{\hat{\sigma}_\phi} \leq S_\alpha$.

Corollary: Non-robustness possible even with correct specification.

Corollary: Leave- $\lfloor \alpha N \rfloor$ -out robustness does not vanish as $N \rightarrow \infty$.

Recall that standard errors reject when $\frac{\Delta}{\hat{\sigma}_\phi} \leq \frac{1.96}{\sqrt{N}}$.

Corollaries.

Report non-robustness if the “**signal to noise ratio**” $\frac{\Delta}{\hat{\sigma}_\phi} \leq S_\alpha$.

Corollary: Non-robustness possible even with correct specification.

Corollary: Leave- $\lfloor \alpha N \rfloor$ -out robustness does not vanish as $N \rightarrow \infty$.

Recall that standard errors reject when $\frac{\Delta}{\hat{\sigma}_\phi} \leq \frac{1.96}{\sqrt{N}}$.

Corollary: Leave- $\lfloor \alpha N \rfloor$ -out is different from standard errors.

Corollaries.

Report non-robustness if the “**signal to noise ratio**” $\frac{\Delta}{\hat{\sigma}_\phi} \leq S_\alpha$.

Corollary: Non-robustness possible even with correct specification.

Corollary: Leave- $\lfloor \alpha N \rfloor$ -out robustness does not vanish as $N \rightarrow \infty$.

Recall that standard errors reject when $\frac{\Delta}{\hat{\sigma}_\phi} \leq \frac{1.96}{\sqrt{N}}$.

Corollary: Leave- $\lfloor \alpha N \rfloor$ -out is different from standard errors.

Corollary: Insignificance is always non-robust.

Take $\Delta = \frac{1.96\hat{\sigma}_\phi}{\sqrt{N}} \rightarrow 0 \leq S_\alpha$.

Corollaries.

Report non-robustness if the “**signal to noise ratio**” $\frac{\Delta}{\hat{\sigma}_\phi} \leq S_\alpha$.

Corollary: Non-robustness possible even with correct specification.

Corollary: Leave- $\lfloor \alpha N \rfloor$ -out robustness does not vanish as $N \rightarrow \infty$.

Recall that standard errors reject when $\frac{\Delta}{\hat{\sigma}_\phi} \leq \frac{1.96}{\sqrt{N}}$.

Corollary: Leave- $\lfloor \alpha N \rfloor$ -out is different from standard errors.

Corollary: Insignificance is always non-robust.

Take $\Delta = \frac{1.96\hat{\sigma}_\phi}{\sqrt{N}} \rightarrow 0 \leq S_\alpha$.

Corollary: Gross outliers primarily affect robustness through $\hat{\sigma}_\phi$.

Cauchy-Schwartz is tight when all the influence scores are the same.

Question 3:

When is our approximation accurate?

The influence function

- Weights as derivatives
- Influence function
- Simulation
- Experiments

Conclusion

- You may be concerned if you could reverse your conclusion by removing a $\lfloor \alpha N \rfloor$ datapoints, for some small α .

Conclusion

- You may be concerned if you could reverse your conclusion by removing a $\lfloor \alpha N \rfloor$ datapoints, for some small α .
- Robustness to removing a $\lfloor \alpha N \rfloor$ datapoints is principally determined by the signal to noise ratio, does not disappear asymptotically, and is distinct from (and typically larger than) standard errors.

Conclusion

- You may be concerned if you could reverse your conclusion by removing a $\lfloor \alpha N \rfloor$ datapoints, for some small α .
- Robustness to removing a $\lfloor \alpha N \rfloor$ datapoints is principally determined by the signal to noise ratio, does not disappear asymptotically, and is distinct from (and typically larger than) standard errors.
- Robustness to removing a $\lfloor \alpha N \rfloor$ datapoints is easy to check! We can quickly and automatically find an approximate influential set which is accurate for small α .

Links and references

Tamara Broderick, Ryan Giordano, Rachael Meager (alphabetical authors)
“An Automatic Finite-Sample Robustness Metric: Can Dropping a Little Data Change Conclusions?”

<https://arxiv.org/abs/2011.14999>

See the paper for applications to:

- Hierarchical meta-analysis of microcredit (?)
- Cash transfers randomized controlled trial (?)
- Oregon Medicaid experiment (?)
- Expository simulations

zaminfluence: R package with leave- α -out robustness for OLS and IV estimators

<https://github.com/rgiordan/zaminfluence>

-
- M. Angelucci and G. De Giorgi. Indirect effects of an aid program: How do cash transfers affect ineligible's consumption? *American Economic Review*, 99(1):486–508, 2009.
- M. Angelucci, D. Karlan, and J. Zinman. Microcredit impacts: Evidence from a randomized microcredit program placement experiment by Compartamos Banco. *American Economic Journal: Applied Economics*, 7(1):151–82, 2015.
- A. Finkelstein, S. Taubman, B. Wright, M. Bernstein, J. Gruber, J. Newhouse, H. Allen, K. Baicker, and Oregon Health Study Group. The Oregon health insurance experiment: Evidence from the first year. *The Quarterly Journal of Economics*, 127(3):1057–1106, 2012.
- R. Giordano, M. I. Jordan, and T. Broderick. A higher-order Swiss army infinitesimal jackknife. *arXiv preprint arXiv:1907.12116*, 2019.
- F. Hampel. *Robust statistics: The approach based on influence functions*, volume 196. Wiley-Interscience, 1986.
- R. Meager. Aggregating distributional treatment effects: A Bayesian hierarchical analysis of the microcredit literature. *LSE working paper*, 2020.