

# Variational Methods for Latent Variable Problems (part 2)

---

Ryan Giordano (for Johns Hopkins Biostats BLAST working group)

Oct, 2021

Massachusetts Institute of Technology

Outline for today:


- What counts as variational inference?
- Kullback-Leibler (KL) divergence and “standard” variational inference
- The classical EM algorithm as a special case of variational inference
- Variational inference as a generalization of the EM algorithm
- A quick and incomplete sketch of further topics in variational inference

# What counts as variational inference?

Lots of very different procedures go by the name “variational inference.” I propose an (idiosyncratic) encompassing definition based on the use cases and the name:

**Variational inference is inference using optimization.**

Think “calculus of variations:” an optimum  $\hat{x} = \operatorname{argmax} \theta f(x)$  is characterized by  $df/dx|_{\hat{x}} = 0$ , i.e. where small variations in  $\hat{x}$  result in no changes to the value of  $f(\hat{x})$ .



By this definition,

- The maximum likelihood estimator (MLE) is VI.
- The Laplace approximation to a Bayesian posterior is VI.
- Markov chain Monte Carlo (MCMC) is not VI.

# What counts as variational inference?

A more common definition of VI is the following.

Suppose we have a random variable  $\xi$  and a distribution  $p(\xi)$  that we want to know.

Let  $y$  denote data and  $\theta$  a parameter. Examples:

- The variable is  $\theta$ , and we wish to know the posterior  $p(\theta|y)$  (Bayes)
- The variable is  $y$ , and we wish to know  $p(y)$  (MLE)
- The variable is  $y$ , and we wish to know the map  $\theta \mapsto p(y|\theta) = \int p(y, z|\theta) dz$  (marginal MLE)

Let  $\mathcal{Q}$  be some class of distributions which may or may not contain  $p(\xi)$ .

**Variational inference finds the distribution in  $\mathcal{Q}$  closest to  $p$  according to some measure of “divergence” between distributions:**

$$q^*(\xi) = \operatorname{argmin}_{q \in \mathcal{Q}} D(q, p).$$

The most common choice of “divergence” is the **Kullback-Leibler** (KL) divergence, though other choices are possible [Li and Turner, 2016, Liu and Wang, 2016, Ambrogioni et al., 2018].

# KL divergence

The KL divergence is defined as:

$$\text{KL}(q||p) := \mathbb{E}_{q(\xi)} [\log q(\xi)] - \mathbb{E}_{q(\xi)} [\log p(\xi)]$$

Some points to be aware of:

- $\text{KL}(q||p) \geq 0$
- $\text{KL}(q||p) = 0 \Rightarrow p = q$
- $\text{KL}(q||p) \neq \text{KL}(p||q)$
- $\text{KL}(q||p)$  is a “strict” measure of closeness [Gibbs and Su, 2002]

Why use KL divergence?

**Phony answer:** The KL divergence has an information theoretic interpretation [Kullback and Leibler, 1951].

**Real answer:** Mathematical convenience (normalizing constants pop out).

**Example: the MLE minimizes KL divergence.** Suppose that  $x_n \stackrel{iid}{\sim} p(\cdot)$ , and  $q(\cdot|\theta) \in \mathcal{Q}$  is a parameteric family of data distributions. Then


# KL divergence exercises

$$\text{KL} (q(\theta)||p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{All bivariate normals}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL} (q(\theta)||p(\theta))$ ?




# KL divergence exercises

$$\text{KL} (q(\theta)||p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{All bivariate normals}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL} (q(\theta)||p(\theta))$ ?



**Sufficiently expressive families recover the target distribution.**


## KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{Independent bivariate normals}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(q(\theta) || p(\theta))$ ?






## KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{Independent bivariate normals}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\operatorname{argmin}} \text{KL}(q(\theta) || p(\theta))$ ?



**KL minimizers “fit inside” the second argument.**


## KL divergence exercises

$$\begin{aligned} \text{KL}(p(\theta) || q(\theta)) = \\ - \mathbb{E}_{p(\theta)} [\log q(\theta)] + \mathbb{E}_{p(\theta)} [\log p(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{Independent bivariate normals}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(p(\theta) || q(\theta))$ ?




## KL divergence exercises

$$\begin{aligned} \text{KL}(p(\theta) || q(\theta)) = \\ - \mathbb{E}_{p(\theta)} [\log q(\theta)] + \mathbb{E}_{p(\theta)} [\log p(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{Independent bivariate normals}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(p(\theta) || q(\theta))$ ?



**KL minimizers “fit inside” the second argument.**


# KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$$\mathcal{Q} = \{\text{Bivariate normals}\}$$

What is  $q^*(\theta) =$   
 $\underset{q \in \mathcal{Q}}{\text{argmin}} \left( - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \right)$ ?




# KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{Bivariate normals}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \left( - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \right)$ ?



**Without the entropy, the KL minimizer concentrates on the maximum of  $\log p(\theta)$ .**


# KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$$\mathcal{Q} = \{\text{Bivariate normals}\}$$

What is  $q^*(\theta) =$   
 $\underset{q \in \mathcal{Q}}{\operatorname{argmin}} \left( - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \right) ?$




# KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{Bivariate normals}\}$

What is  $q^*(\theta) =$   
 $\underset{q \in \mathcal{Q}}{\text{argmin}} \left( - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \right)$ ?



**Without  $\log p(\theta)$ , the KL minimizer is infinitely dispersed.**


# KL divergence exercises

$$\text{KL}(q(\theta) || p(\theta)) = -\mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{Point masses}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\operatorname{argmin}} \text{KL}(q(\theta) || p(\theta))$ ?






## KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{Point masses}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\operatorname{argmin}} \text{KL}(q(\theta) || p(\theta))$ ?



**Without a common dominating measure, the KL divergence is undefined.**

## KL divergence exercises


$$\text{KL}(q(\theta) || p(\theta)) =$$

$$- \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{BVN with small, fixed variance}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(q(\theta) || p(\theta))$ ?



## KL divergence exercises


$$\text{KL}(q(\theta) || p(\theta)) =$$

$$- \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$  = Correlated bivariate normal

$\mathcal{Q} = \{\text{BVN with small, fixed variance}\}$

What is  $q^*(\theta) = \underset{q \in \mathcal{Q}}{\operatorname{argmin}} \text{KL}(q(\theta) || p(\theta))$ ?



**Sufficiently concentrated distributions with constant entropy act like a point mass at the maximum of  $\log p(\theta)$ .**

- Luca Ambrogioni, Umut Güçlü, Yağmur Güçlütürk, Max Hinne, Eric Maris, and Marcel AJ van Gerven. Wasserstein variational inference. *arXiv preprint arXiv:1805.11284*, 2018.
- Alison L Gibbs and Francis Edward Su. On choosing and bounding probability metrics. *International statistical review*, 70(3):419–435, 2002.
- Solomon Kullback and Richard A Leibler. On information and sufficiency. *The annals of mathematical statistics*, 22(1):79–86, 1951.
- Yingzhen Li and Richard E Turner. Variational inference with rényi divergence. *stat*, 1050:6, 2016.
- Qiang Liu and Dilin Wang. Stein variational gradient descent: A general purpose bayesian inference algorithm. *arXiv preprint arXiv:1608.04471*, 2016.