

Locally Equivalent Weights for Bayesian MrP

Ryan Giordano, Alice Cima, Erin Hartman, Jared Murray, Avi Feller

UT Austin Statistics Seminar

September 2025



Are US non-voters becoming more Republican?

Blue Rose research says yes:

“Politically disengaged voters have become much more Republican, and because less-engaged voters swung away from [Democrats], an expanded electorate meant a more Republican electorate.”

(Blue Rose Research 2024)
(major professional pollsters)

On Data and Democracy says no:

“Claims of a decisive pro-Republican shift among the overall non-voting population are not supported by the most reliable, large-scale post-election data currently available.”

(Bonica et al. 2025)
(major professional researchers)

Are US non-voters becoming more Republican?

Blue Rose research says yes:

“Politically disengaged voters have become much more Republican, and because less-engaged voters swung away from [Democrats], an expanded electorate meant a more Republican electorate.”

(Blue Rose Research 2024)
(major professional pollsters)

On Data and Democracy says no:

“Claims of a decisive pro-Republican shift among the overall non-voting population are not supported by the most reliable, large-scale post-election data currently available.”

(Bonica et al. 2025)
(major professional researchers)

-
- The problem is very hard (it's difficult to accurately poll non-voters)
 - Different data sources
 - ★★★ **Different statistical methods**
 - Blue Rose uses Bayesian hierarchical modeling (MrP)
 - On Data and Democracy is using calibration weighting (CW)

Are US non-voters becoming more Republican?

Blue Rose research says yes:

“Politically disengaged voters have become much more Republican, and because less-engaged voters swung away from [Democrats], an expanded electorate meant a more Republican electorate.”

(Blue Rose Research 2024)
(major professional pollsters)

On Data and Democracy says no:

“Claims of a decisive pro-Republican shift among the overall non-voting population are not supported by the most reliable, large-scale post-election data currently available.”

(Bonica et al. 2025)
(major professional researchers)

-
- The problem is very hard (it’s difficult to accurately poll non-voters)
 - Different data sources
 - *** **Different statistical methods**
 - Blue Rose uses Bayesian hierarchical modeling (MrP)
 - On Data and Democracy is using calibration weighting (CW)

Our contribution

We define “MrP local equivalent weights” (MrPlew) that:

- Are easily computable from MCMC draws and standard software, and
- Provide MrP versions of key diagnostics that motivate calibration weighting.

⇒ **MrPlew provides direct comparisons between MrP and calibration weighting.**

- Introduce the statistical problem
 - Contrast CW and MrP
 - Prior work: Equivalent weights for linear models
 - Our key idea: Locally equivalent weights for non-linear models

- Introduce the statistical problem
 - Contrast CW and MrP
 - Prior work: Equivalent weights for linear models
 - Our key idea: Locally equivalent weights for non-linear models
- Locally equivalent weights for covariate balance
 - Describe covariate balance
 - Define MrPlew weights and connect them to covariate balance
 - Theoretical support
 - Example of real-world results


- Introduce the statistical problem
 - Contrast CW and MrP
 - Prior work: Equivalent weights for linear models
 - Our key idea: Locally equivalent weights for non-linear models
- Locally equivalent weights for covariate balance
 - Describe covariate balance
 - Define MrPlew weights and connect them to covariate balance
 - Theoretical support
 - Example of real-world results
- Other uses of locally equivalent weights
 - Partial pooling
 - The meaning of negative weights
 - Frequentist variance estimation
- Future directions

The basic problem


We have a survey population, for whom we observe:

- Covariates \mathbf{x} (e.g. race, gender, zip code, age, education level)
- Responses y (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe (\mathbf{x}_i, y_i) for $i = 1, \dots, N_S$




Observe \mathbf{x}_j for $j = 1, \dots, N_T$

The basic problem


We have a survey population, for whom we observe:

- Covariates \mathbf{x} (e.g. race, gender, zip code, age, education level)
- Responses y (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe (\mathbf{x}_i, y_i) for $i = 1, \dots, N_S$



Observe \mathbf{x}_j for $j = 1, \dots, N_T$


The problem is that the populations may be very different.

The basic problem


We have a survey population, for whom we observe:

- Covariates \mathbf{x} (e.g. race, gender, zip code, age, education level)
- Responses y (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe (\mathbf{x}_i, y_i) for $i = 1, \dots, N_S$



Observe \mathbf{x}_j for $j = 1, \dots, N_T$

The problem is that the populations may be very different.

Our survey results may be biased.

How can we use the covariates to say something about the target responses?

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target population y_j .

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target population y_j .

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- Choose “calibration weights” w_i
using only the regressors \mathbf{x}
(e.g. raking weights)

Bayesian hierarchical modeling (MrP)

- Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$,
choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$
(e.g. Hierarchical logistic regression)

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target population y_j .

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- Choose “calibration weights” w_i
using only the regressors \mathbf{x}
(e.g. raking weights)
- Take $\hat{\mu}_{\text{CW}} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$

Bayesian hierarchical modeling (MrP)

- Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$,
choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$
(e.g. Hierarchical logistic regression)
- Take $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$ and
 $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target population y_j .

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- Choose “calibration weights” w_i using only the regressors \mathbf{x} (e.g. raking weights)
- Take $\hat{\mu}_{CW} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$
- Dependence on y_i is clear

Bayesian hierarchical modeling (MrP)

- Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$, choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$ (e.g. Hierarchical logistic regression)
- Take $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$ and $\hat{\mu}_{MrP} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$
- Dependence on y_i very complicated (Typically via MCMC draws from $\mathcal{P}(\theta|\text{Survey data})$)

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target population y_j .

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- ▶ Choose “calibration weights” w_i using only the regressors \mathbf{x} (e.g. raking weights)

- ▶ Take $\hat{\mu}_{\text{CW}} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$

- ▶ Dependence on y_i is clear

- ▶ Weights give interpretable diagnostics:

- Frequentist variability
- Partial pooling
- Regressor balance

Bayesian hierarchical modeling (MrP)

- ▶ Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$, choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$

(e.g. Hierarchical logistic regression)

- ▶ Take $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$ and $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$

- ▶ Dependence on y_i very complicated (Typically via MCMC draws from $\mathcal{P}(\theta|\text{Survey data})$)

- ▶ **Black box**

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target population y_j .

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- ▶ Choose “calibration weights” w_i using only the regressors \mathbf{x} (e.g. raking weights)

- ▶ Take $\hat{\mu}_{CW} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$

- ▶ Dependence on y_i is clear

- ▶ Weights give interpretable diagnostics:

- Frequentist variability
- Partial pooling
- Regressor balance

Bayesian hierarchical modeling (MrP)

- ▶ Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$, choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$ (e.g. Hierarchical logistic regression)

- ▶ Take $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$ and $\hat{\mu}_{MrP} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$

- ▶ Dependence on y_i very complicated (Typically via MCMC draws from $\mathcal{P}(\theta|\text{Survey data})$)

▶ Black box

- ← We open this box, providing analogues of all these diagnostics

Gelman (2007b) observes that MrP is a CW estimator when one uses linear regression to form \hat{y} :

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j = \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\mathbf{x}_j^T \hat{\boldsymbol{\theta}}}_{\text{Linear in } y_i}$$

Most existing literature on comparing CW and MrP focus on such linear models.¹

Let's spend some time discussing why it is reasonable to even attempt such a thing as forming approximate equivalent weights for non-linear estimators.

¹For example, Gelman (2007b), B., F., and H. (2021), and Chattopadhyay and Zubizarreta (2023).

Gelman (2007b) observes that MrP is a CW estimator when one uses linear regression to form \hat{y} :

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j = \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\mathbf{x}_j^\top \hat{\boldsymbol{\theta}}}_{\text{Linear in } y_i}$$

Most existing literature on comparing CW and MrP focus on such linear models.¹

But what if you use a non-linear link function? Or a hierarchical model?

“It would also be desirable to use nonlinear methods ... but then it would seem difficult to construct even approximately equivalent weights. Weighting and fully nonlinear models would seem to be completely incompatible methods.” — (Gelman 2007a)

Let’s spend some time discussing why it is reasonable to even attempt such a thing as forming approximate equivalent weights for non-linear estimators.

¹For example, Gelman (2007b), B., F., and H. (2021), and Chattopadhyay and Zubizarreta (2023).

Equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose $\mathbf{x} \in \mathcal{X}$ is discrete and saturated.

Then MrP is a CW estimator.

Equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose $\mathbf{x} \in \mathcal{X}$ is discrete and saturated.

Then MrP is a CW estimator.

- Let \bar{y}_S^c denote the survey average among $\mathbf{x} = c$ for $c \in \mathcal{X}$
- For $\mathbf{x} = c$, $m(\hat{\theta}^\top \mathbf{x}) = \bar{y}_S^c$
- Let N_S^c (or N_S^c) denote the # of survey (or target) observations with $\mathbf{x}_n = c$.

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_T} \sum_{c \in \mathcal{X}} \underbrace{N_T^c \bar{y}_S^c}_{\text{Linear in } y_i} = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i^{\text{MrP}} y_i$$

$$\text{For } w_i^{\text{MrP}} = \frac{N_T^c / N_T}{N_S^c / N_S} \text{ when } \mathbf{x}_i = c.$$

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

Then MrP is *approximately* a CW estimator.

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

Then MrP is *approximately* a CW estimator.

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$$

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

Then MrP is *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} \quad (\text{Law of large numbers})\end{aligned}$$

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

Then MrP is *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)}\end{aligned}$$

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

Then MrP is *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)}\end{aligned}$$

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

Then MrP is *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)} \\ &\approx \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i m(\mathbf{x}_i^\top \hat{\theta}) && \text{(Law of large numbers)}\end{aligned}$$

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

Then MrP is *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}_{\text{MrP}} &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)} \\ &\approx \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i m(\mathbf{x}_i^\top \hat{\theta}) && \text{(Law of large numbers)} \\ &= \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i y_i && \text{(Property of exponential family MLEs)}\end{aligned}$$

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \alpha^\top \mathbf{x}_i y_i + \text{Small error}$$

We don't observe $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$, so it's hard to estimate α directly.

²Krantz and Parks 2012; G., Stephenson, et al. 2019.

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \alpha^\top \mathbf{x}_i y_i + \text{Small error}$$

We don't observe $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$, so it's hard to estimate α directly.

Key idea (informal)

If $\hat{\mu}_{\text{MrP}} \approx \frac{1}{N_S} \sum_{i=1}^{N_S} w_i^{\text{MrP}} y_i$ for some w_i^{MrP} , then

$$w_i^{\text{MrP}} \approx \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$$

²Krantz and Parks 2012; G., Stephenson, et al. 2019.

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.
- Suppose there exists α such that $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$.

$$\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \alpha^\top \mathbf{x}_i y_i + \text{Small error}$$

We don't observe $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$, so it's hard to estimate α directly.

Key idea (informal)

If $\hat{\mu}_{\text{MrP}} \approx \frac{1}{N_S} \sum_{i=1}^{N_S} w_i^{\text{MrP}} y_i$ for some w_i^{MrP} , then

$$w_i^{\text{MrP}} \approx \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$$

For logistic regression, could compute and analyze $\frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$ using the implicit function theorem.²

²Krantz and Parks 2012; G., Stephenson, et al. 2019.

Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$.
- Set a hierarchical prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$, use MCMC to draw from $\mathcal{P}(\theta|\text{Survey data})$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[m(\mathbf{x}_j^\top \theta) \right]$.

³Gustafson 1996; G., Broderick, and Jordan 2018.

Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$.
- Set a hierarchical prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$, use MCMC to draw from $\mathcal{P}(\theta|\text{Survey data})$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[m(\mathbf{x}_j^\top \theta) \right]$.

MrP locally equivalent weights (MrPlew)

For new data \tilde{Y}_S , form a series expansion

$$\hat{\mu}_{\text{MrP}}(\tilde{Y}_S) \approx \hat{\mu}_{\text{MrP}}(Y_S) + \sum_{i=1}^{N_S} w_i^{\text{MrP}} (\tilde{y}_i - y_i) \quad \text{where} \quad w_i^{\text{MrP}} := \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}.$$

Our task is to rigorously show that even such local weights can be used diagnostically.

³Gustafson 1996; G., Broderick, and Jordan 2018.

Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$.
- Set a hierarchical prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$, use MCMC to draw from $\mathcal{P}(\theta|\text{Survey data})$.
- MrP is $\hat{\mu}_{\text{MrP}} = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[m(\mathbf{x}_j^\top \theta) \right]$.

MrP locally equivalent weights (MrPlew)

For new data \tilde{Y}_S , form a series expansion

$$\hat{\mu}_{\text{MrP}}(\tilde{Y}_S) \approx \hat{\mu}_{\text{MrP}}(Y_S) + \sum_{i=1}^{N_S} w_i^{\text{MrP}} (\tilde{y}_i - y_i) \quad \text{where} \quad w_i^{\text{MrP}} := \frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}.$$

Our task is to rigorously show that even such local weights can be used diagnostically.

- If $\hat{\mu}_{\text{MrP}}$ is linear in Y_S , this definition recovers the true weights
- Whether or not $\hat{\mu}_{\text{MrP}}$ is linear in Y_S , this definition is valid “locally” for \tilde{Y}_S “nearby” Y_S .
- For MCMC, compute and analyze $\frac{\partial \hat{\mu}_{\text{MrP}}}{\partial y_i}$ using Bayesian sensitivity analysis.³

³Gustafson 1996; G., Broderick, and Jordan 2018.

The weights can look very different!

Does this mean anything? Are the differences important?




Figure 1: Comparison between raking and MrPlew weights for a particular example

Note that there was no talk of correct specification for the data you have.

That was a foregone conclusion when we started looking at equivalent weights!

How do you perform model checking with sensitivity analysis?

Existing methods evaluate whether the analysis changes “a lot” when you:

- Parametrically perturb the model (e.g. fit a richer model class)
- Non-parametrically perturb the data (e.g. produce gross outliers)

The problem is:

- How much is “a lot”?
- Non-parametric data perturbations are hard to reason about
- It’s hard to say whether parametric model changes are enough

Instead, we

- Parametrically perturb the data
- Observe whether our model could detect the change
- Know exactly the expected change (don’t have to decide on what “a lot” means)
- Easy to reason about whether the data perturbation is reasonable
- Don’t need to propose an alternative model, instead study the model you have

Student contributions and future work:

- **Alice Cima** contributed significantly to this work
- **Vladimir Palmin** is working on extending MrPlew to lme4
- **Sequoia Andrade** is working on generalizing to other local sensitivity checks
- **Lucas Schwengber** is working on novel flow-based techniques for local sensitivity




Alice Cima

No picture!
Vladimir Palmin



Sequoia Andrade



Lucas Schwengber

- 
- B., Eli, Avi F., and Erin H. (2021). *Multilevel calibration weighting for survey data*. arXiv: 2102.09052 [stat.ME].
- 
- Blue Rose Research (2024). *2024 Election Retrospective Presentation*. <https://data.blueroseresearch.org/2024retro-download>. Accessed on 2024-10-26.
- 
- Bonica, A. et al. (Apr. 2025). *Did Non-Voters Really Flip Republican in 2024? The Evidence Says No*. <https://data4democracy.substack.com/p/did-non-voters-really-flip-republican>.
- 
- Chattopadhyay, A. and J. Zubizarreta (2023). “On the implied weights of linear regression for causal inference”. In: *Biometrika* 110.3, pp. 615–629.
- 
- G., T. Broderick, and M. I. Jordan (2018). “Covariances, robustness and variational bayes”. In: *Journal of machine learning research* 19.51.
- 
- G., W. Stephenson, et al. (2019). “A swiss army infinitesimal jackknife”. In: *The 22nd International Conference on Artificial Intelligence and Statistics*. PMLR, pp. 1139–1147.
- 
- Gelman, A. (2007a). “Rejoinder: Struggles with survey weighting and regression modelling”. In: *Statistical Science* 22.2, pp. 184–188.
- 
- (2007b). “Struggles with survey weighting and regression modeling”. In.
- 
- Gustafson, P. (1996). “Local sensitivity of posterior expectations”. In: *The Annals of Statistics* 24.1, pp. 174–195.
- 
- Krantz, S. and H. Parks (2012). *The Implicit Function Theorem: History, Theory, and Applications*. Springer Science & Business Media.