

Locally Equivalent Weights for Bayesian MrP

Ryan Giordano, Alice Cima, Erin Hartman, Jared Murray, Avi Feller

University of British Columbia Statistics Seminar

October 2025



Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

But the high level idea can be extended much more widely:

1. Assume your initial model was accurate
2. Select some perturbation your model should be able to capture
3. Use local sensitivity to detect whether the change is what you expect
4. If the change is not what you expect, either (1) or (2) was wrong

Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

But the high level idea can be extended much more widely:

1. Assume your initial model was accurate
2. Select some perturbation your model should be able to capture
3. Use local sensitivity to detect whether the change is what you expect
4. If the change is not what you expect, either (1) or (2) was wrong

Checks of this form give generalized versions of many standard linear model diagnostics.

Regression

Regression

General models

Regression

General models

Consistency /
Unbiased

$$y = \theta^\top \mathbf{x} + \varepsilon$$

$$\tilde{y} = (\theta + \delta)^\top \mathbf{x} + \varepsilon$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

Regression

Consistency /
Unbiased

$$y = \theta^\top \mathbf{x} + \varepsilon$$

$$\tilde{y} = (\theta + \delta)^\top \mathbf{x} + \varepsilon$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

General models

$$y = f(\mathbf{x}, \varepsilon, \theta)$$

$$\tilde{y} = f(\mathbf{x}, \varepsilon, \theta + \delta)$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

Regression

Consistency /
Unbiased

$$y = \theta^\top \mathbf{x} + \varepsilon$$

$$\tilde{y} = (\theta + \delta)^\top \mathbf{x} + \varepsilon$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

General models

$$y = f(\mathbf{x}, \varepsilon, \theta)$$

$$\tilde{y} = f(\mathbf{x}, \varepsilon, \theta + \delta)$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

Exogenous
residuals

$$y = \theta^\top \mathbf{x} + \varepsilon$$

$$\tilde{y} = y + \varepsilon z$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y)$$

Regression

General models

Consistency /
Unbiased

$$y = \theta^\top \mathbf{x} + \varepsilon$$

$$\tilde{y} = (\theta + \delta)^\top \mathbf{x} + \varepsilon$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

$$y = f(\mathbf{x}, \varepsilon, \theta)$$

$$\tilde{y} = f(\mathbf{x}, \varepsilon, \theta + \delta)$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

Exogenous
residuals

$$y = \theta^\top \mathbf{x} + \varepsilon$$

$$\tilde{y} = y + \varepsilon z$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y)$$

$$y \sim \mathcal{P}(y|\mathbf{x}) \text{ and } \mathcal{P}(\mathbf{x}) = w$$

$$\tilde{w} = w + \delta z$$

$$\hat{\theta}(\tilde{w}) \stackrel{\text{check}}{=} \hat{\theta}(w)$$

Regression

General models

Consistency /
Unbiased

$$y = \theta^T \mathbf{x} + \varepsilon$$

$$\tilde{y} = (\theta + \delta)^T \mathbf{x} + \varepsilon$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

$$y = f(\mathbf{x}, \varepsilon, \theta)$$

$$\tilde{y} = f(\mathbf{x}, \varepsilon, \theta + \delta)$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

Exogenous
residuals

$$y = \theta^T \mathbf{x} + \varepsilon$$

$$\tilde{y} = y + \varepsilon z$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y)$$

$$y \sim \mathcal{P}(y|\mathbf{x}) \text{ and } \mathcal{P}(\mathbf{x}) = w$$

$$\tilde{w} = w + \delta z$$

$$\hat{\theta}(\tilde{w}) \stackrel{\text{check}}{=} \hat{\theta}(w)$$

Fisher
information

$$\mathcal{I} := \text{Fisher information}$$

$$\Sigma := \text{Score covariance}$$

$$\mathcal{I}^{-1} \stackrel{\text{check}}{=} \Sigma$$

Regression

General models

Consistency /
Unbiased

$$y = \theta^T \mathbf{x} + \varepsilon$$

$$\tilde{y} = (\theta + \delta)^T \mathbf{x} + \varepsilon$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

$$y = f(\mathbf{x}, \varepsilon, \theta)$$

$$\tilde{y} = f(\mathbf{x}, \varepsilon, \theta + \delta)$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

Exogenous
residuals

$$y = \theta^T \mathbf{x} + \varepsilon$$

$$\tilde{y} = y + \varepsilon z$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y)$$

$$y \sim \mathcal{P}(y|\mathbf{x}) \text{ and } \mathcal{P}(\mathbf{x}) = w$$

$$\tilde{w} = w + \delta z$$

$$\hat{\theta}(\tilde{w}) \stackrel{\text{check}}{=} \hat{\theta}(w)$$

Fisher
information

$\mathcal{I} :=$ Fisher information

$\Sigma :=$ Score covariance

$$\mathcal{I}^{-1} \stackrel{\text{check}}{=} \Sigma$$

$$y \sim \mathcal{P}(y|\theta)$$

$\tilde{y} \sim$ Importance sample y

$$\text{using } \tilde{w} = \frac{\mathcal{P}(y|\hat{\theta} + \delta)}{\mathcal{P}(y|\hat{\theta})}$$

$$\hat{\theta}(\tilde{w}) \stackrel{\text{check}}{=} \hat{\theta}(1) + \delta$$

Regression

General models

Consistency /
Unbiased

$$y = \theta^T \mathbf{x} + \varepsilon$$

$$\tilde{y} = (\theta + \delta)^T \mathbf{x} + \varepsilon$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

$$y = f(\mathbf{x}, \varepsilon, \theta)$$

$$\tilde{y} = f(\mathbf{x}, \varepsilon, \theta + \delta)$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y) + \delta$$

Exogenous
residuals

$$y = \theta^T \mathbf{x} + \varepsilon$$

$$\tilde{y} = y + \varepsilon z$$

$$\hat{\theta}(\tilde{y}) \stackrel{\text{check}}{=} \hat{\theta}(y)$$

$$y \sim \mathcal{P}(y|\mathbf{x}) \text{ and } \mathcal{P}(\mathbf{x}) = w$$

$$\tilde{w} = w + \delta z$$

$$\hat{\theta}(\tilde{w}) \stackrel{\text{check}}{=} \hat{\theta}(w)$$

Fisher
information

$\mathcal{I} :=$ Fisher information

$\Sigma :=$ Score covariance

$$\mathcal{I}^{-1} \stackrel{\text{check}}{=} \Sigma$$

$$y \sim \mathcal{P}(y|\theta)$$

$\tilde{y} \sim$ Importance sample y

$$\text{using } \tilde{w} = \frac{\mathcal{P}(y|\hat{\theta} + \delta)}{\mathcal{P}(y|\hat{\theta})}$$

$$\hat{\theta}(\tilde{w}) \stackrel{\text{check}}{=} \hat{\theta}(1) + \delta$$

Student contributions and ongoing work:


- **Vladimir Palmin** is working on extending MrPlew to `lme4`
- **Sequoia Andrade** is working on generalizing to other local sensitivity checks
- **Lucas Schwengber** is working on novel flow-based techniques for local sensitivity
- **(Currently recruiting!)** Doubly-robust Bayesian Hierarchical MrP



Vladimir Palmin



Sequoia Andrade



Lucas Schwengber

Preprint and R package (hopefully) coming soon!

