

# Discussion of “The Shrinkage-Delinkage Trade-off”

Restricted variational families (mean field) can lead to poor posterior approximations.

Two very common responses to this problem are:

- “Machine learning”: Ignore it (evaluate using some other criteria, like prediction)
- “Modern VI”: Use more expressive families (at a computational cost)

This paper tries to understand *how* the restricted family goes wrong. \*fire emoji\*

# Discussion of “The Shrinkage-Delinkage Trade-off”

Restricted variational families (mean field) can lead to poor posterior approximations.

Two very common responses to this problem are:

- “Machine learning”: Ignore it (evaluate using some other criteria, like prediction)
- “Modern VI”: Use more expressive families (at a computational cost)

This paper tries to understand *how* the restricted family goes wrong. \*fire emoji\*

---

This paper’s most (initially) surprising conclusion is probably this:

## Theorem 3.6, paraphrased

Let the target distribution have the constant  $\varepsilon$ -correlation matrix.

As the dimension  $n$  of the matrix goes to infinity:

- Each marginal mean field variance is wrong by  $\varepsilon$
- The per-component entropy gap  $\rightarrow 0$

# Discussion of “The Shrinkage-Delinkage Trade-off”

Restricted variational families (mean field) can lead to poor posterior approximations.

Two very common responses to this problem are:

- “Machine learning”: Ignore it (evaluate using some other criteria, like prediction)
- “Modern VI”: Use more expressive families (at a computational cost)

This paper tries to understand *how* the restricted family goes wrong. \*fire emoji\*

---

This paper’s most (initially) surprising conclusion is probably this:

## Theorem 3.6, paraphrased

Let the target distribution have the constant  $\varepsilon$ -correlation matrix.

As the dimension  $n$  of the matrix goes to infinity:

- Each marginal mean field variance is wrong by  $\varepsilon$
- The per-component entropy gap  $\rightarrow 0$

The key is “per-component entropy gap” means **entropy difference** /  $n$ .

In fact, the entropy gap is  $O(\log n) \rightarrow \infty$ . **Why is  $n$  the “right” scaling?**

# Discussion of “The Shrinkage-Delinkage Trade-off”

Restricted variational families (mean field) can lead to poor posterior approximations.

Two very common responses to this problem are:

- “Machine learning”: Ignore it (evaluate using some other criteria, like prediction)
- “Modern VI”: Use more expressive families (at a computational cost)

This paper tries to understand *how* the restricted family goes wrong. \*fire emoji\*

---

This paper’s most (initially) surprising conclusion is probably this:

## Theorem 3.6, paraphrased

Let the target distribution have the constant  $\varepsilon$ -correlation matrix.

As the dimension  $n$  of the matrix goes to infinity:

- Each marginal mean field variance is wrong by  $\varepsilon$
- The per-component entropy gap  $\rightarrow 0$

The key is “per-component entropy gap” means **entropy difference** /  $n$ .

In fact, the entropy gap is  $O(\log n) \rightarrow \infty$ . **Why is  $n$  the “right” scaling?**

Why do the relative values across dimensions of the entropy gap matter?

**It’s clear why variance is useful. Less so the entropy gap, especially as  $n$  changes.**