


# **Approximate data deletion and replication with the Bayesian influence function**

---

Ryan Giordano (rgiordano@berkeley.edu, UC Berkeley), Tamara Broderick (MIT)  
**Stanford Statistics Seminar May 2024**

# Economist 2016 Election Model [Gelman and Heidemanns, 2020]



A time series model to predict the 2016 US presidential election outcome from polling data.


Model:

- $X = x_1, \dots, x_N =$  Polling data ( $N = 361$ ).
- $\theta =$  Lots of random effects (day, pollster, etc.)
- $f(\theta) =$  Democratic % of vote on election day

Typically, we compute Markov chain Monte Carlo (MCMC) draws from the posterior  $p(\theta|X)$ .

We want to know  $\mathbb{E}_{p(\theta|X)} [f(\theta)]$ .

# Economist 2016 Election Model [Gelman and Heidemanns, 2020]



A time series model to predict the 2016 US presidential election outcome from polling data.

Model:

- $X = x_1, \dots, x_N =$  Polling data ( $N = 361$ ).
- $\theta =$  Lots of random effects (day, pollster, etc.)
- $f(\theta) =$  Democratic % of vote on election day

Typically, we compute Markov chain Monte Carlo (MCMC) draws from the posterior  $p(\theta|X)$ .


We want to know  $\mathbb{E}_{p(\theta|X)} [f(\theta)]$ .

The people who responded to the polls were randomly selected.

If we had selected a different random sample, how much would our estimate have changed?

**Idea:** Re-fit with bootstrap samples of data [Huggins and Miller, 2023]

# Economist 2016 Election Model [Gelman and Heidemanns, 2020]



A time series model to predict the 2016 US presidential election outcome from polling data.

Model:

- $X = x_1, \dots, x_N =$  Polling data ( $N = 361$ ).
- $\theta =$  Lots of random effects (day, pollster, etc.)
- $f(\theta) =$  Democratic % of vote on election day

Typically, we compute Markov chain Monte Carlo (MCMC) draws from the posterior  $p(\theta|X)$ .

We want to know  $\mathbb{E}_{p(\theta|X)} [f(\theta)]$ .

The people who responded to the polls were randomly selected.



If we had selected a different random sample, how much would our estimate have changed?

**Idea:** Re-fit with bootstrap samples of data [Huggins and Miller, 2023]

**Problem:** Each MCMC run takes about 10 hours (Stan, six cores).



# Results

Proposal: Use full-data posterior draws to form a linear approximation to *data reweightings*.



# Results

Proposal: Use full-data posterior draws to form a linear approximation to *data reweightings*.



Compute time for 100 bootstraps: 51 days

Compute time for the linear approximation: Seconds  
(But note the approximation has some error)

- Data reweighting
  - Write the change in the posterior expectation as **linear component** + **error**
  - The **linear component** can be computed from a single run of MCMC

- Data reweighting
  - Write the change in the posterior expectation as **linear component** + **error**
  - The **linear component** can be computed from a single run of MCMC
- Finite-dimensional problems with posteriors which concentrate asymptotically
  - As  $N \rightarrow \infty$ , the linear component provides an arbitrarily good approximation
  - Consistent variance estimates via a uniform Bernstein–von Mises theorem



- Data reweighting
  - Write the change in the posterior expectation as **linear component** + **error**
  - The **linear component** can be computed from a single run of MCMC
- Finite-dimensional problems with posteriors which concentrate asymptotically
  - As  $N \rightarrow \infty$ , the linear component provides an arbitrarily good approximation
  - Consistent variance estimates via a uniform Bernstein–von Mises theorem
- High-dimensional problems
  - The linear component is the same order as the error
  - Even for parameters which concentrate, even as  $N \rightarrow \infty$
  - Study the variance estimates via a Bayesian von–Mises expansion

- Data reweighting
  - Write the change in the posterior expectation as **linear component** + **error**
  - The **linear component** can be computed from a single run of MCMC
- Finite-dimensional problems with posteriors which concentrate asymptotically
  - As  $N \rightarrow \infty$ , the linear component provides an arbitrarily good approximation
  - Consistent variance estimates via a uniform Bernstein–von Mises theorem
- High-dimensional problems
  - The linear component is the same order as the error
  - Even for parameters which concentrate, even as  $N \rightarrow \infty$
  - Study the variance estimates via a Bayesian von–Mises expansion
- Some implications and future work

## Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .

$$\ell_n(\theta) := \log p(x_n|\theta) \qquad \log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$

## Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .

$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$

Original weights:



# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .


$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$

Original weights:



Leave-one-out weights:



# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .


$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$


Original weights:



Leave-one-out weights:



Bootstrap weights:



# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .


$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$



Original weights:



Leave-one-out weights:



Bootstrap weights:



# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .


$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$



Original weights:



Leave-one-out weights:



Bootstrap weights:





# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .


$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$



Original weights:



Leave-one-out weights:



Bootstrap weights:



# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .

$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$


Original weights:



Leave-one-out weights:



Bootstrap weights:



# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .

$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$


Original weights:



Leave-one-out weights:



Bootstrap weights:



# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .

$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$



Original weights:



Leave-one-out weights:



Bootstrap weights:



# Data re-weighting.

Augment the problem with *data weights*  $w_1, \dots, w_N$ . We can write  $\mathbb{E}_{p(\theta|X,w)}[f(\theta)]$ .


$$\ell_n(\theta) := \log p(x_n|\theta)$$

$$\log p(X|\theta, w) = \sum_{n=1}^N w_n \ell_n(\theta)$$



Original weights:



Leave-one-out weights:



Bootstrap weights:



The re-scaled slope  $N\psi_n$  is known as the “influence function” at data point  $x_n$ .

$$\mathbb{E}_{p(\theta|X,w)}[f(\theta)] - \mathbb{E}_{p(\theta|X)}[f(\theta)] = \sum_{n=1}^N \psi_n(w_n - 1) + \mathcal{E}(w_n)$$

How can we use the approximation?

Assume the **slope** is computable and **error** is small.

$$\mathbb{E}_{p(\theta|X,w)}[f(\theta)] - \mathbb{E}_{p(\theta|X)}[f(\theta)] = \sum_{n=1}^N \psi_n(w_n - 1) + \mathcal{E}(w_n)$$

How can we use the approximation?

Assume the **slope** is computable and **error** is small.

$$\mathbb{E}_{p(\theta|X,w)}[f(\theta)] - \mathbb{E}_{p(\theta|X)}[f(\theta)] = \sum_{n=1}^N \psi_n(w_n - 1) + \mathcal{E}(w_n)$$

**Bootstrap.** Draw bootstrap weights  $w \sim p(w) = \text{Multinomial}(N, N^{-1})$ .

$$\text{Bootstrap variance} = \text{Var}_{p(w)} \left( \mathbb{E}_{p(\theta|X,w)}[f(\theta)] \right)$$

How can we use the approximation?

Assume the **slope** is computable and **error** is small.

$$\mathbb{E}_{p(\theta|X,w)}[f(\theta)] - \mathbb{E}_{p(\theta|X)}[f(\theta)] = \sum_{n=1}^N \psi_n(w_n - 1) + \mathcal{E}(w_n)$$

**Bootstrap.** Draw bootstrap weights  $w \sim p(w) = \text{Multinomial}(N, N^{-1})$ .

$$\begin{aligned} \text{Bootstrap variance} &= \text{Var}_{p(w)} \left( \mathbb{E}_{p(\theta|X,w)}[f(\theta)] \right) \\ &= \text{Var}_{p(w)} \left( \sum_{n=1}^N \psi_n(w_n - 1) + \mathcal{E}(w_n) \right) \\ &= \frac{1}{N^2} \sum_{n=1}^N \left( \psi_n - \bar{\psi} \right)^2 + \text{Term involving } \mathcal{E}(w_n) \text{ for } n = 1, \dots, N \\ &\approx \frac{1}{N} \underbrace{\left( \frac{1}{N} \sum_{n=1}^N \left( \psi_n - \bar{\psi} \right)^2 \right)}_{\text{"Infinitesimal jackknife variance estimate"}} \end{aligned}$$



## Expressions for the slope and error

How to compute the slopes  $\psi_n$ ? How large is the error  $\mathcal{E}(w)$ ?

For simplicity, let us consider a single weight for the moment.

$$\mathbb{E}_{p(\theta|X, w_n)} [f(\theta)] - \mathbb{E}_{p(\theta|X)} [f(\theta)] = \psi_n(w_n - 1) + \mathcal{E}(w_n)$$

## Expressions for the slope and error

How to compute the slopes  $\psi_n$ ? How large is the error  $\mathcal{E}(w)$ ?

For simplicity, let us consider a single weight for the moment.

$$\mathbb{E}_{p(\theta|X, w_n)} [f(\theta)] - \mathbb{E}_{p(\theta|X)} [f(\theta)] = \psi_n(w_n - 1) + \mathcal{E}(w_n)$$

Let an overbar denote “posterior–mean zero.” For example,  $\bar{f}(\theta) := f(\theta) - \mathbb{E}_{p(\theta|X)} [f(\theta)]$ .

By dominated convergence and the mean value theorem, for some  $\tilde{w}_n \in [0, w_n]$ :

$$\psi_n = \underbrace{\mathbb{E}_{p(\theta|X)} [\bar{f}(\theta) \bar{\ell}_n(\theta)]}_{\text{Estimatable with MCMC!}} \quad \mathcal{E}(w_n) = \frac{1}{2} \underbrace{\mathbb{E}_{p(\theta|X, \tilde{w}_n)} [\bar{f}(\theta) \bar{\ell}_n(\theta) \bar{\ell}_n(\theta)]}_{\text{Cannot compute directly (don't know } \tilde{w})} (w_n - 1)^2$$

# Expressions for the slope and error

How to compute the slopes  $\psi_n$ ? How large is the error  $\mathcal{E}(w)$ ?

For simplicity, let us consider a single weight for the moment.

$$\mathbb{E}_{p(\theta|X, w_n)} [f(\theta)] - \mathbb{E}_{p(\theta|X)} [f(\theta)] = \psi_n(w_n - 1) + \mathcal{E}(w_n)$$

Let an overbar denote “posterior–mean zero.” For example,  $\bar{f}(\theta) := f(\theta) - \mathbb{E}_{p(\theta|X)} [f(\theta)]$ .

By dominated convergence and the mean value theorem, for some  $\tilde{w}_n \in [0, w_n]$ :

$$\begin{aligned} \psi_n &= \underbrace{\mathbb{E}_{p(\theta|X)} [\bar{f}(\theta) \bar{\ell}_n(\theta)]}_{\text{Estimatable with MCMC!}} & \mathcal{E}(w_n) &= \frac{1}{2} \underbrace{\mathbb{E}_{p(\theta|X, \tilde{w}_n)} [\bar{f}(\theta) \bar{\ell}_n(\theta) \bar{\ell}_n(\theta)]}_{\text{Cannot compute directly (don't know } \tilde{w})} (w_n - 1)^2 \\ &= O_p(N^{-1}) \text{ under posterior concentration} & &= O_p(N^{-2}) \text{ under posterior concentration} \end{aligned}$$

# Expressions for the slope and error

How to compute the slopes  $\psi_n$ ? How large is the error  $\mathcal{E}(w)$ ?

For simplicity, let us consider a single weight for the moment.

$$\mathbb{E}_{p(\theta|X, w_n)} [f(\theta)] - \mathbb{E}_{p(\theta|X)} [f(\theta)] = \psi_n (w_n - 1) + \mathcal{E}(w_n)$$

Let an overbar denote “posterior–mean zero.” For example,  $\bar{f}(\theta) := f(\theta) - \mathbb{E}_{p(\theta|X)} [f(\theta)]$ .

By dominated convergence and the mean value theorem, for some  $\tilde{w}_n \in [0, w_n]$ :

$$\begin{aligned} \psi_n &= \underbrace{\mathbb{E}_{p(\theta|X)} [\bar{f}(\theta) \bar{\ell}_n(\theta)]}_{\text{Estimatable with MCMC!}} & \mathcal{E}(w_n) &= \frac{1}{2} \underbrace{\mathbb{E}_{p(\theta|X, \tilde{w}_n)} [\bar{f}(\theta) \bar{\ell}_n(\theta) \bar{\ell}_n(\theta)]}_{\text{Cannot compute directly (don't know } \tilde{w})} (w_n - 1)^2 \\ &= O_p(N^{-1}) \text{ under posterior concentration} & &= O_p(N^{-2}) \text{ under posterior concentration} \end{aligned}$$

## Theorem 1 [Giordano and Broderick, 2023] (paraphrase):

If the posterior  $p(\theta|X)$  “concentrates” (e.g. as in the Bernstein–von Mises theorem),<sup>a</sup> then

$$w_n \mapsto N \left( \mathbb{E}_{p(\theta|X, w_n)} [f(\theta)] - \mathbb{E}_{p(\theta|X)} [f(\theta)] \right)$$

becomes linear as  $N \rightarrow \infty$ , with slope  $\lim_{N \rightarrow \infty} \psi_n$ .

<sup>a</sup>Existing results are sufficient for a *particular weight* [Kass et al., 1990]. Giordano and Broderick [2023] proves that the result holds when averaged over all weights, as needed for variance estimation.

## Variance consistency theorem

How do the results for a single weight translate into variance estimates?

$$\text{Var}_{p(w)} \left( \mathbb{E}_{p(\theta|X,w)} [f(\theta)] \right) = \frac{1}{N^2} \sum_{n=1}^N \left( \psi_n - \bar{\psi} \right)^2 + \text{Term involving } \mathcal{E}(w_n) \text{ for } n = 1, \dots, N$$

# Variance consistency theorem

How do the results for a single weight translate into variance estimates?

$$\text{Var}_{p(w)} \left( \mathbb{E}_{p(\theta|X,w)} [f(\theta)] \right) = \frac{1}{N^2} \sum_{n=1}^N \left( \psi_n - \bar{\psi} \right)^2 + \text{Term involving } \mathcal{E}(w_n) \text{ for } n = 1, \dots, N$$

- Assume (sketch): A well-behaved MAP *maximum a posteriori* estimator  $\hat{\theta}$  exists.
  - The dimension of  $\theta$  is fixed as  $N \rightarrow \infty$
  - The expected log likelihood has a strict maximum at  $\theta_\infty$
  - The observed log likelihood satisfies  $\hat{\theta} \rightarrow \theta_\infty$
  - The expected log likelihood Hessian is negative definite at  $\theta_\infty$
- Assume (sketch): We can apply standard asymptotics.
  - The log prior and log likelihood are four times continuously differentiable
  - The prior is proper, and a technical set of squared expectations are finite
  - The log likelihood derivatives are dominated by a square-integrable envelope function in a neighborhood of  $\theta_\infty$ .

# Variance consistency theorem

How do the results for a single weight translate into variance estimates?

$$\text{Var}_{p(w)} \left( \mathbb{E}_{p(\theta|X, w)} [f(\theta)] \right) = \frac{1}{N^2} \sum_{n=1}^N \left( \psi_n - \bar{\psi} \right)^2 + \text{Term involving } \mathcal{E}(w_n) \text{ for } n = 1, \dots, N$$

- Assume (sketch): A well-behaved MAP *maximum a posteriori* estimator  $\hat{\theta}$  exists.
  - The dimension of  $\theta$  is fixed as  $N \rightarrow \infty$
  - The expected log likelihood has a strict maximum at  $\theta_\infty$
  - The observed log likelihood satisfies  $\hat{\theta} \rightarrow \theta_\infty$
  - The expected log likelihood Hessian is negative definite at  $\theta_\infty$
- Assume (sketch): We can apply standard asymptotics.
  - The log prior and log likelihood are four times continuously differentiable
  - The prior is proper, and a technical set of squared expectations are finite
  - The log likelihood derivatives are dominated by a square-integrable envelope function in a neighborhood of  $\theta_\infty$ .

**Theorem 2 [Giordano and Broderick, 2023]:** Under the above assumptions,

$$\sqrt{N} \left( \mathbb{E}_{p(\theta|X)} [g(\theta)] - g(\theta_\infty) \right) \xrightarrow[N \rightarrow \infty]{dist} \mathcal{N}(0, V^g) \quad [\text{Kleijn and Van der Vaart, 2012}]$$

$$\text{and } V^{\text{IJ}} := \frac{1}{N} \sum_{n=1}^N \left( \psi_n - \bar{\psi} \right)^2 \xrightarrow[N \rightarrow \infty]{prob} V^g. \quad (\text{Our contribution})$$

## Negative binomial experiment

Example: Negative binomial models with an unknown parameter  $\gamma$ .

For  $n = 1, \dots, N$  let  $x_n | \gamma \stackrel{iid}{\sim} \text{NegativeBinomial}(\alpha, \gamma)$  for fixed  $\alpha$ .

$$\text{Write } \log p(X | \lambda, \gamma, w) = \sum_{n=1}^N w_n \ell_n(\gamma).$$




# Negative binomial experiment

Example: Negative binomial models with an unknown parameter  $\gamma$ .

For  $n = 1, \dots, N$  let  $x_n | \gamma \stackrel{iid}{\sim} \text{NegativeBinomial}(\alpha, \gamma)$  for fixed  $\alpha$ .

$$\text{Write } \log p(X | \lambda, \gamma, w) = \sum_{n=1}^N w_n \ell_n(\gamma).$$

Negative Binomial model  
leaving out single datapoints with  $N = 800$




# Data Analysis Using Regression and Multilevel/Hierarchical Models.

We ran `rstanarm` on 56 different models on 13 different datasets from Gelman and Hill [2006], including Gaussian and logistic regression, fixed and mixed-effects models.

Across all models, we estimate 799 distinct covariances (regression coefficients and log scale parameters).

Using the bootstrap as ground truth, compute the relative errors:

$$\frac{V_{\text{Bayes}} - V_{\text{Boot}}}{|V_{\text{Boot}}|} \quad \text{and} \quad \frac{V_{\text{IJ}} - V_{\text{Boot}}}{|V_{\text{Boot}}|}.$$



**Figure 1:** The distribution of the relative errors. Log scale parameters include all variances or covariances that involve at least one log scale parameters.



**Total compute time for all models:**

Initial fit: 1.6 hours

Bootstrap: 381.5 hours

# How to connect to the election data?

Problem: MCMC is only interesting when the posterior doesn't concentrate.



Example: Exponential families with random effects (REs)  $\lambda$  and fixed effects  $\gamma$ .

If the observations per random effect remains bounded as  $N \rightarrow \infty$ , then

- Parameter  $\lambda$  (“local”) grows in dimension with  $N$ .
- Parameter  $\gamma$  (“global”) is finite-dimensional.
- Marginally  $p(\lambda|X)$  does not concentrate.
- Marginally,  $p(\gamma|X)$  concentrates.

# High dimensional problems

Example: Exponential families with random effects (REs)  $\lambda$  and fixed effects  $\gamma$ .

If the observations per random effect remains bounded as  $N \rightarrow \infty$ , then

- Parameter  $\lambda$  (“local”) grows in dimension with  $N$ .
- Parameter  $\gamma$  (“global”) is finite-dimensional.
- Marginally  $p(\lambda|X)$  does not concentrate.
- Marginally,  $p(\gamma|X)$  concentrates.

In general, we cannot hope for an asymptotic analysis of  $\mathbb{E}_{p(\lambda, \gamma|X)} [f(\lambda)]$ .

Can we save the approximation when *some* parameters concentrate?

Does the residual vanish asymptotically for  $w_n \mapsto \mathbb{E}_{p(\gamma|X, w_n)} [f(\gamma)]$ ?

## High dimensional problems

We assume that  $p(\gamma|X)$  concentrates but  $p(\lambda|X)$  does not. By our series expansion:

$$\begin{aligned} \mathbb{E}_{p(\gamma, \lambda|X, w_n)}[\gamma] - \mathbb{E}_{p(\gamma, \lambda|X)}[\gamma] = \\ \psi_n(w_n - 1) + \mathcal{E}(w_n) \end{aligned}$$

## High dimensional problems

We assume that  $p(\gamma|X)$  concentrates but  $p(\lambda|X)$  does not. By our series expansion:

$$\begin{aligned} & \mathbb{E}_{p(\gamma, \lambda|X, w_n)} [\gamma] - \mathbb{E}_{p(\gamma, \lambda|X)} [\gamma] = \\ & \quad \psi_n(w_n - 1) + \mathcal{E}(w_n) \\ = & \mathbb{E}_{p(\gamma, \lambda|X)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)] (w_n - 1) + \frac{1}{2} \mathbb{E}_{p(\gamma, \lambda|X, \tilde{w}_n)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)^2] (w_n - 1)^2 \end{aligned}$$

# High dimensional problems

We assume that  $p(\gamma|X)$  concentrates but  $p(\lambda|X)$  does not. By our series expansion:

$$\begin{aligned}
 & \mathbb{E}_{p(\gamma, \lambda|X, w_n)} [\gamma] - \mathbb{E}_{p(\gamma, \lambda|X)} [\gamma] = \\
 & \quad \psi_n(w_n - 1) + \mathcal{E}(w_n) \\
 = & \mathbb{E}_{p(\gamma, \lambda|X)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)] (w_n - 1) + \frac{1}{2} \mathbb{E}_{p(\gamma, \lambda|X, \tilde{w}_n)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)^2] (w_n - 1)^2 \\
 = & \mathbb{E}_{p(\gamma|X)} \left[ \bar{\gamma} \underbrace{\mathbb{E}_{p(\lambda|\gamma, X)} [\bar{\ell}_n(\gamma, \lambda)]}_{F_1(\gamma)} \right] (w_n - 1) + \frac{1}{2} \mathbb{E}_{p(\gamma|X, \tilde{w}_n)} \left[ \bar{\gamma} \underbrace{\mathbb{E}_{p(\lambda|X, \gamma, \tilde{w}_n)} [\bar{\ell}_n(\gamma, \lambda)^2]}_{F_2(\gamma)} \right] (w_n - 1)
 \end{aligned}$$



# High dimensional problems

We assume that  $p(\gamma|X)$  concentrates but  $p(\lambda|X)$  does not. By our series expansion:

$$\begin{aligned}
 & p(\gamma, \lambda|X, w_n) \mathbb{E}[\gamma] - p(\gamma, \lambda|X) \mathbb{E}[\gamma] = \\
 & \quad \psi_n(w_n - 1) + \mathcal{E}(w_n) \\
 = & \quad \mathbb{E}_{p(\gamma, \lambda|X)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)] (w_n - 1) + \frac{1}{2} \mathbb{E}_{p(\gamma, \lambda|X, \tilde{w}_n)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)^2] (w_n - 1)^2 \\
 = & \quad \mathbb{E}_{p(\gamma|X)} \left[ \bar{\gamma} \underbrace{\mathbb{E}_{p(\lambda|\gamma, X)} [\bar{\ell}_n(\gamma, \lambda)]}_{F_1(\gamma)} \right] (w_n - 1) + \frac{1}{2} \mathbb{E}_{p(\gamma|X, \tilde{w}_n)} \left[ \bar{\gamma} \underbrace{\mathbb{E}_{p(\lambda|X, \gamma, \tilde{w}_n)} [\bar{\ell}_n(\gamma, \lambda)^2]}_{F_2(\gamma)} \right] (w_n - 1)^2 \\
 = & \quad \underbrace{\mathbb{E}_{p(\gamma|X)} [\bar{\gamma} F_1(\gamma)]}_{O_p(N^{-1})} (w_n - 1) + \frac{1}{2} \underbrace{\mathbb{E}_{p(\gamma|X, \tilde{w}_n)} [\bar{\gamma} F_2(\gamma)]}_{O_p(N^{-1})} (w_n - 1)^2 \\
 & \quad \text{(by } p(\gamma|X) \text{ concentration)} \quad \text{(by } p(\gamma|X) \text{ concentration)} \\
 \Rightarrow &
 \end{aligned}$$

# High dimensional problems

We assume that  $p(\gamma|X)$  concentrates but  $p(\lambda|X)$  does not. By our series expansion:

$$\begin{aligned}
 & p(\gamma, \lambda|X, w_n) [\gamma] - p(\gamma, \lambda|X) [\gamma] = \\
 & \quad \psi_n(w_n - 1) + \mathcal{E}(w_n) \\
 = & \underbrace{p(\gamma, \lambda|X) [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)]}_{F_1(\gamma)} (w_n - 1) + \frac{1}{2} \underbrace{p(\gamma, \lambda|X, \tilde{w}_n) [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)^2]}_{F_2(\gamma)} (w_n - 1)^2 \\
 = & \underbrace{p(\gamma|X) [\bar{\gamma} \underbrace{p(\lambda|\gamma, X) [\bar{\ell}_n(\gamma, \lambda)]}_{F_1(\gamma)}]}_{O_p(N^{-1})} (w_n - 1) + \frac{1}{2} \underbrace{p(\gamma|X, \tilde{w}_n) [\bar{\gamma} \underbrace{p(\lambda|X, \gamma, \tilde{w}_n) [\bar{\ell}_n(\gamma, \lambda)^2]}_{F_2(\gamma)}]}_{O_p(N^{-1})} (w_n - 1)^2 \\
 = & \underbrace{p(\gamma|X) [\bar{\gamma} F_1(\gamma)]}_{O_p(N^{-1})} (w_n - 1) + \frac{1}{2} \underbrace{p(\gamma|X, \tilde{w}_n) [\bar{\gamma} F_2(\gamma)]}_{O_p(N^{-1})} (w_n - 1)^2 \\
 & \text{(by } p(\gamma|X) \text{ concentration)} \quad \text{(by } p(\gamma|X) \text{ concentration)} \\
 \Rightarrow & \psi_n = O_p(N^{-1}) \quad \mathcal{E}(w_n) = O_p(N^{-1})
 \end{aligned}$$

# High dimensional problems

We assume that  $p(\gamma|X)$  concentrates but  $p(\lambda|X)$  does not. By our series expansion:

$$\begin{aligned}
 & p(\gamma, \lambda|X, w_n) [\gamma] - p(\gamma, \lambda|X) [\gamma] = \\
 & \quad \psi_n(w_n - 1) + \mathcal{E}(w_n) \\
 = & \quad \mathbb{E}_{p(\gamma, \lambda|X)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)] (w_n - 1) + \frac{1}{2} \mathbb{E}_{p(\gamma, \lambda|X, \tilde{w}_n)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)^2] (w_n - 1)^2 \\
 = & \quad \mathbb{E}_{p(\gamma|X)} \left[ \bar{\gamma} \underbrace{\mathbb{E}_{p(\lambda|\gamma, X)} [\bar{\ell}_n(\gamma, \lambda)]}_{F_1(\gamma)} \right] (w_n - 1) + \frac{1}{2} \mathbb{E}_{p(\gamma|X, \tilde{w}_n)} \left[ \bar{\gamma} \underbrace{\mathbb{E}_{p(\lambda|X, \gamma, \tilde{w}_n)} [\bar{\ell}_n(\gamma, \lambda)^2]}_{F_2(\gamma)} \right] (w_n - 1) \\
 = & \quad \underbrace{\mathbb{E}_{p(\gamma|X)} [\bar{\gamma} F_1(\gamma)]}_{O_p(N^{-1})} (w_n - 1) + \frac{1}{2} \underbrace{\mathbb{E}_{p(\gamma|X, \tilde{w}_n)} [\bar{\gamma} F_2(\gamma)]}_{O_p(N^{-1})} (w_n - 1)^2 \\
 & \quad \text{(by } p(\gamma|X) \text{ concentration)} \quad \text{(by } p(\gamma|X) \text{ concentration)} \\
 \Rightarrow & \quad \psi_n = O_p(N^{-1}) \quad \mathcal{E}(w_n) = O_p(N^{-1})
 \end{aligned}$$

## Corollary [Giordano and Broderick, 2023]:

In general,  $w_n \mapsto N \left( \mathbb{E}_{p(\gamma|X, w_n)} [\gamma] - \mathbb{E}_{p(\gamma|X)} [\gamma] \right)$  remains non-linear as  $N \rightarrow \infty$ .

How can we apply the single-weight result to variance computations?

# Bayesian von-Mises Expansion

How can we apply the single-weight result to variance computations?

Define the “generalized posterior” functional

$$T(\mathbb{G}, N) := \frac{\int g(\theta) \exp \left( N \int \ell(x_0|\theta) \mathbb{G}(dx_0) \right) \pi(\theta) d\theta}{\int \exp \left( N \int \ell(x_0|\theta) \mathbb{G}(dx_0) \right) \pi(\theta) d\theta}.$$

Let  $\mathbb{F}_N$  denote the empirical distribution. Then

$$\mathbb{E}_{p(\theta|X)}[g(\theta)] = \frac{\int g(\theta) \exp \left( N \frac{1}{N} \sum_{n=1}^N \ell(x_n|\theta) \right) \pi(\theta) d\theta}{\int \exp \left( N \frac{1}{N} \sum_{n=1}^N \ell(x_n|\theta) \right) \pi(\theta) d\theta} = T(\mathbb{F}_N, N).$$

# Bayesian von-Mises Expansion

How can we apply the single-weight result to variance computations?

Define the “generalized posterior” functional

$$T(\mathbb{G}, N) := \frac{\int g(\theta) \exp \left( N \int \ell(x_0|\theta) \mathbb{G}(dx_0) \right) \pi(\theta) d\theta}{\int \exp \left( N \int \ell(x_0|\theta) \mathbb{G}(dx_0) \right) \pi(\theta) d\theta}.$$

Let  $\mathbb{F}_N$  denote the empirical distribution. Then

$$\mathbb{E}_{p(\theta|X)}[g(\theta)] = \frac{\int g(\theta) \exp \left( N \frac{1}{N} \sum_{n=1}^N \ell(x_n|\theta) \right) \pi(\theta) d\theta}{\int \exp \left( N \frac{1}{N} \sum_{n=1}^N \ell(x_n|\theta) \right) \pi(\theta) d\theta} = T(\mathbb{F}_N, N).$$

---

Let  $\mathbb{F}$  denote the true distribution of  $x_n$ , and let  $\mathbb{F}_N^t = t\mathbb{F}_N + (1-t)\mathbb{F}$ .

We can study the *von Mises expansion*:

$$\begin{aligned} \sqrt{N} \left( \mathbb{E}_{p(\theta|X)}[g(\theta)] - T(\mathbb{F}, N) \right) &= \sqrt{N} \left. \frac{\partial T(\mathbb{F}_N^t, N)}{\partial t} \right|_{t=0} (\mathbb{F}_N - \mathbb{F}) + \mathcal{E}(\tilde{t}) \\ &= \underbrace{\sqrt{N} \sum_{n=1}^N (\psi_n - \bar{\psi})}_{\text{Infinitesimal jackknife estimator}} + o_p(1) + \mathcal{E}(\tilde{t}). \end{aligned}$$

Inconsistency is suggested if  $\mathcal{E}(\tilde{t})$  fails to vanish.

**Theorem 3 [Giordano and Broderick, 2023] (sketch):**

**(Consistency of the von-Mises expansion in finite dimensions)**

Under slightly stronger conditions our original finite-dimensional posterior consistency result,

$$\sup_{\tilde{t} \in [0,1]} |\mathcal{E}(\tilde{t})| \rightarrow 0 \quad \text{in the Bayesian von-Mises expansion.}$$

# Bayesian von-Mises Expansion Results

**Theorem 3 [Giordano and Broderick, 2023] (sketch):**

**(Consistency of the von-Mises expansion in finite dimensions)**

Under slightly stronger conditions our original finite-dimensional posterior consistency result,

$$\sup_{\tilde{t} \in [0,1]} |\mathcal{E}(\tilde{t})| \rightarrow 0 \quad \text{in the Bayesian von-Mises expansion.}$$

**Theorem 4 [Giordano and Broderick, 2023] (sketch, not yet on arxiv):**

**(Inconsistency of the von-Mises expansion in infinite dimensions)**

Assume that  $x_n$  comes with a random group assignment  $g_n \in 1, \dots, G$ . Conditional on  $g, x_n$  is modeled as a finite-dimensional exponential family given  $\lambda, \gamma$ :

$$\log p(x_n | g_n = g, \gamma, \lambda) = \tau(x_n)^\top \eta_g(\gamma, \lambda) + \text{Constant.}$$

Define the average product of second moments:

$$\mathcal{V}_{\mathcal{N}}(\gamma) := \frac{1}{G} \sum_{g=1}^G \text{tr} \left( \mathbb{E}_{\mathbb{P}(x_n)} [\tau(x_n) \tau(x_n)^\top] \text{Cov}_{p(\lambda|\gamma, \mathbb{F})}(\eta_g(\gamma, \lambda)) \right).$$

If  $N \mathbb{E}_{p(\gamma|\mathbb{F})} [\bar{f}(\gamma) \mathcal{V}_{\mathcal{N}}(\gamma)]$  is strictly bounded away from 0 as  $N \rightarrow \infty$ , then


$$\sup_{\tilde{t} \in [0,1]} |\mathcal{E}(\tilde{t})| \rightarrow \infty \quad \text{in the Bayesian von-Mises expansion.}$$



# More experimental results for Gamma–Poisson mixtures

We ran simulations of the Gamma–Poisson mixture with different ratios of  $N/G$  (average observations per group).

- When  $N/G$  is small:
  - IJ is biased significantly downwards
  - Bootstrap is biased somewhat downwards
- When  $N/G$  is larger:
  - Both improve
  - Both remain somewhat biased
  - The IJ and bootstrap perform similarly



**Figure 2:** The error of the IJ and bootstrap covariances for different values of  $N$  and  $G$ . The y-axis shows the difference between  $N(V - \hat{V}_{\text{sim}})$ , where  $V$  is either  $\hat{V}_{\text{IJ}}$  or  $\hat{V}_{\text{Boot}}$ .

## Example: Poisson regression with Gamma-distributed random effects

For  $g = 1, \dots, G$ ,  $\lambda_g \stackrel{iid}{\sim} \text{Gamma}(\alpha, \beta)$  for fixed  $\alpha, \beta$

For  $n = 1, \dots, N$ ,  $g_n \stackrel{iid}{\sim} \text{Categorical}(1, \dots, G)$ ,  $y_n | \lambda_n, \gamma, g_n \stackrel{iid}{\sim} \text{Poisson}(\gamma \lambda_{g_n})$ .

$x_n = (y_n, g_n)$  are IID given  $\lambda, \gamma$ . Write  $\log p(X | \lambda, \gamma, w) = \sum_{n=1}^N w_n \ell_n(\lambda, \gamma)$ .

# Experiments


## Example: Poisson regression with Gamma-distributed random effects

For  $g = 1, \dots, G$ ,  $\lambda_g \stackrel{iid}{\sim} \text{Gamma}(\alpha, \beta)$  for fixed  $\alpha, \beta$

For  $n = 1, \dots, N$ ,  $g_n \stackrel{iid}{\sim} \text{Categorical}(1, \dots, G)$ ,  $y_n | \lambda_n, \gamma, g_n \stackrel{iid}{\sim} \text{Poisson}(\gamma \lambda_{g_n})$ .

$x_n = (y_n, g_n)$  are IID given  $\lambda, \gamma$ . Write  $\log p(X | \lambda, \gamma, w) = \sum_{n=1}^N w_n \ell_n(\lambda, \gamma)$ .

Poisson random effect model  
leaving out single datapoints with  $N = 800$



## Exchangeable units. (A contradiction?)

Negative binomial observations.

Asymptotically linear in  $w$ .

Poisson observations with random effects.

Asymptotically non-linear in  $w$ .

## Exchangeable units. (A contradiction?)

**Negative binomial observations.**

**Poisson observations with random effects.**

**Asymptotically linear in  $w$ .**

**Asymptotically non-linear in  $w$ .**

With a constant regressor, Gamma REs, and one RE per observation,  
these are the same model, with the same  $p(\gamma|X)$ .

**Is  $\mathbb{E}_{p(\gamma|X,w)} [\gamma]$  linear in the data weights or not?**

## Exchangeable units. (A contradiction?)

Negative binomial observations.

Poisson observations with random effects.

Asymptotically linear in  $w$ .

Asymptotically non-linear in  $w$ .

With a constant regressor, Gamma REs, and one RE per observation, these are the same model, with the same  $p(\gamma|X)$ .

Is  $\mathbb{E}_{p(\gamma|X,w)} [\gamma]$  linear in the **data weights** or not?

## Exchangeable units. (A contradiction?)

Negative binomial observations.

Poisson observations with random effects.

Asymptotically linear in  $w$ .

Asymptotically non-linear in  $w$ .

With a constant regressor, Gamma REs, and one RE per observation, these are the same model, with the same  $p(\gamma|X)$ .

Is  $\mathbb{E}_{p(\gamma|X,w)}[\gamma]$  linear in the **data weights** or not?

**Trick question!** We weight a log likelihood contribution, not a datapoint.

$$\log p(X|\gamma, w^m) = \sum_{n=1}^N w_n^m \log p(x_n|\gamma) \quad \log p(X|\gamma, \lambda, w^c) = \sum_{n=1}^N w_n^c \log p(x_n|\lambda, \gamma)$$

The two weightings are not equivalent in general.

What is the right exchangeable unit for a particular problem?


# Exchangeable units: Experimental results revisited

Our results were actually computed on **identical datasets** with  $G = N$  and  $g_n = n$ .

Uses  $\log p(x_n | \gamma)$ :

$$\psi_n = \mathbb{E}_{p(\gamma|X)} [\bar{\gamma} \bar{\ell}_n(\gamma)]$$


Negative Binomial model  
leaving out single datapoints with  $N = 800$



Uses  $\log p(x_n | \gamma, \lambda)$ :

$$\psi_n = \mathbb{E}_{p(\gamma, \lambda|X)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)]$$

Poisson random effect model  
leaving out single datapoints with  $N = 800$





# Exchangeable units: Experimental results revisited

Our results were actually computed on **identical datasets** with  $G = N$  and  $g_n = n$ .

Uses  $\log p(x_n | \gamma)$ :

$$\psi_n = \mathbb{E}_{p(\gamma|X)} [\bar{\gamma} \bar{\ell}_n(\gamma)]$$


Not easily computable from  
 $\gamma, \lambda \sim p(\gamma, \lambda | X)$   
in general.

Uses  $\log p(x_n | \gamma, \lambda)$ :


$$\psi_n = \mathbb{E}_{p(\gamma, \lambda | X)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)]$$

Easily computable from  
 $\gamma, \lambda \sim p(\gamma, \lambda | X)$ .

Negative Binomial model  
leaving out single datapoints with  $N = 800$



Poisson random effect model  
leaving out single datapoints with  $N = 800$



# Exchangeable units: Experimental results revisited

Our results were actually computed on **identical datasets** with  $G = N$  and  $g_n = n$ .

Uses  $\log p(x_n | \gamma)$ :

$$\psi_n = \mathbb{E}_{p(\gamma|X)} [\bar{\gamma} \bar{\ell}_n(\gamma)]$$

Not easily computable from  
 $\gamma, \lambda \sim p(\gamma, \lambda|X)$   
in general.


Uses  $\log p(x_n | \gamma, \lambda)$ :

$$\psi_n = \mathbb{E}_{p(\gamma, \lambda|X)} [\bar{\gamma} \bar{\ell}_n(\gamma, \lambda)]$$


Easily computable from  
 $\gamma, \lambda \sim p(\gamma, \lambda|X)$ .

May still be useful when  $p(\lambda|X)$   
is *somewhat* concentrated.

Negative Binomial model  
leaving out single datapoints with  $N = 800$



Poisson random effect model  
leaving out single datapoints with  $N = 800$



# Observations and consequences

- For finite-dimensional models which concentrate asymptotically:
  - Posterior expectations are approximately linear in data weights
  - The linearized variance estimate (infinitesimal jackknife) is consistent
  - The residual of the von Mises expansion vanishes
- For high-dimensional models which marginally concentrate only asymptotically:
  - Posterior expectations are not approximately linear in data weights
  - The linearized variance estimate (infinitesimal jackknife) is inconsistent
  - The residual of the von Mises expansion does not vanish
  - Even if the error  $\mathcal{E}(w)$  does not vanish, it can still be small enough in practice.  
... Especially given the linear approximation's huge computational advantage.

# Observations and consequences

- For finite-dimensional models which concentrate asymptotically:
  - Posterior expectations are approximately linear in data weights
  - The linearized variance estimate (infinitesimal jackknife) is consistent
  - The residual of the von Mises expansion vanishes
- For high-dimensional models which marginally concentrate only asymptotically:
  - Posterior expectations are not approximately linear in data weights
  - The linearized variance estimate (infinitesimal jackknife) is inconsistent
  - The residual of the von Mises expansion does not vanish
  - Even if the error  $\mathcal{E}(w)$  does not vanish, it can still be small enough in practice.  
... Especially given the linear approximation's huge computational advantage.
- When the weighting is linear, there are many other applications:
  - Cross-validation
  - Conformal inference
  - Identification of influential subsets
- When the weighting is non-linear, the inconsistency results should apply more widely:
  - The EM algorithm
  - The nonparametric bootstrap
  - Local prior sensitivity measures

# Observations and consequences

- For finite-dimensional models which concentrate asymptotically:
  - Posterior expectations are approximately linear in data weights
  - The linearized variance estimate (infinitesimal jackknife) is consistent
  - The residual of the von Mises expansion vanishes
- For high-dimensional models which marginally concentrate only asymptotically:
  - Posterior expectations are not approximately linear in data weights
  - The linearized variance estimate (infinitesimal jackknife) is inconsistent
  - The residual of the von Mises expansion does not vanish
  - Even if the error  $\mathcal{E}(w)$  does not vanish, it can still be small enough in practice.  
... Especially given the linear approximation's huge computational advantage.
- When the weighting is linear, there are many other applications:
  - Cross-validation
  - Conformal inference
  - Identification of influential subsets
- When the weighting is non-linear, the inconsistency results should apply more widely:
  - The EM algorithm
  - The nonparametric bootstrap
  - Local prior sensitivity measures

**Preprint:** Giordano and Broderick [2023] ([arXiv:2305.06466](https://arxiv.org/abs/2305.06466))

(Major update in progress, coming soon.)

- A. Gelman and M. Heidemanns. The Economist: Forecasting the US elections., 2020. URL <https://projects.economist.com/us-2020-forecast/president>. Data and model accessed Oct., 2020.
- A. Gelman and J. Hill. *Data analysis using regression and multilevel/hierarchical models*. Cambridge university press, 2006.
- R. Giordano and T. Broderick. The Bayesian infinitesimal jackknife for variance. *arXiv preprint arXiv:2305.06466*, 2023.
- J. Huggins and J. Miller. Reproducible model selection using bagged posteriors. *Bayesian Analysis*, 18(1):79–104, 2023.
- R. Kass, L. Tierney, and J. Kadane. The validity of posterior expansions based on Laplace’s method. *Bayesian and Likelihood Methods in Statistics and Econometrics*, 1990.
- B. Kleijn and A. Van der Vaart. The Bernstein-von-Mises theorem under misspecification. *Electronic Journal of Statistics*, 6: 354–381, 2012.