

Variational Methods for Latent Variable Problems

Ryan Giordano (for Johns Hopkins Biostats BLAST working group)

Oct, 2021

Massachusetts Institute of Technology

Outline for today:

- Some examples of latent variable models
- A template: The Neyman-Scott “paradox” and marginalization
- Bayesian versus frequentist approaches to marginalization
- The classical EM algorithm (in brief)

Next week, we will build on these ideas to present more general variational inference.


KL divergence exercises

$$\text{KL} (q(\theta)||p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{All bivariate normals}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL} (q(\theta)||p(\theta))$?




KL divergence exercises

$$\text{KL}(q(\theta)||p(\theta)) = -\mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{All bivariate normals}\}$

What is $q^*(\theta) = \operatorname{argmin}_{q \in \mathcal{Q}} \text{KL}(q(\theta)||p(\theta))$?



Sufficiently expressive families recover the target distribution.


KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{Independent bivariate normals}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(q(\theta) || p(\theta))$?




KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{Independent bivariate normals}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(q(\theta) || p(\theta))$?



KL minimizers “fit inside” the second argument.


KL divergence exercises

$$\begin{aligned} \text{KL}(\mathbf{p}(\theta) \parallel \mathbf{q}(\theta)) = \\ - \mathbb{E}_{\mathbf{p}(\theta)} [\log \mathbf{q}(\theta)] + \mathbb{E}_{\mathbf{p}(\theta)} [\log \mathbf{p}(\theta)] \end{aligned}$$

$\mathbf{p}(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{Independent bivariate normals}\}$

What is $\mathbf{q}^*(\theta) = \underset{\mathbf{q} \in \mathcal{Q}}{\text{argmin}} \text{KL}(\mathbf{p}(\theta) \parallel \mathbf{q}(\theta))$?




KL divergence exercises

$$\begin{aligned} \text{KL}(p(\theta) || q(\theta)) = \\ - \mathbb{E}_{p(\theta)} [\log q(\theta)] + \mathbb{E}_{p(\theta)} [\log p(\theta)] \end{aligned}$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{Independent bivariate normals}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(p(\theta) || q(\theta))$?



KL minimizers “fit inside” the second argument.


KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$ = Correlated bivariate normal

$$\mathcal{Q} = \{\text{Bivariate normals}\}$$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \left(- \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \right)$?




KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{Bivariate normals}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \left(- \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \right)$?



Without the entropy, the KL minimizer concentrates on the maximum of $\log p(\theta)$.


KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$ = Correlated bivariate normal

$$\mathcal{Q} = \{\text{Bivariate normals}\}$$

What is $q^*(\theta) =$
 $\underset{q \in \mathcal{Q}}{\operatorname{argmin}} \left(- \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \right) ?$




KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{Bivariate normals}\}$

What is $q^*(\theta) =$
 $\underset{q \in \mathcal{Q}}{\operatorname{argmin}} \left(- \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \right)$?



Without $\log p(\theta)$, the KL minimizer is infinitely dispersed.


KL divergence exercises

$$\text{KL}(q(\theta) || p(\theta)) = -\mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{Point masses}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(q(\theta) || p(\theta))$?




KL divergence exercises

$$\begin{aligned} \text{KL}(q(\theta) || p(\theta)) = \\ - \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)] \end{aligned}$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{Point masses}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\operatorname{argmin}} \text{KL}(q(\theta) || p(\theta))$?



Without a common dominating measure, the KL divergence is undefined.

KL divergence exercises


$$\text{KL}(q(\theta) || p(\theta)) =$$

$$- \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{BVN with small, fixed variance}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\text{argmin}} \text{KL}(q(\theta) || p(\theta))$?



KL divergence exercises


$$\text{KL}(q(\theta) || p(\theta)) =$$

$$- \mathbb{E}_{q(\theta)} [\log p(\theta)] + \mathbb{E}_{q(\theta)} [\log q(\theta)]$$

$p(\theta)$ = Correlated bivariate normal

$\mathcal{Q} = \{\text{BVN with small, fixed variance}\}$

What is $q^*(\theta) = \underset{q \in \mathcal{Q}}{\operatorname{argmin}} \text{KL}(q(\theta) || p(\theta))$?



Sufficiently concentrated distributions with constant entropy act like a point mass at the maximum of $\log p(\theta)$.

