

## Black box variational Bayes

“Black box variational Bayes” (BBVI) is a set of techniques for quickly and *automatically* approximating Bayesian posteriors using optimization. I’ll consider “mean field automatic differentiation variational inference” (ADVI).

# Black box variational Bayes

“Black box variational Bayes” (BBVI) is a set of techniques for quickly and *automatically* approximating Bayesian posteriors using optimization. I’ll consider “mean field automatic differentiation variational inference” (ADVI).

Let  $\theta$  denote model parameters and  $y$  some data and the joint generating distribution be  $\mathbb{P}(\theta, y) = \mathbb{P}(y|\theta) \mathbb{P}(\theta)$ . Let  $\mathbb{Q}(\theta|\eta)$  be a family of candidate approximate posteriors, here taken to be independent normals.

ADVI aims to find

$$\eta^* := \operatorname{argmin}_{\eta} \operatorname{KL}(\mathbb{Q}(\theta|\eta) || \mathbb{P}(\theta|y)) = \operatorname{argmin}_{\eta} \mathbb{E}_{\mathcal{N}(z)} [f(z|\eta)]$$

for a cleverly constructed, automatically-differentiable  $\eta \mapsto f(z|\eta)$ .

# Black box variational Bayes

“Black box variational Bayes” (BBVI) is a set of techniques for quickly and *automatically* approximating Bayesian posteriors using optimization. I’ll consider “mean field automatic differentiation variational inference” (ADVI).

Let  $\theta$  denote model parameters and  $y$  some data and the joint generating distribution be  $\mathbb{P}(\theta, y) = \mathbb{P}(y|\theta)\mathbb{P}(\theta)$ . Let  $\mathbb{Q}(\theta|\eta)$  be a family of candidate approximate posteriors, here taken to be independent normals.

ADVI aims to find

$$\eta^* := \underset{\eta}{\operatorname{argmin}} \operatorname{KL}(\mathbb{Q}(\theta|\eta) || \mathbb{P}(\theta|y)) = \underset{\eta}{\operatorname{argmin}} \mathbb{E}_{\mathcal{N}(z)} [f(z|\eta)]$$

for a cleverly constructed, automatically-differentiable  $\eta \mapsto f(z|\eta)$ .

Unfortunately,  $\mathbb{E}_{\mathcal{N}(z)} [f(z|\eta)]$  is typically intractable. So ADVI uses stochastic gradient (SG). This leads to the following problems:

- You have to tune the step size carefully
- You can’t assess convergence directly
- You can’t compute sensitivity, so you can’t use linear response covariances.

⇒ Optimization is slow and imprecise, and the posterior uncertainty is no good. Not so black box actually!

**We propose a simple alternative to SG that resolves these problems (sometimes).**

## Optimization of intractable expectations

Suppose you want to minimize an objective function of the form

$$\eta^* := \operatorname{argmin}_{\eta} \mathbb{E}_{\mathbb{P}(z)} [f(z|\eta)] := \operatorname{argmin}_{\eta} \ell(\eta),$$

where  $\mathbb{P}(z)$  is known, but the expectation is not available in closed form.

# Optimization of intractable expectations

Suppose you want to minimize an objective function of the form

$$\eta^* := \operatorname{argmin}_{\eta} \mathbb{E}_{\mathbb{P}(z)} [f(z|\eta)] := \operatorname{argmin}_{\eta} \ell(\eta),$$

where  $\mathbb{P}(z)$  is known, but the expectation is not available in closed form.

When does this happen?

- Black box variational inference
- Stochastic control (e.g. you have a factory, and supply and demand are random)

# Optimization of intractable expectations

Suppose you want to minimize an objective function of the form

$$\eta^* := \operatorname{argmin}_{\eta} \mathbb{E}_{\mathbb{P}(z)} [f(z|\eta)] := \operatorname{argmin}_{\eta} \ell(\eta),$$

where  $\mathbb{P}(z)$  is known, but the expectation is not available in closed form.

When does this happen?

- Black box variational inference
- Stochastic control (e.g. you have a factory, and supply and demand are random)

What can you do? There are two options, both using the Monte Carlo (MC) estimate

$$\hat{\ell}(\eta) := \frac{1}{N} \sum_{n=1}^N f(z_n|\eta) \approx \ell(\eta).$$

# Optimization of intractable expectations

Suppose you want to minimize an objective function of the form

$$\eta^* := \operatorname{argmin}_{\eta} \mathbb{E}_{\mathbb{P}(z)} [f(z|\eta)] := \operatorname{argmin}_{\eta} \ell(\eta),$$

where  $\mathbb{P}(z)$  is known, but the expectation is not available in closed form.

When does this happen?

- Black box variational inference
- Stochastic control (e.g. you have a factory, and supply and demand are random)

What can you do? There are two options, both using the Monte Carlo (MC) estimate

$$\hat{\ell}(\eta) := \frac{1}{N} \sum_{n=1}^N f(z_n|\eta) \approx \ell(\eta).$$

- Stochastic gradient (SG)
  - Update with  $\eta^i = \eta^{i-1} - \rho \nabla_{\eta} \hat{\ell}(\eta)$  for some step size  $\rho$  (new  $z_n$  every step)
  - Approximately minimizes the exact objective

# Optimization of intractable expectations

Suppose you want to minimize an objective function of the form

$$\eta^* := \operatorname{argmin}_{\eta} \mathbb{E}_{\mathbb{P}(z)} [f(z|\eta)] := \operatorname{argmin}_{\eta} \ell(\eta),$$

where  $\mathbb{P}(z)$  is known, but the expectation is not available in closed form.

When does this happen?

- Black box variational inference
- Stochastic control (e.g. you have a factory, and supply and demand are random)

What can you do? There are two options, both using the Monte Carlo (MC) estimate

$$\hat{\ell}(\eta) := \frac{1}{N} \sum_{n=1}^N f(z_n|\eta) \approx \ell(\eta).$$

- Stochastic gradient (SG)
  - Update with  $\eta^i = \eta^{i-1} - \rho \nabla_{\eta} \hat{\ell}(\eta)$  for some step size  $\rho$  (new  $z_n$  every step)
  - Approximately minimizes the exact objective
- Sample average approximation (SAA)
  - Find  $\hat{\eta} := \operatorname{argmin}_{\eta} \hat{\ell}(\eta)$  for fixed  $z_n$
  - Exactly minimizes approximate objective



# Optimization of intractable expectations

Suppose you want to minimize an objective function of the form

$$\eta^* := \operatorname{argmin}_{\eta} \mathbb{E}_{\mathbb{P}(z)} [f(z|\eta)] := \operatorname{argmin}_{\eta} \ell(\eta),$$

where  $\mathbb{P}(z)$  is known, but the expectation is not available in closed form.

When does this happen?

- Black box variational inference
- Stochastic control (e.g. you have a factory, and supply and demand are random)

What can you do? There are two options, both using the Monte Carlo (MC) estimate

$$\hat{\ell}(\eta) := \frac{1}{N} \sum_{n=1}^N f(z_n|\eta) \approx \ell(\eta).$$

- Stochastic gradient (SG)
  - Update with  $\eta^i = \eta^{i-1} - \rho \nabla_{\eta} \hat{\ell}(\eta)$  for some step size  $\rho$  (new  $z_n$  every step)
  - Approximately minimizes the exact objective
- Sample average approximation (SAA)
  - Find  $\hat{\eta} := \operatorname{argmin}_{\eta} \hat{\ell}(\eta)$  for fixed  $z_n$
  - Exactly minimizes approximate objective

Which is better? **In general, it depends.**

As far as we can tell, the BBVI literature has only ever considered SG.

# Optimization of intractable expectations

## Sample average approximation (SAA)

- Find  $\hat{\eta} := \operatorname{argmin}_{\eta} \hat{\ell}(\eta)$
- Fixed  $z_n$  for whole procedure
- Exactly minimizes approximate objective

## Stochastic gradient (SG)

- $\eta^i = \eta^{i-1} - \rho \nabla_{\eta} \hat{\ell}(\eta)$
- New  $z_n$  every step
- Approximately minimizes the exact objective

# Optimization of intractable expectations

## Sample average approximation (SAA)

- Find  $\hat{\eta} := \operatorname{argmin}_{\eta} \hat{\ell}(\eta)$
- Fixed  $z_n$  for whole procedure
- Exactly minimizes approximate objective

### Advantages:

- Can use fast off-the-shelf second-order optimization (great for poorly-conditioned problems)
- Can evaluate the objective function exactly to check for convergence
- Can compute sensitivity (linear response covariances  $\Rightarrow$  more accurate posterior covariances for mean field approximations)

## Stochastic gradient (SG)

- $\eta^i = \eta^{i-1} - \rho \nabla_{\eta} \hat{\ell}(\eta)$
- New  $z_n$  every step
- Approximately minimizes the exact objective

# Optimization of intractable expectations

## Sample average approximation (SAA)

- Find  $\hat{\eta} := \operatorname{argmin}_{\eta} \hat{\ell}(\eta)$
- Fixed  $z_n$  for whole procedure
- Exactly minimizes approximate objective

### Advantages:

- Can use fast off-the-shelf second-order optimization (great for poorly-conditioned problems)
- Can evaluate the objective function exactly to check for convergence
- Can compute sensitivity (linear response covariances  $\Rightarrow$  more accurate posterior covariances for mean field approximations)

## Stochastic gradient (SG)

- $\eta^i = \eta^{i-1} - \rho \nabla_{\eta} \hat{\ell}(\eta)$
- New  $z_n$  every step
- Approximately minimizes the exact objective

### Advantages:

- Uses each draw  $z_n$  only once (for a single gradient step)

# Optimization of intractable expectations

## Sample average approximation (SAA)

- Find  $\hat{\eta} := \operatorname{argmin}_{\eta} \hat{\ell}(\eta)$
- Fixed  $z_n$  for whole procedure
- Exactly minimizes approximate objective

### Advantages:

- Can use fast off-the-shelf second-order optimization (great for poorly-conditioned problems)
- Can evaluate the objective function exactly to check for convergence
- Can compute sensitivity (linear response covariances  $\Rightarrow$  more accurate posterior covariances for mean field approximations)

## Stochastic gradient (SG)

- $\eta^i = \eta^{i-1} - \rho \nabla_{\eta} \hat{\ell}(\eta)$
- New  $z_n$  every step
- Approximately minimizes the exact objective

### Advantages:

- Uses each draw  $z_n$  only once (for a single gradient step)

This is actually a big one! Because if  $\eta \in \mathbb{R}^D$ , in general, both SG and SAA have accuracy  $(D/N)^{-1/2}$ , where  $N$  is the *total* number of draws of  $z_n$  used.

SAA uses each draw at each step of optimization. SG uses each draw once.  
 $\Rightarrow$  In general, SG is much more efficient in high dimensions!

# Optimization of intractable expectations

## Sample average approximation (SAA)

- Find  $\hat{\eta} := \operatorname{argmin}_{\eta} \hat{\ell}(\eta)$
- Fixed  $z_n$  for whole procedure
- Exactly minimizes approximate objective

### Advantages:

- Can use fast off-the-shelf second-order optimization (great for poorly-conditioned problems)
- Can evaluate the objective function exactly to check for convergence
- Can compute sensitivity (linear response covariances  $\Rightarrow$  more accurate posterior covariances for mean field approximations)

## Stochastic gradient (SG)

- $\eta^i = \eta^{i-1} - \rho \nabla_{\eta} \hat{\ell}(\eta)$
- New  $z_n$  every step
- Approximately minimizes the exact objective

### Advantages:

- Uses each draw  $z_n$  only once (for a single gradient step)


This is actually a big one! Because if  $\eta \in \mathbb{R}^D$ , in general, both SG and SAA have accuracy  $(D/N)^{-1/2}$ , where  $N$  is the *total* number of draws of  $z_n$  used.

SAA uses each draw at each step of optimization. SG uses each draw once.  
 $\Rightarrow$  In general, SG is much more efficient in high dimensions!


---

**Theorem (us).** If  $\log \mathbb{P}(\theta, y)$  is high dimensional due to a large number of “local” variables, then the accuracy is  $(\log D/N)^{-1/2}$ , rendering SAA feasible.

## Experimental results: Runtime



## Experimental results: Means





## Experimental results: Standard deviations

