

Locally Equivalent Weights for Bayesian MrP

Ryan Giordano, Alice Cima, Erin Hartman, Jared Murray, Avi Feller

UT Austin Statistics Seminar

September 2025



Standard error estimation

Does this mean anything?

Yes: We can meaningfully sum these weights against regressors.

What else might it mean?

Does the spread relate to frequentist variance?




Figure 1: Comparison between raking and MrPlew weights for the Name Change dataset




Figure 2: Comparison between raking and MrPlew weights for the Gay Marriage dataset

Standard error consistency theorem: (sketch)

For Bayesian hierarchical logistic regression, define

$$\varepsilon_n = y_n - \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} [m(\mathbf{x}_n^\top \theta)] \quad \text{and} \quad \psi_n := N_S w_n^{\text{MrP}} \varepsilon_n.$$

We state mild conditions under which, as $N \rightarrow \infty$,

$$\begin{aligned} \sqrt{N} (\hat{\mu}_{\text{MrP}} - \mu_\infty) &\rightarrow \mathcal{N}(0, V) \quad \text{for some } \mu_\infty \text{ and variance } V, \text{ and} \\ \frac{1}{N_S} \sum_{i=1}^{N_S} (\psi_n - \bar{\psi})^2 &\rightarrow V. \end{aligned}$$

The use of w_n^{MrP} is exactly analogous to the use of raking weights for standard error estimation. This builds on our earlier work on the Bayesian infinitesimal jackknife¹.

¹G. and Broderick 2024.

Standard error estimation




Figure 3: Frequentist standard deviation estimates

Future work

Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

But the high level idea can be extended much more widely:

1. Assume your initial model was accurate
2. Select some perturbation your model should be able to capture
3. Use local sensitivity to detect whether the change is what you expect
4. If the change is not what you expect, either (1) or (2) was wrong

Future work

Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

But the high level idea can be extended much more widely:

1. Assume your initial model was accurate
2. Select some perturbation your model should be able to capture
3. Use local sensitivity to detect whether the change is what you expect
4. If the change is not what you expect, either (1) or (2) was wrong

Such checks recover generalized versions of many standard diagnostics for linear models.

Examples:

- Additive parameter shifts \leftrightarrow Unbiasedness
- Invariance to survey data weighting \leftrightarrow Regressor + residual orthogonality
- Importance sampling \leftrightarrow Sandwich covariance $\stackrel{?}{=}$ Inverse Fisher information

Student contributions and ongoing work:

- **Alice Cima** contributed significantly to this work
- **Vladimir Palmin** is working on extending MrPlew to lme4
- **Sequoia Andrade** is working on generalizing to other local sensitivity checks
- **Lucas Schwengber** is working on novel flow-based techniques for local sensitivity
- **(Currently recruiting!)** Doubly-robust Bayesian Hierarchical MrP




Alice Cima



Vladimir Palmin



Sequoia Andrade



Lucas Schwengber

Preprint and R package (hopefully) coming soon!



G. and T. Broderick (2024). *The Bayesian Infinitesimal Jackknife for Variance*. arXiv: 2305.06466 [stat.ME]. URL: <https://arxiv.org/abs/2305.06466>.