

Locally Equivalent Weights for Bayesian MrP

Ryan Giordano, Alice Cima, Erin Hartman, Jared Murray, Avi Feller

University of British Columbia Statistics Seminar

October 2025




The basic problem


We have a survey population, for whom we observe:

- Covariates \mathbf{x} (e.g. race, gender, zip code, age, education level)
- Responses y (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe (\mathbf{x}_i, y_i) for $i = 1, \dots, N_S$



Observe \mathbf{x}_j for $j = 1, \dots, N_T$

¹Photo copyright: Mark Taylor / naturepl.com

The basic problem


We have a survey population, for whom we observe:

- Covariates \mathbf{x} (e.g. race, gender, zip code, age, education level)
- Responses y (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe (\mathbf{x}_i, y_i) for $i = 1, \dots, N_S$



Observe \mathbf{x}_j for $j = 1, \dots, N_T$

The problem is that the populations may be very different, maybe leading to bias. ¹

¹Photo copyright: Mark Taylor / naturepl.com

The basic problem


We have a survey population, for whom we observe:

- Covariates \mathbf{x} (e.g. race, gender, zip code, age, education level)
- Responses y (e.g. A binary response to “do you support Trump”)

We want the average response in a target population, in which we observe only covariates.



Observe (\mathbf{x}_i, y_i) for $i = 1, \dots, N_S$



Observe \mathbf{x}_j for $j = 1, \dots, N_T$

The problem is that the populations may be very different, maybe leading to bias. ¹

How can we use the covariates to say something about the target responses?

¹Photo copyright: Mark Taylor / naturepl.com

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target y_j . Let $Y_S = \{y_1, \dots, y_{N_S}\}$.

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target y_j . Let $Y_S = \{y_1, \dots, y_{N_S}\}$.

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- Choose “calibration weights” w_i
using only the regressors \mathbf{x}
(e.g. raking weights)

Bayesian hierarchical modeling (MrP)

- Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$,
choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$
(e.g. Hierarchical logistic regression)

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target y_j . Let $Y_S = \{y_1, \dots, y_{N_S}\}$.

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- Choose “calibration weights” w_i
using only the regressors \mathbf{x}
(e.g. raking weights)
- Take $\hat{\mu}^{\text{CW}}(Y_S) = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$

Bayesian hierarchical modeling (MrP)

- Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$,
choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$
(e.g. Hierarchical logistic regression)
- Take $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$ and
 $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target y_j . Let $Y_S = \{y_1, \dots, y_{N_S}\}$.

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- ▶ Choose “calibration weights” w_i using only the regressors \mathbf{x} (e.g. raking weights)
- ▶ Take $\hat{\mu}^{CW}(Y_S) = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$
- ▶ Dependence on y_i is clear

Bayesian hierarchical modeling (MrP)

- ▶ Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$, choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$ (e.g. Hierarchical logistic regression)
- ▶ Take $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$ and $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$
- ▶ Dependence on y_i very complicated (Typically via MCMC draws from $\mathcal{P}(\theta|\text{Survey data})$)

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target y_j . Let $Y_S = \{y_1, \dots, y_{N_S}\}$.

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- ▶ Choose “calibration weights” w_i using only the regressors \mathbf{x} (e.g. raking weights)
- ▶ Take $\hat{\mu}^{CW}(Y_S) = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$
 - ▶ Dependence on y_i is clear
- ▶ Weights give interpretable diagnostics:
 - Regressor balance
 - Frequentist variability
 - Partial pooling
 - Extrapolation

Bayesian hierarchical modeling (MrP)

- ▶ Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$, choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$ (e.g. Hierarchical logistic regression)
- ▶ Take $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$ and $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$
 - ▶ Dependence on y_i very complicated (Typically via MCMC draws from $\mathcal{P}(\theta|\text{Survey data})$)
- ▶ **Black box**

Two approaches

We want $\mu := \frac{1}{N_T} \sum_{j=1}^{N_T} y_j$, but don't observe target y_j . Let $Y_S = \{y_1, \dots, y_{N_S}\}$.

- Assume $p(y|\mathbf{x})$ is the same in both populations,
- But the distribution of \mathbf{x} may be different in the survey and target.

Calibration weighting (CW)

- ▶ Choose “calibration weights” w_i using only the regressors \mathbf{x} (e.g. raking weights)
- ▶ Take $\hat{\mu}^{CW}(Y_S) = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$
 - ▶ Dependence on y_i is clear
- ▶ Weights give interpretable diagnostics:
 - Regressor balance
 - Frequentist variability
 - Partial pooling
 - Extrapolation

Bayesian hierarchical modeling (MrP)

- ▶ Choose $\mathbb{E}[y|\mathbf{x}, \theta] = m(\theta^\top \mathbf{x})$, choose prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$ (e.g. Hierarchical logistic regression)
- ▶ Take $\hat{y}_j = \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})}[y|\mathbf{x}_j]$ and $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j$
 - ▶ Dependence on y_i very complicated (Typically via MCMC draws from $\mathcal{P}(\theta|\text{Survey data})$)
- ▶ **Black box**
 - ← We open this box, providing analogues of all these diagnostics

Prior work: Equivalent weights for linear models

Gelman (2007b) observes that MrP is a CW estimator when one uses linear regression to form \hat{y} :

$$\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j = \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\mathbf{x}_j^\top \hat{\boldsymbol{\theta}}}_{\text{Linear in } Y_S}$$

Most existing literature on comparing CW and MrP focus on such linear models. ²

²For example, Gelman (2007b), B., F., and H. (2021), and Chattopadhyay and Zubizarreta (2023).

Prior work: Equivalent weights for linear models

Gelman (2007b) observes that MrP is a CW estimator when one uses linear regression to form \hat{y} :

$$\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \hat{y}_j = \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\mathbf{x}_j^\top \hat{\boldsymbol{\theta}}}_{\text{Linear in } Y_S}$$

Most existing literature on comparing CW and MrP focus on such linear models. ²

But what if you use a non-linear link function? Or a hierarchical model?

“It would also be desirable to use nonlinear methods ... but then it would seem difficult to construct even approximately equivalent weights. Weighting and fully nonlinear models would seem to be completely incompatible methods.” — (Gelman 2007a)

²For example, Gelman (2007b), B., F., and H. (2021), and Chattopadhyay and Zubizarreta (2023).

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

The map from $Y_S \mapsto m(\mathbf{x}_j^\top \hat{\theta})$ is *inherently nonlinear*.

But *some sample averages* of $m(\mathbf{x}_j^\top \hat{\theta})$ can be approximately linear.

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is a *approximately* a CW estimator.

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is a *approximately* a CW estimator.

$$\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$$

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . **Then MrP is a *approximately* a CW estimator.**

$$\begin{aligned}\hat{\mu}^{\text{MrP}}(Y_S) &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} \quad (\text{Law of large numbers})\end{aligned}$$

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}^{\text{MrP}}(Y_S) &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)}\end{aligned}$$

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is a *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}^{\text{MrP}}(Y_S) &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)}\end{aligned}$$

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is a *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}^{\text{MrP}}(Y_S) &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)} \\ &\approx \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i m(\mathbf{x}_i^\top \hat{\theta}) && \text{(Law of large numbers)}\end{aligned}$$

Approximately equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is *approximately* a CW estimator.

$$\begin{aligned}\hat{\mu}^{\text{MrP}}(Y_S) &= \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) \\ &\approx \int m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_T(\mathbf{x}) d\mathbf{x} && \text{(Law of large numbers)} \\ &= \int \frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(Multiply by } \mathcal{P}_S(\mathbf{x})/\mathcal{P}_S(\mathbf{x}) \text{)} \\ &\approx \int (\alpha^\top \mathbf{x}) m(\mathbf{x}^\top \hat{\theta}) \mathcal{P}_S(\mathbf{x}) d\mathbf{x} && \text{(By assumption)} \\ &\approx \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i m(\mathbf{x}_i^\top \hat{\theta}) && \text{(Law of large numbers)} \\ &= \alpha^\top \frac{1}{N_S} \sum_{i=1}^{N_S} \mathbf{x}_i y_i && \text{(Property of exponential family MLEs)}\end{aligned}$$

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is *approximately* a CW estimator.

$$\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \underbrace{w_i^{\text{MrP}}}_{\alpha^\top \mathbf{x}_i} y_i + \text{Small error}$$

³Krantz and Parks 2012; G., Stephenson, et al. 2019.

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is *approximately* a CW estimator.

$$\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \underbrace{w_i^{\text{MrP}}}_{\alpha^\top \mathbf{x}_i} y_i + \text{Small error}$$

But what are the weights? We don't observe $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$, so can't estimate α directly.

³Krantz and Parks 2012; G., Stephenson, et al. 2019.

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is *approximately* a CW estimator.

$$\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \underbrace{w_i^{\text{MrP}}}_{\alpha^\top \mathbf{x}_i} y_i + \text{Small error}$$

But what are the weights? We don't observe $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$, so can't estimate α directly.

Key idea (informal)

If $\hat{\mu}^{\text{MrP}}(Y_S)$ is approximately linear, then $w_i^{\text{MrP}} \approx N_S \frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i}$.

³Krantz and Parks 2012; G., Stephenson, et al. 2019.

Nearly equivalent weights for (some) logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$, with MLE $\hat{\theta}$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta})$.

Example

Suppose $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})} \approx \alpha^\top \mathbf{x}$ for some α . Then MrP is *approximately* a CW estimator.

$$\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} m(\mathbf{x}_j^\top \hat{\theta}) = \frac{1}{N_S} \sum_{i=1}^{N_S} \underbrace{w_i^{\text{MrP}}}_{\alpha^\top \mathbf{x}_i} y_i + \text{Small error}$$

But what are the weights? We don't observe $\frac{\mathcal{P}_T(\mathbf{x})}{\mathcal{P}_S(\mathbf{x})}$, so can't estimate α directly.

Key idea (informal)

If $\hat{\mu}^{\text{MrP}}(Y_S)$ is approximately linear, then $w_i^{\text{MrP}} \approx N_S \frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i}$.

For logistic regression, compute and analyze $\frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i}$ using the implicit function theorem.³

³Krantz and Parks 2012; G., Stephenson, et al. 2019.

Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$.
- Set a hierarchical prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$, use MCMC to draw from $\mathcal{P}(\theta|\text{Survey data})$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[m(\mathbf{x}_j^\top \theta) \right]$.

No reason to think $Y_S \mapsto \hat{\mu}^{\text{MrP}}(Y_S)$ is even approximately **globally** linear.

⁴Diaconis and Freedman 1986; Gustafson 1996; Efron 2015; G., Broderick, and Jordan 2018.

Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$.
- Set a hierarchical prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$, use MCMC to draw from $\mathcal{P}(\theta|\text{Survey data})$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[m(\mathbf{x}_j^\top \theta) \right]$.

No reason to think $Y_S \mapsto \hat{\mu}^{\text{MrP}}(Y_S)$ is even approximately **globally** linear.

But can still compute and analyze $w_i^{\text{MrP}} := N_S \frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i}$ using Bayesian sensitivity analysis!⁴

MrP weights for MCMC

$$w_i^{\text{MrP}} := N_S \frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i} = N_S \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\text{Cov}_{\mathcal{P}(\theta|\text{Survey data})} \left(m(\mathbf{x}_j^\top \theta), \theta^\top \mathbf{x}_i \right)}_{\text{Can estimate without rerunning MCMC!}}$$

⁴Diaconis and Freedman 1986; Gustafson 1996; Efron 2015; G., Broderick, and Jordan 2018.

Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$.
- Set a hierarchical prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$, use MCMC to draw from $\mathcal{P}(\theta|\text{Survey data})$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[m(\mathbf{x}_j^\top \theta) \right]$.

No reason to think $Y_S \mapsto \hat{\mu}^{\text{MrP}}(Y_S)$ is even approximately **globally** linear.

But can still compute and analyze $w_i^{\text{MrP}} := N_S \frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i}$ using Bayesian sensitivity analysis!⁴

MrP weights for MCMC

$$w_i^{\text{MrP}} := N_S \frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i} = N_S \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\text{Cov}_{\mathcal{P}(\theta|\text{Survey data})} \left(m(\mathbf{x}_j^\top \theta), \theta^\top \mathbf{x}_i \right)}_{\text{Can estimate without rerunning MCMC!}}$$

What do these weights mean? There are now two distinct possibilities:

- “Locally implicit weights”
 - An estimator of $\mathcal{P}_T(\mathbf{x})/\mathcal{P}_S(\mathbf{x})$ (via Riesz regression applied to the Gateaux derivative)
- “Locally equivalent weights”
 - A characterization of $Y_S \mapsto \hat{\mu}^{\text{MrP}}(Y_S)$ for diagnostics and interpretation

⁴Diaconis and Freedman 1986; Gustafson 1996; Efron 2015; G., Broderick, and Jordan 2018.

Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$.
- Set a hierarchical prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$, use MCMC to draw from $\mathcal{P}(\theta|\text{Survey data})$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[m(\mathbf{x}_j^\top \theta) \right]$.

No reason to think $Y_S \mapsto \hat{\mu}^{\text{MrP}}(Y_S)$ is even approximately **globally** linear.

But can still compute and analyze $w_i^{\text{MrP}} := N_S \frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i}$ using Bayesian sensitivity analysis!⁴

MrP weights for MCMC

$$w_i^{\text{MrP}} := N_S \frac{\partial \hat{\mu}^{\text{MrP}}(Y_S)}{\partial y_i} = N_S \frac{1}{N_T} \sum_{j=1}^{N_T} \underbrace{\text{Cov}_{\mathcal{P}(\theta|\text{Survey data})} \left(m(\mathbf{x}_j^\top \theta), \theta^\top \mathbf{x}_i \right)}_{\text{Can estimate without rerunning MCMC!}}$$

What do these weights mean? There are now two distinct possibilities:

- “Locally implicit weights”
 - An estimator of $\mathcal{P}_T(\mathbf{x})/\mathcal{P}_S(\mathbf{x})$ (via Riesz regression applied to the Gateaux derivative)
- “Locally equivalent weights” \leftarrow **The present talk will focus on this interpretation**
 - A characterization of $Y_S \mapsto \hat{\mu}^{\text{MrP}}(Y_S)$ for diagnostics and interpretation

⁴Diaconis and Freedman 1986; Gustafson 1996; Efron 2015; G., Broderick, and Jordan 2018.

Locally equivalent weights for hierarchical logistic regression MrP

- Suppose the model is $m(\mathbf{x}^\top \theta) = \text{Logistic}(\mathbf{x}^\top \theta)$.
- Set a hierarchical prior $\mathcal{P}(\theta|\Sigma)\mathcal{P}(\Sigma)$, use MCMC to draw from $\mathcal{P}(\theta|\text{Survey data})$.
- MrP is $\hat{\mu}^{\text{MrP}}(Y_S) = \frac{1}{N_T} \sum_{j=1}^{N_T} \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} \left[m(\mathbf{x}_j^\top \theta) \right]$.

MrP locally equivalent weights (MrPlew)

For new data \tilde{Y}_S , form a **MrP locally equivalent weighting**:

$$\hat{\mu}^{\text{MrP}}(\tilde{Y}_S) \approx \hat{\mu}^{\text{MrP}}(Y_S) + \sum_{i=1}^{N_S} w_i^{\text{MrP}} (\tilde{y}_i - y_i)$$

Our task is to rigorously show that even such local weights can be meaningfully used diagnostically in the same ways we use global weights.

The weights can look very different!

Does this mean anything?




Figure 1: Comparison between raking and MrPlew weights for the Name Change dataset




Figure 2: Comparison between raking and MrPlew weights for the Gay Marriage dataset

The weights can look very different!

Does this mean anything?

Does the spread relate to frequentist variance?




Figure 1: Comparison between raking and MrPlew weights for the Name Change dataset




Figure 2: Comparison between raking and MrPlew weights for the Gay Marriage dataset

Let $\hat{\text{Var}}(\cdot)$ denote the sample variance.

Calibration weighting standard errors: (sketch) ⁵

Suppose we have $\hat{\mu}^{\text{CW}}(Y_S) = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$ and a consistent residual estimate ε_i .

Then $\hat{\text{Var}}(w_i \varepsilon_i) \approx \text{Var}(\sqrt{N_S} \hat{\mu}^{\text{CW}}(Y_S))$.

⁵E.g. , Deville, Särndal, and Sautory (1993) and Fuller (2011).

⁶G. and Broderick 2024.

Standard error estimation

Let $\hat{\text{Var}}(\cdot)$ denote the sample variance.

Calibration weighting standard errors: (sketch)⁵

Suppose we have $\hat{\mu}^{\text{CW}}(Y_S) = \frac{1}{N_S} \sum_{i=1}^{N_S} w_i y_i$ and a consistent residual estimate ε_i .

Then $\hat{\text{Var}}(w_i \varepsilon_i) \approx \text{Var}(\sqrt{N_S} \hat{\mu}^{\text{CW}}(Y_S))$.

Standard error consistency theorem: (sketch)

For Bayesian hierarchical logistic regression, define $\varepsilon_i = y_i - \mathbb{E}_{\mathcal{P}(\theta|\text{Survey data})} [m(\mathbf{x}_i^T \theta)]$.

We state mild conditions under which, as $N_S \rightarrow \infty$, for some μ_∞ and variance V ,

$$\sqrt{N_S} (\hat{\mu}^{\text{MRP}}(Y_S) - \mu_\infty) \rightarrow \mathcal{N}(0, V) \quad \text{and} \quad \hat{\text{Var}}(w_i^{\text{MRP}} \varepsilon_i) \rightarrow V.$$

The use of w_i^{MRP} is exactly analogous to the use of raking weights for standard error estimation.

This builds on our earlier work on the Bayesian infinitesimal jackknife.⁶

⁵E.g., Deville, Särndal, and Sautory (1993) and Fuller (2011).

⁶G. and Broderick 2024.

Standard error estimation




Figure 3: Frequentist standard deviation estimates

Does this mean anything?

Yes: The “spread” relates to frequentist variance just as in calibration weighting.




Figure 4: Comparison between raking and MrPlew weights for the Name Change dataset




Figure 5: Comparison between raking and MrPlew weights for the Gay Marriage dataset

Does this mean anything?

Yes: The “spread” relates to frequentist variance just as in calibration weighting.

What about covariate balance?




Figure 4: Comparison between raking and MrPlew weights for the Name Change dataset




Figure 5: Comparison between raking and MrPlew weights for the Gay Marriage dataset

Future work

Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

But the high level idea can be extended much more widely:

1. Assume your initial model was accurate
2. Select some perturbation your model should be able to capture
3. Use local sensitivity to detect whether the change is what you expect
4. If the change is not what you expect, either (1) or (2) was wrong

Future work

Notice that there was no discussion of misspecification!

Calibration weights (typically) do not depend on Y_S .

But the high level idea can be extended much more widely:

1. Assume your initial model was accurate
2. Select some perturbation your model should be able to capture
3. Use local sensitivity to detect whether the change is what you expect
4. If the change is not what you expect, either (1) or (2) was wrong

Such checks recover generalized versions of many standard diagnostics for linear models.

Examples:

- Additive parameter shifts \leftrightarrow Unbiasedness
- Invariance to survey data weighting \leftrightarrow Regressor + residual orthogonality
- Importance sampling \leftrightarrow Sandwich covariance $\stackrel{?}{=}$ Inverse Fisher information

Student contributions and ongoing work:


- **Vladimir Palmin** is working on extending MrPlew to `lme4`
- **Sequoia Andrade** is working on generalizing to other local sensitivity checks
- **Lucas Schwengber** is working on novel flow-based techniques for local sensitivity
- **(Currently recruiting!)** Doubly-robust Bayesian Hierarchical MrP



Vladimir Palmin



Sequoia Andrade



Lucas Schwengber

Preprint and R package (hopefully) coming soon!



B., Eli, Avi F., and Erin H. (2021). *Multilevel calibration weighting for survey data*. arXiv: 2102.09052 [stat.ME].



Chattopadhyay, A. and J. Zubizarreta (2023). “On the implied weights of linear regression for causal inference”. In: *Biometrika* 110.3, pp. 615–629.



Deville, J., C. Särndal, and O. Sautory (1993). “Generalized raking procedures in survey sampling”. In: *Journal of the American statistical Association* 88.423, pp. 1013–1020.



Diaconis, P. and D. Freedman (1986). “On the consistency of Bayes estimates”. In: *The Annals of Statistics*, pp. 1–26.



Efron, B. (2015). “Frequentist accuracy of Bayesian estimates”. In: *Journal of the Royal Statistical Society Series B: Statistical Methodology* 77.3, pp. 617–646.



Fuller, W. (2011). *Sampling statistics*. John Wiley & Sons.



G. and T. Broderick (2024). *The Bayesian Infinitesimal Jackknife for Variance*. arXiv: 2305.06466 [stat.ME]. URL: <https://arxiv.org/abs/2305.06466>.



G., T. Broderick, and M. I. Jordan (2018). “Covariances, robustness and variational bayes”. In: *Journal of machine learning research* 19.51.



G., W. Stephenson, et al. (2019). “A swiss army infinitesimal jackknife”. In: *The 22nd International Conference on Artificial Intelligence and Statistics*. PMLR, pp. 1139–1147.



Gelman, A. (2007a). “Rejoinder: Struggles with survey weighting and regression modelling”. In: *Statistical Science* 22.2, pp. 184–188.



— (2007b). “Struggles with survey weighting and regression modeling”. In:



Gustafson, P. (1996). “Local sensitivity of posterior expectations”. In: *The Annals of Statistics* 24.1, pp. 174–195.



Krantz, S. and H. Parks (2012). *The Implicit Function Theorem: History, Theory, and Applications*. Springer Science & Business Media.