

```
## Error in `prettyNum()`:  
## ! invalid value 0 for 'digits' argument
```

An Automatic Finite-Sample Robustness Metric: Can Dropping a Little Data Make a Big Difference?

Ryan Giordano (rgiordano@berkeley.edu)¹

JSM 2026: Data Selection Beyond Estimation Efficiency

Dropping data: Mexico Microcredit

Example: Angelucci et al. [2015], a randomized controlled trial study of the efficacy of microcredit in Mexico based on $N=16,560$ data points. A regression was run to estimate the average effect of microcredit.

Original result: Treatment effect statistically insignificant at 95%.

Policy implication: Disinvest in microcredit initiatives.

Dropping data: Mexico Microcredit

Example: Angelucci et al. [2015], a randomized controlled trial study of the efficacy of microcredit in Mexico based on $N = 16,560$ data points. A regression was run to estimate the average effect of microcredit.

Original result: Treatment effect statistically insignificant at 95%.

Policy implication: Disinvest in microcredit initiatives.

Data dropping: Can produce both positive and negative statistically significant results dropping no more than 15 data points ($< 0.1\%$).

Policy implication: Run a higher-powered study (not just larger N).

Dropping data: Mexico Microcredit

Example: Angelucci et al. [2015], a randomized controlled trial study of the efficacy of microcredit in Mexico based on $N = 16,560$ data points. A regression was run to estimate the average effect of microcredit.

Original result: Treatment effect statistically insignificant at 95%.

Policy implication: Disinvest in microcredit initiatives.

Data dropping: Can produce both positive and negative statistically significant results dropping no more than 15 data points ($< 0.1\%$).

Policy implication: Run a higher-powered study (not just larger N).

Cannot find influential subsets by brute force!

We provide a fast, automatic tool to approximately identify the most influential set of points.

- How do we identify influential sets? When is our method accurate?
- Data dropping as distributional robustness
- Do we identify “the most important datapoints?”

Formalizing the question.

Example: Least squares

Formalizing the question.

Example: Least squares

A data point d_n has regressors x_n and response y_n : $d_n = (x_n, y_n)$.

Formalizing the question.

Example: Least squares

A data point d_n has regressors x_n and response y_n : $d_n = (x_n, y_n)$.

The estimator $\hat{\theta} \in \mathbb{R}^P$ satisfies:

$$\begin{aligned}\hat{\theta} &:= \arg \min_{\theta} \frac{1}{2} \sum_{n=1}^N (y_n - \theta^T x_n)^2 \\ &\Leftrightarrow \sum_{n=1}^N (y_n - \hat{\theta}^T x_n) x_n = 0.\end{aligned}$$

Formalizing the question.

Example: Least squares

A data point d_n has regressors x_n and response y_n : $d_n = (x_n, y_n)$.

The estimator $\hat{\theta} \in \mathbb{R}^P$ satisfies:

$$\begin{aligned}\hat{\theta} &:= \arg \min_{\theta} \frac{1}{2} \sum_{n=1}^N (y_n - \theta^T x_n)^2 \\ \Leftrightarrow \sum_{n=1}^N (y_n - \hat{\theta}^T x_n) x_n &= 0.\end{aligned}$$

Make a qualitative decision using:

- A particular component: $\hat{\theta}_k$
- The end of a confidence interval: $\hat{\theta}_k + \frac{1.96}{\sqrt{N}} \hat{\sigma}(\hat{\theta})$

Formalizing the question.

Example: Least squares

A data point d_n has regressors x_n and response y_n : $d_n = (x_n, y_n)$.

The estimator $\hat{\theta} \in \mathbb{R}^P$ satisfies:

$$\begin{aligned}\hat{\theta} &:= \arg \min_{\theta} \frac{1}{2} \sum_{n=1}^N (y_n - \theta^T x_n)^2 \\ \Leftrightarrow \sum_{n=1}^N (y_n - \hat{\theta}^T x_n) x_n &= 0.\end{aligned}$$

Make a qualitative decision using:

- A particular component: $\hat{\theta}_k$
- The end of a confidence interval: $\hat{\theta}_k + \frac{1.96}{\sqrt{N}} \hat{\sigma}(\hat{\theta})$

General setup: Z-estimators

Formalizing the question.

Example: Least squares

A data point d_n has regressors x_n and response y_n : $d_n = (x_n, y_n)$.

The estimator $\hat{\theta} \in \mathbb{R}^P$ satisfies:

$$\begin{aligned}\hat{\theta} &:= \arg \min_{\theta} \frac{1}{2} \sum_{n=1}^N (y_n - \theta^T x_n)^2 \\ \Leftrightarrow \sum_{n=1}^N (y_n - \hat{\theta}^T x_n) x_n &= 0.\end{aligned}$$

Make a qualitative decision using:

- A particular component: $\hat{\theta}_k$
- The end of a confidence interval: $\hat{\theta}_k + \frac{1.96}{\sqrt{N}} \hat{\sigma}(\hat{\theta})$

General setup: Z-estimators

We observe N data points $d_1, \dots, d_N$ (in any domain).

Formalizing the question.

Example: Least squares

A data point d_n has regressors x_n and response y_n : $d_n = (x_n, y_n)$.

The estimator $\hat{\theta} \in \mathbb{R}^P$ satisfies:

$$\begin{aligned}\hat{\theta} &:= \arg \min_{\theta} \frac{1}{2} \sum_{n=1}^N (y_n - \theta^T x_n)^2 \\ \Leftrightarrow \sum_{n=1}^N (y_n - \hat{\theta}^T x_n) x_n &= 0.\end{aligned}$$

Make a qualitative decision using:

- A particular component: $\hat{\theta}_k$
- The end of a confidence interval: $\hat{\theta}_k + \frac{1.96}{\sqrt{N}} \hat{\sigma}(\hat{\theta})$

General setup: Z-estimators

We observe N data points $d_1, \dots, d_N$ (in any domain).

The estimator $\hat{\theta} \in \mathbb{R}^P$ satisfies:

$$\sum_{n=1}^N G(\hat{\theta}, d_n) = 0_P.$$

$G(\cdot, d_n)$ is “nice,” $\mathbb{R}^P$ -valued.
E.g. MLE, MAP, VB, IV &c.

Formalizing the question.

Example: Least squares

A data point d_n has regressors x_n and response y_n : $d_n = (x_n, y_n)$.

The estimator $\hat{\theta} \in \mathbb{R}^P$ satisfies:

$$\begin{aligned}\hat{\theta} &:= \arg \min_{\theta} \frac{1}{2} \sum_{n=1}^N (y_n - \theta^T x_n)^2 \\ \Leftrightarrow \sum_{n=1}^N (y_n - \hat{\theta}^T x_n) x_n &= 0.\end{aligned}$$

Make a qualitative decision using:

- A particular component: $\hat{\theta}_k$
- The end of a confidence interval: $\hat{\theta}_k + \frac{1.96}{\sqrt{N}} \hat{\sigma}(\hat{\theta})$

General setup: Z-estimators

We observe N data points $d_1, \dots, d_N$ (in any domain).

The estimator $\hat{\theta} \in \mathbb{R}^P$ satisfies:

$$\sum_{n=1}^N G(\hat{\theta}, d_n) = 0_P.$$

$G(\cdot, d_n)$ is “nice,” $\mathbb{R}^P$ -valued.
E.g. MLE, MAP, VB, IV &c.

Make a qualitative decision using $\phi(\hat{\theta})$ for a smooth, real-valued ϕ .

(WLOG try to increase $\phi(\hat{\theta})$.)

Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ?

Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ? **Two big problems:**

Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ? **Two big problems:**

- There are $\binom{N}{\lfloor \alpha N \rfloor}$ sets to check. (E.g. $\binom{16,560}{15} \approx 1.5 \cdot 10^{51}$)

Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ? **Two big problems:**

- There are $\binom{N}{\lfloor \alpha N \rfloor}$ sets to check. (E.g. $\binom{16,560}{15} \approx 1.5 \cdot 10^{51}$)
- Evaluating $\hat{\theta}$ re-solving the estimating equation.
 - E.g., re-running OLS. ($0.001s \cdot 1.5 \cdot 10^{51} \approx 4.8 \cdot 10^{40}$ years)
 - Other examples are even harder (VB, machine learning)

Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ? **Two big problems:**

- There are $\binom{N}{\lfloor \alpha N \rfloor}$ sets to check. (E.g. $\binom{16,560}{15} \approx 1.5 \cdot 10^{51}$)
- Evaluating $\hat{\theta}$ re-solving the estimating equation.
 - E.g., re-running OLS. ($0.001s \cdot 1.5 \cdot 10^{51} \approx 4.8 \cdot 10^{40}$ years)
 - Other examples are even harder (VB, machine learning)

Our idea: Smoothly approximate the effect of leaving out points.

Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ? **Two big problems:**

- There are $\binom{N}{\lfloor \alpha N \rfloor}$ sets to check. (E.g. $\binom{16,560}{15} \approx 1.5 \cdot 10^{51}$)
- Evaluating $\hat{\theta}$ re-solving the estimating equation.
 - E.g., re-running OLS. ($0.001s \cdot 1.5 \cdot 10^{51} \approx 4.8 \cdot 10^{40}$ years)
 - Other examples are even harder (VB, machine learning)

Our idea: Smoothly approximate the effect of leaving out points.

We have N data points $d_1, \dots, d_N$, a quantity of interest $\phi(\cdot)$, and

$$\sum_{n=1}^N G(\hat{\theta}, d_n) = 0_P \quad .$$

Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ? **Two big problems:**

- There are $\binom{N}{\lfloor \alpha N \rfloor}$ sets to check. (E.g. $\binom{16,560}{15} \approx 1.5 \cdot 10^{51}$)
- Evaluating $\hat{\theta}$ re-solving the estimating equation.
 - E.g., re-running OLS. ($0.001s \cdot 1.5 \cdot 10^{51} \approx 4.8 \cdot 10^{40}$ years)
 - Other examples are even harder (VB, machine learning)

Our idea: Smoothly approximate the effect of leaving out points.

We have N data points $d_1, \dots, d_N$, a quantity of interest $\phi(\cdot)$, and

$$\sum_{n=1}^N w_n G(\hat{\theta}(w), d_n) = 0_P \text{ for a weight vector } w \in \mathbb{R}^N.$$

Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ? **Two big problems:**


- There are $\binom{N}{\lfloor \alpha N \rfloor}$ sets to check. (E.g. $\binom{16,560}{15} \approx 1.5 \cdot 10^{51}$)
- Evaluating $\hat{\theta}$ re-solving the estimating equation.
 - E.g., re-running OLS. ($0.001s \cdot 1.5 \cdot 10^{51} \approx 4.8 \cdot 10^{40}$ years)
 - Other examples are even harder (VB, machine learning)

Our idea: Smoothly approximate the effect of leaving out points.

We have N data points $d_1, \dots, d_N$, a quantity of interest $\phi(\cdot)$, and

$$\sum_{n=1}^N w_n G(\hat{\theta}(w), d_n) = 0_P \text{ for a weight vector } w \in \mathbb{R}^N.$$

Original weights: $\vec{1} = (1, \dots, 1)$ Leave points out by setting their elements of w to zero.



Data dropping as data reweighting.

Question: Can we make a big change in $\phi(\hat{\theta})$ by dropping $\lfloor \alpha N \rfloor$ datapoints, for some small proportion α ? **Two big problems:**


- There are $\binom{N}{\lfloor \alpha N \rfloor}$ sets to check. (E.g. $\binom{16,560}{15} \approx 1.5 \cdot 10^{51}$)
- Evaluating $\hat{\theta}$ re-solving the estimating equation.
 - E.g., re-running OLS. ($0.001s \cdot 1.5 \cdot 10^{51} \approx 4.8 \cdot 10^{40}$ years)
 - Other examples are even harder (VB, machine learning)

Our idea: Smoothly approximate the effect of leaving out points.

We have N data points $d_1, \dots, d_N$, a quantity of interest $\phi(\cdot)$, and


$$\sum_{n=1}^N w_n G(\hat{\theta}(w), d_n) = 0_P \text{ for a weight vector } w \in \mathbb{R}^N.$$

Original weights: $\vec{1} = (1, \dots, 1)$ Leave points out by setting their elements of w to zero.




The map $w \mapsto \phi(\hat{\theta}(w))$ is well-defined even for continuous weights.


Taylor series approximation.



Taylor series approximation.




Taylor series approximation.



The values $N\psi_n$ are the **empirical influence function** [Hampel, 1986]. We call ψ_n an “influence score.”

Taylor series approximation.




The values $N\psi_n$ are the **empirical influence function** [Hampel, 1986]. We call ψ_n an “influence score.”

We can use ψ_n to form a Taylor series approximation:

$$\phi(\hat{\theta}(w)) \approx \phi^{\text{lin}}(w) := \phi(\hat{\theta}(\vec{1})) + \sum_{n=1}^N \psi_n(w_n - 1)$$

Taylor series approximation.



The values $N\psi_n$ are the **empirical influence function** [Hampel, 1986]. We call ψ_n an “influence score.”


We can use ψ_n to form a Taylor series approximation:

$$\phi(\hat{\theta}(w)) \approx \phi^{\text{lin}}(w) := \phi(\hat{\theta}(\vec{1})) + \sum_{n=1}^N \psi_n (w_n - 1)$$

The most influential points to drop from the linear approximation are easy to identify — they are simply the points with the largest ψ_n !


How accurate is the approximation?

Checking the approximation for Mexico microcredit.




How accurate is the approximation?

Checking the approximation for Mexico microcredit.



Distributional robustness



- You may be concerned if you could reverse your conclusion by removing a small proportion of your data.

Conclusion

- You may be concerned if you could reverse your conclusion by removing a small proportion of your data.
- We can quickly and automatically find an approximate influential set which is accurate for small sets.

- You may be concerned if you could reverse your conclusion by removing a small proportion of your data.
- We can quickly and automatically find an approximate influential set which is accurate for small sets.
- Data dropping robustness is principally determined by the signal to noise ratio, and captures sensitivity distinct from sampling and gross error sensitivity.

Links and references

Tamara Broderick, Ryan Giordano, Rachael Meager (alphabetical authors)
“An Automatic Finite-Sample Robustness Metric: Can Dropping a Little Data Change Conclusions?”

<https://arxiv.org/abs/2011.14999>

Select blog posts with more details: <https://rgiordan.github.io>

- Data dropping sensitivity overcomes p-hacking
- Collinearity in OLS after dropping
- Influence functions and sums
- Connections to the bootstrap

Related software on github:

- [rgiordan/zaminfluence](#) (for R)
- [rgiordan/vittles](#) (for Python)

Some of my work on other forms of robustness:

- Prior sensitivity in Bayesian nonparametrics [Giordano et al., 2021]
- Approximate cross-validation (and other reweightings) [Giordano et al., 2019b,a]
- Covariances and prior sensitivity for mean field VB [Giordano et al., 2015, 2018]
- Model sensitivity of MCMC output [Giordano et al., 2018]
- Frequentist variances of MCMC posteriors (in progress)

- M. Angelucci, D. Karlan, and J. Zinman. Microcredit impacts: Evidence from a randomized microcredit program placement experiment by Compartamos Banco. *American Economic Journal: Applied Economics*, 7(1):151–82, 2015.
- R. Giordano, T. Broderick, and M. I. Jordan. Linear response methods for accurate covariance estimates from mean field variational Bayes. *Advances in Neural Information Processing Systems*, 28:1441–1449, 2015.
- R. Giordano, T. Broderick, and M. I. Jordan. Covariances, robustness and variational Bayes. *The Journal of Machine Learning Research*, 19(1):1981–2029, 2018.
- R. Giordano, M. I. Jordan, and T. Broderick. A higher-order Swiss army infinitesimal jackknife. *arXiv preprint arXiv:1907.12116*, 2019a.
- R. Giordano, W. Stephenson, R. Liu, M. I. Jordan, and T. Broderick. A Swiss army infinitesimal jackknife. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1139–1147. PMLR, 2019b.
- R. Giordano, R. Liu, M. I. Jordan, and T. Broderick. Evaluating sensitivity to the stick-breaking prior in Bayesian nonparametrics. 2021.
- F. Hampel. *Robust statistics: The approach based on influence functions*, volume 196. Wiley-Interscience, 1986.