

Class09

Ralph Goguanco

```
DataExp.df <- read.csv("E:\\Downloads\\DataExportSummary.csv")
DataExp.df
```

##	Molecular.Type	X.ray	NMR	EM	Multiple.methods	Neutron	Other	
## 1	Protein (only)	144616	11881	6759		185	70	32
## 2	Protein/Oligosaccharide	8551	31	1133		5	0	0
## 3	Protein/NA	7623	274	2183		3	0	0
## 4	Nucleic acid (only)	2396	1399	61		8	2	1
## 5	Other	154	31	3		0	0	0
## 6	Oligosaccharide (only)	11	6	0		1	0	4
##	Total							
## 1	163543							
## 2	9720							
## 3	10083							
## 4	3867							
## 5	188							
## 6	22							

Q1: What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

```
(163351+10139)/187423
```

```
## [1] 0.9256601
```

Q2: What proportion of structures in the PDB are protein?

```
163543/187423
```

```
## [1] 0.8725877
```

Q3: Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB?

There are 4496 total structures in the current PDB

Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?

It would be too crowded if we added three atoms per water molecule since there's a lot of them.

Q5: There is a conserved water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have (see note below)?

The residue number of the atom associated with the water molecule is ALA28:CA

##3. Introduction to BIO3D in R

```
library(bio3d)

read.pdb(file = "1hsg")

## Note: Accessing on-line PDB file

##
## Call: read.pdb(file = "1hsg")
##
## Total Models#: 1
## Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)
##
## Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
## Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
##
## Non-protein/nucleic Atoms#: 172 (residues: 128)
## Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
##
## Protein sequence:
## PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
## QILIEICGHKAIGTVLVGPTPVNIIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
## ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
## VNIIGRNLLTQIGCTLNF
##
## + attr: atom, xyz, seqres, helix, sheet,
## calpha, remark, call
```

Q7: How many amino acid residues are there in this pdb object?

There are 198 amino acid residues in this pdb object.

Q8: Name one of the two non-protein residues?

Water and MK1

Q9: How many protein chains are in this structure?

There are 2 protein chains in this structure

Q10. Which of the packages above is found only on BioConductor and not CRAN?

msa

Q11. Which of the above packages is not found on BioConductor or CRAN?:

bio3d-view

Q12. True or False? Functions from the devtools package can be used to install packages from GitHub and BitBucket?

TRUE

```
library(bio3d)
aa <- get.seq("lake_A")
```

```
## Warning in get.seq("lake_A"): Removing existing file: seqs.fasta
```

```
## Fetching... Please wait. Done.
```

```
aa
```

```
##           1           .           .           .           .           .           60
## pdb|1AKE|A  MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMMLRAAVKSGSELGKQAKDIMDAGKLV
##           1           .           .           .           .           .           60
##
##           61           .           .           .           .           .           120
## pdb|1AKE|A  DELVIALVKERIAQEDCRNGFLLDGFPRTIPQADAMKEAGINVDYVLEFDVPDELIVDRI
##           61           .           .           .           .           .           120
##
##           121          .           .           .           .           .           180
## pdb|1AKE|A  VGRRVHAPSGRVYHVKNPPKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG
##           121          .           .           .           .           .           180
##
##           181          .           .           .           .           .           214
## pdb|1AKE|A  YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
##           181          .           .           .           .           .           214
##
## Call:
##   read.fasta(file = outfile)
##
## Class:
##   fasta
##
## Alignment dimensions:
##   1 sequence rows; 214 position columns (214 non-gap, 0 gap)
##
## + attr: id, ali, call
```

Q13. How many amino acids are in this sequence, i.e. how long is this sequence?


214

```
# Blast or hammer search
b <- blast.pdb(aa)
```

```
## Searching ... please wait (updates every 5 seconds) RID = 1FYHBM5D016
## .
## Reporting 100 hits
```

```
# Plot a summary of search results
hits <- plot(b)
```

```
## * Possible cutoff values: 197 -3
##      Yielding Nhits: 16 100
##
## * Chosen cutoff value of: 197
##      Yielding Nhits: 16
```



```
hits <- NULL
hits$pdb.id <- c('1AKE_A', '6S36_A', '6RZE_A', '3HPR_A', '1E4V_A', '5EJE_A', '1E4Y_A', '3X2S_A', '6HAP_A', '6HAM_A')
```

```
#Download related PDB files
```

```
files <- get.pdb(hits$pdb.id, path="pdbs", split=TRUE, gzip=TRUE)
```

```
## Warning in get.pdb(hits$pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 1AKE.pdb exists. Skipping download
```

```
## Warning in get.pdb(hits$pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 6S36.pdb exists. Skipping download
```

```
## Warning in get.pdb(hits$pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 6RZE.pdb exists. Skipping download
```

```
## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 3HPR.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 1E4V.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 5EJE.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 1E4Y.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 3X2S.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 6HAP.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 6HAM.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 4K46.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 3GMT.pdb exists. Skipping download

## Warning in get.pdb(hits$ pdb.id, path = "pdbs", split = TRUE, gzip = TRUE): pdbs/
## 4PZL.pdb exists. Skipping download

##      |
```

```
# Align related PDBs
#pdbs <- pdbaln(files, fit = TRUE), exefile="msa")
```

```
# Vector containing PDB codes for figure axis
#ids <- basename.pdb(pdb$ id)

# Draw schematic alignment
#(pdbs, labels=ids)
```

What do you note about this plot? Are the black and colored lines similar or different? Where do you think they differ most and why?

The black and colored lines are different, they differ in fluctuation heights but the patterns are almost similar.