

# ML in Economics and Finance: Where do We Go Now? - Part I

---

Raul Riva

FGV EPGE

December, 2025

INSPER - São Paulo

# Intro

---

# Who is this guy?

- I have just joined [FGV EPGE](#) as an Assistant Professor;
- I got my PhD in Finance at [Northwestern University](#);
- Asset Pricing + Macro-Finance + Econometrics;

# Who is this guy?

- I have just joined [FGV EPGE](#) as an Assistant Professor;
- I got my PhD in Finance at [Northwestern University](#);
- Asset Pricing + Macro-Finance + Econometrics;

I am **not** an ML developer, but maybe a mildly sophisticated economist consumer

## Where are we?

- Last 20-30 years: explosion of computation power and popularization of ML techniques;
- Last 15 years: we economists imported several techniques from CS and Stats;

# Where are we?

- Last 20-30 years: explosion of computation power and popularization of ML techniques;
- Last 15 years: we economists imported several techniques from CS and Stats;
- Many challenges in this translation:
  - Causality vs pattern recognition;
  - Sophisticated notions of equilibrium;
  - Interpretability;
  - Time series dynamics;

# Where are we?

- Last 20-30 years: explosion of computation power and popularization of ML techniques;
- Last 15 years: we economists imported several techniques from CS and Stats;
- Many challenges in this translation:
  - Causality vs pattern recognition;
  - Sophisticated notions of equilibrium;
  - Interpretability;
  - Time series dynamics;

## Right now:

- Better understanding of the limitations of "plug and play" ML;
- Great stuff: new hybrid methods designed by and for economists;
- Bad stuff: we are flooded with tutorials, books, videos, bootcamps...

## Where do we go now?

- The Econ/Finance forecasting crowd was really fast in adopting ML...
- But what else? What is worth knowing about ML in Econ and Finance?



# Where do we go now?

- The Econ/Finance forecasting crowd was really fast in adopting ML...
- But what else? What is worth knowing about ML in Econ and Finance?

## **What I will do:**

- My own economist-crafted definition of ML methods and how to think about them;
- Three very cool agendas where ML can help economists
- Causality in HD, seriously heterogeneous treatment effects, and solving large models;

# Where do we go now?

- The Econ/Finance forecasting crowd was really fast in adopting ML...
- But what else? What is worth knowing about ML in Econ and Finance?

## What I will do:

- My own economist-crafted definition of ML methods and how to think about them;
- Three very cool agendas where ML can help economists
- Causality in HD, seriously heterogeneous treatment effects, and solving large models;

## What I will not do:

- Teach you how to code;

# Where do we go now?

- The Econ/Finance forecasting crowd was really fast in adopting ML...
- But what else? What is worth knowing about ML in Econ and Finance?

## What I will do:

- My own economist-crafted definition of ML methods and how to think about them;
- Three very cool agendas where ML can help economists
- Causality in HD, seriously heterogeneous treatment effects, and solving large models;

## What I will not do:

- Teach you how to code;
- Pretend I know how to prove the complicated theorems and walk you through proofs;

# Where do we go now?

- The Econ/Finance forecasting crowd was really fast in adopting ML...
- But what else? What is worth knowing about ML in Econ and Finance?

## What I will do:

- My own economist-crafted definition of ML methods and how to think about them;
- Three very cool agendas where ML can help economists
- Causality in HD, seriously heterogeneous treatment effects, and solving large models;

## What I will not do:

- Teach you how to code;
- Pretend I know how to prove the complicated theorems and walk you through proofs;
- Lie to you and say you can easily perform any of this in Stata! 🙄

## Who is this for?

- Students starting their empirical research agendas;
- Fellow empirical researchers trying to grasp what kind of ML tools might be useful;
- Someone coming from Stats or CS into Economics;

## Who is this for?

- Students starting their empirical research agendas;
- Fellow empirical researchers trying to grasp what kind of ML tools might be useful;
- Someone coming from Stats or CS into Economics;

## Who is this not for?

- Super sophisticated economists already deploying these techniques everywhere;
- Hardcore econometricians looking for open theoretical problems;

## Who is this for?

- Students starting their empirical research agendas;
- Fellow empirical researchers trying to grasp what kind of ML tools might be useful;
- Someone coming from Stats or CS into Economics;

## Who is this not for?

- Super sophisticated economists already deploying these techniques everywhere;
- Hardcore econometricians looking for open theoretical problems;

DISCLAIMER: These are **my** own views, based on **my** experience, and **my** own readings.  
Other people will disagree.

- 1. What is ML, anyway?
  - 2. Causality in High Dimensions
  - 3. (Seriously) Heterogeneous Partial Effects
  - 4. Solving Large-Scale General Equilibrium Models
- } **Today**
- } **Tomorrow**



- 1. What is ML, anyway?
  - 2. Causality in High Dimensions
  - 3. (Seriously) Heterogeneous Partial Effects
  - 4. Solving Large-Scale General Equilibrium Models
- } **Today**
- } **Tomorrow**

**Please bring questions at any time!**

**Questions?**

## **A General Framework**

---

# What is *Machine Learning*?

- Different fields = different definitions: CS, Stats, Operations Research, ...
- Many types: Supervised, Unsupervised, Reinforcement Learning, Deep Learning, ...
- More buzzwords = better consulting gigs! 🤖

# What is *Machine Learning*?

- Different fields = different definitions: CS, Stats, Operations Research, ...
- Many types: Supervised, Unsupervised, Reinforcement Learning, Deep Learning, ...
- More buzzwords = better consulting gigs! 🤖
- Today and tomorrow: **Supervised Learning**;

# What is *Machine Learning*?

- Different fields = different definitions: CS, Stats, Operations Research, ...
- Many types: Supervised, Unsupervised, Reinforcement Learning, Deep Learning, ...
- More buzzwords = better consulting gigs! 🤖
- Today and tomorrow: **Supervised Learning**;
- I will be brave enough and provide the one I think is really useful for Economists:

# What is *Machine Learning*?

- Different fields = different definitions: CS, Stats, Operations Research, ...
- Many types: Supervised, Unsupervised, Reinforcement Learning, Deep Learning, ...
- More buzzwords = better consulting gigs! 🤖
- Today and tomorrow: **Supervised Learning**;
- I will be brave enough and provide the one I think is really useful for Economists:

(Supervised) **Machine Learning** is a set of tools that enable computationally-feasible data-driven search over high-dimensional functional spaces.

## A General Framework

$$y = f(\mathbf{x}) + \varepsilon$$

- $y \in \mathbb{R}^k$  is some "target" or "outcome";
- $\mathbf{x} \in \mathbb{R}^p$  is a vector of "features", or "predictors", or "covariates";
- $f : \mathbb{R}^p \rightarrow \mathbb{R}^k$  is some unknown function;
- $\varepsilon$  is some unobserved noise because the world is messy;



## A General Framework

$$y = f(\mathbf{x}) + \varepsilon$$

- $y \in \mathbb{R}^k$  is some "target" or "outcome";
- $\mathbf{x} \in \mathbb{R}^p$  is a vector of "features", or "predictors", or "covariates";
- $f : \mathbb{R}^p \rightarrow \mathbb{R}^k$  is some unknown function;
- $\varepsilon$  is some unobserved noise because the world is messy;

**Question:** given a function space  $\mathcal{F}$ , how to find  $\hat{f} \in \mathcal{F}$  that approximates  $f$  well?

$$y = f(\mathbf{x}) + \varepsilon$$

- $y \in \mathbb{R}^k$  is some "target" or "outcome";
- $\mathbf{x} \in \mathbb{R}^p$  is a vector of "features", or "predictors", or "covariates";
- $f : \mathbb{R}^p \rightarrow \mathbb{R}^k$  is some unknown function;
- $\varepsilon$  is some unobserved noise because the world is messy;

**Question:** given a function space  $\mathcal{F}$ , how to find  $\hat{f} \in \mathcal{F}$  that approximates  $f$  well?

- Collect data  $\{(y_1, \mathbf{x}_1), \dots, (y_n, \mathbf{x}_n)\}$ ;
- Define some notion of "approximates well"  $\implies$  (a loss function);
- Be explicit about  $\mathcal{F}$ ;
- Be explicit about your optimization mechanism;

# You are already doing ML!

Consider an outcome  $y_i$ , and a set of covariates  $\mathbf{x}_i$  for  $i = 1, \dots, n$ :

$$y_i = \alpha + \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i$$

- This is a linear regression model;
- The function space  $\mathcal{F}$  is the set of all affine functions of the treatment and covariates;
- The loss function is the MSE:  $\mathcal{L}(y_i, \hat{y}_i) = (y_i - \hat{y}_i)^2$ ;
- OLS: minimize a convex loss function over the space of parameters;

# You are already doing ML!

Consider an outcome  $y_i$ , and a set of covariates  $\mathbf{x}_i$  for  $i = 1, \dots, n$ :

$$y_i = \alpha + \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i$$

- This is a linear regression model;
- The function space  $\mathcal{F}$  is the set of all affine functions of the treatment and covariates;
- The loss function is the MSE:  $\mathcal{L}(y_i, \hat{y}_i) = (y_i - \hat{y}_i)^2$ ;
- OLS: minimize a convex loss function over the space of parameters;

**Conclusion:** Linear regression is a (very simple) ML method! But there is so much more...

## Hold on... Isn't Machine Learning just Non-Parametric Estimation?

- The general framework I used could be used in a non-parametric estimation class...
- Why do we need new tools? We already have the good and old kernel regression!

## Hold on... Isn't Machine Learning just Non-Parametric Estimation?

- The general framework I used could be used in a non-parametric estimation class...
- Why do we need new tools? We already have the good and old kernel regression!
- Well... there is the curse of dimensionality! If  $p \approx 6$ , you are already in trouble!

# Hold on... Isn't Machine Learning just Non-Parametric Estimation?

- The general framework I used could be used in a non-parametric estimation class...
- Why do we need new tools? We already have the good and old kernel regression!
- Well... there is the curse of dimensionality! If  $p \approx 6$ , you are already in trouble!

## OLS

- Leverages linearity (strong!);
- Easy to compute and interpret;

## Fully Non-Parametric Methods

- Extreme flexibility;
- Super data hungry!

Machine Learning = a *compromise*: richer parametrizations while still computationally feasible in high dimensions.

**Questions?**




## **Causality in High Dimensions**

---

- No estimator will lead to causality by itself – only careful design will;

- No estimator will lead to causality by itself – only careful design will;
- ML methods were *not* created to tackle causality problems;
- See [Varian \(2014\)](#), [Mullainathan and Spiess \(2017\)](#), [Athey and Imbens \(2019\)](#), and [Masini et al. \(2023\)](#);

- No estimator will lead to causality by itself – only careful design will;
- ML methods were *not* created to tackle causality problems;
- See [Varian \(2014\)](#), [Mullainathan and Spiess \(2017\)](#), [Athey and Imbens \(2019\)](#), and [Masini et al. \(2023\)](#);
- [Kleinberg et al. \(2015\)](#): many policy-relevant questions are prediction problems!

- No estimator will lead to causality by itself – only careful design will;
- ML methods were *not* created to tackle causality problems;
- See [Varian \(2014\)](#), [Mullainathan and Spiess \(2017\)](#), [Athey and Imbens \(2019\)](#), and [Masini et al. \(2023\)](#);
- [Kleinberg et al. \(2015\)](#): many policy-relevant questions are prediction problems!
- Belloni , Chernozhukov, Hansen and co-authors took it even further:
  - Computing the propensity score *is* forecasting!
  - The first-stage regression in an IV context *is* forecasting!

# Treatment Effects in High Dimensions

Suppose you're interested in the treatment effect  $\theta_0 \in \mathbb{R}$ :

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- $y_i \in \mathbb{R}$  is an outcome;
- $d_i \in \mathbb{R}$  is a treatment;
- $\mathbf{x}_i \in \mathbb{R}^p$  is a vector of available covariates;
- $\varepsilon_i$  is some unobserved noise with  $\mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] = 0$ ;

# Treatment Effects in High Dimensions

Suppose you're interested in the treatment effect  $\theta_0 \in \mathbb{R}$ :

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- $y_i \in \mathbb{R}$  is an outcome;
- $d_i \in \mathbb{R}$  is a treatment;
- $\mathbf{x}_i \in \mathbb{R}^p$  is a vector of available covariates;
- $\varepsilon_i$  is some unobserved noise with  $\mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] = 0$ ;
- You have an i.i.d. sample  $\{y_i, d_i, \mathbf{x}_i\}_{i=1}^n$  and we allow for  $p \gg n$ ;

# Treatment Effects in High Dimensions

Suppose you're interested in the treatment effect  $\theta_0 \in \mathbb{R}$ :

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- $y_i \in \mathbb{R}$  is an outcome;
- $d_i \in \mathbb{R}$  is a treatment;
- $\mathbf{x}_i \in \mathbb{R}^p$  is a vector of available covariates;
- $\varepsilon_i$  is some unobserved noise with  $\mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] = 0$ ;
- You have an i.i.d. sample  $\{y_i, d_i, \mathbf{x}_i\}_{i=1}^n$  and we allow for  $p \gg n$ ;
- **Goal:** estimate  $\theta_0$  and get a confidence interval;



# Treatment Effects in High Dimensions

Suppose you're interested in the treatment effect  $\theta_0 \in \mathbb{R}$ :

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- $y_i \in \mathbb{R}$  is an outcome;
- $d_i \in \mathbb{R}$  is a treatment;
- $\mathbf{x}_i \in \mathbb{R}^p$  is a vector of available covariates;
- $\varepsilon_i$  is some unobserved noise with  $\mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] = 0$ ;
- You have an i.i.d. sample  $\{y_i, d_i, \mathbf{x}_i\}_{i=1}^n$  and we allow for  $p \gg n$ ;
- **Goal:** estimate  $\theta_0$  and get a confidence interval;

**Question:** what will happen if you try OLS here?

## Treatment Effects in High Dimensions

- Let's say you believe only a few  $\beta_j$ 's are  $\neq 0 \implies$  “*sparsity*” in  $\beta$ ;
- But you do not know which ones!

## Treatment Effects in High Dimensions

- Let's say you believe only a few  $\beta_j$ 's are  $\neq 0 \implies$  “*sparsity*” in  $\beta$ ;
- But you do not know which ones!
- What about using your economic intuition to select a subset of controls?

# Treatment Effects in High Dimensions

- Let's say you believe only a few  $\beta_j$ 's are  $\neq 0 \implies$  “*sparsity*” in  $\beta$ ;
- But you do not know which ones!
- What about using your economic intuition to select a subset of controls?
- Applying Econ theory is always a good idea, but:
  - You might not get a meaningful reduction with theory alone;
  - Your referee might not agree with your choices;
  - You might get lost in a sea of robustness checks...

# Treatment Effects in High Dimensions

- Let's say you believe only a few  $\beta_j$ 's are  $\neq 0 \implies$  “*sparsity*” in  $\beta$ ;
- But you do not know which ones!
- What about using your economic intuition to select a subset of controls?
- Applying Econ theory is always a good idea, but:
  - You might not get a meaningful reduction with theory alone;
  - Your referee might not agree with your choices;
  - You might get lost in a sea of robustness checks...
- Good news: ML researchers devoted a lot of attention to *sparse regressions*!

The Least Absolute Shrinkage and Selection Operator (LASSO) estimator solves:

$$\hat{\boldsymbol{\delta}} \equiv \arg \min_{\boldsymbol{\delta} \in \mathbb{R}^p} \left\{ \frac{1}{n} \sum_{i=1}^n (y_i - \mathbf{w}_i' \boldsymbol{\delta})^2 + \lambda \sum_{j=1}^p |\delta_j| \right\}$$

- $\lambda \geq 0$  is a tuning parameter that controls the amount of penalization (“*regularization*”);
- $\mathbf{w}_i$  is a general vector of regressors of size  $p$ ;

The Least Absolute Shrinkage and Selection Operator (LASSO) estimator solves:

$$\hat{\boldsymbol{\delta}} \equiv \arg \min_{\boldsymbol{\delta} \in \mathbb{R}^p} \left\{ \frac{1}{n} \sum_{i=1}^n (y_i - \mathbf{w}_i' \boldsymbol{\delta})^2 + \lambda \sum_{j=1}^p |\delta_j| \right\}$$

- $\lambda \geq 0$  is a tuning parameter that controls the amount of penalization (“*regularization*”);
- $\mathbf{w}_i$  is a general vector of regressors of size  $p$ ;
- The  $\ell_1$  penalty  $\sum_{j=1}^p |\delta_j|$  induces sparsity in  $\hat{\boldsymbol{\delta}}$ ;

The Least Absolute Shrinkage and Selection Operator (LASSO) estimator solves:

$$\hat{\boldsymbol{\delta}} \equiv \arg \min_{\boldsymbol{\delta} \in \mathbb{R}^p} \left\{ \frac{1}{n} \sum_{i=1}^n (y_i - \mathbf{w}_i' \boldsymbol{\delta})^2 + \lambda \sum_{j=1}^p |\delta_j| \right\}$$

- $\lambda \geq 0$  is a tuning parameter that controls the amount of penalization (“*regularization*”);
- $\mathbf{w}_i$  is a general vector of regressors of size  $p$ ;
- The  $\ell_1$  penalty  $\sum_{j=1}^p |\delta_j|$  induces sparsity in  $\hat{\boldsymbol{\delta}}$ ;
- If  $\lambda = 0$ , we get OLS; if  $\lambda \rightarrow \infty$ , we get  $\hat{\boldsymbol{\delta}} = \mathbf{0}$ ;



## Welcome to SBE, Mr. LASSO

The Least Absolute Shrinkage and Selection Operator (LASSO) estimator solves:

$$\hat{\boldsymbol{\delta}} \equiv \arg \min_{\boldsymbol{\delta} \in \mathbb{R}^p} \left\{ \frac{1}{n} \sum_{i=1}^n (y_i - \mathbf{w}_i' \boldsymbol{\delta})^2 + \lambda \sum_{j=1}^p |\delta_j| \right\}$$

- $\lambda \geq 0$  is a tuning parameter that controls the amount of penalization (“*regularization*”);
- $\mathbf{w}_i$  is a general vector of regressors of size  $p$ ;
- The  $\ell_1$  penalty  $\sum_{j=1}^p |\delta_j|$  induces sparsity in  $\hat{\boldsymbol{\delta}}$ ;
- If  $\lambda = 0$ , we get OLS; if  $\lambda \rightarrow \infty$ , we get  $\hat{\boldsymbol{\delta}} = \mathbf{0}$ ;
- For intermediate values of  $\lambda$ , some  $\hat{\delta}_j$ 's will be exactly zero!

# Welcome to SBE, Mr. LASSO

The Least Absolute Shrinkage and Selection Operator (LASSO) estimator solves:

$$\hat{\boldsymbol{\delta}} \equiv \arg \min_{\boldsymbol{\delta} \in \mathbb{R}^p} \left\{ \frac{1}{n} \sum_{i=1}^n (y_i - \mathbf{w}_i' \boldsymbol{\delta})^2 + \lambda \sum_{j=1}^p |\delta_j| \right\}$$

- $\lambda \geq 0$  is a tuning parameter that controls the amount of penalization (“*regularization*”);
- $\mathbf{w}_i$  is a general vector of regressors of size  $p$ ;
- The  $\ell_1$  penalty  $\sum_{j=1}^p |\delta_j|$  induces sparsity in  $\hat{\boldsymbol{\delta}}$ ;
- If  $\lambda = 0$ , we get OLS; if  $\lambda \rightarrow \infty$ , we get  $\hat{\boldsymbol{\delta}} = \mathbf{0}$ ;
- For intermediate values of  $\lambda$ , some  $\hat{\delta}_j$ 's will be exactly zero!
- $\hat{\boldsymbol{\delta}}$  gives up unbiasedness for much lower variance;
- This problem is still feasible if  $p \gg n$  and it is convex  $\implies$  fast computation;


# The Geometry of LASSO

For  $c > 0$ , consider the following:

$$\tilde{\delta} \equiv \arg \min_{\delta \in \mathbb{R}^p} \left\{ \frac{1}{n} \sum_{i=1}^n (y_i - \mathbf{w}_i' \delta)^2 \right\}$$

subject to  $\sum_{j=1}^p |\delta_j| \leq c$

- Think about the Lagrangian of this problem!
- For every  $\lambda$ , there is a  $c$  such that  $\hat{\delta} = \tilde{\delta}$ ;



Recall our treatment effects model:

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- **Approach 1:** run LASSO of  $y_i$  on  $d_i$  and  $\mathbf{x}_i$ . Is this a good idea?

## Exploring Options

Recall our treatment effects model:

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- **Approach 1:** run LASSO of  $y_i$  on  $d_i$  and  $\mathbf{x}_i$ . Is this a good idea?
- $\hat{\theta}_0$  will be severely biased and might even be set to zero!

Recall our treatment effects model:

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- **Approach 1:** run LASSO of  $y_i$  on  $d_i$  and  $\mathbf{x}_i$ . Is this a good idea?
- $\hat{\theta}_0$  will be severely biased and might even be set to zero!
- **Approach 2:** run LASSO but with no penalty on  $\theta_0$ . Is this a good idea?

Recall our treatment effects model:

$$y_i = d_i\theta_0 + \mathbf{x}'_i\boldsymbol{\beta} + \varepsilon_i$$

- **Approach 1:** run LASSO of  $y_i$  on  $d_i$  and  $\mathbf{x}_i$ . Is this a good idea?
- $\hat{\theta}_0$  will be severely biased and might even be set to zero!
- **Approach 2:** run LASSO but with no penalty on  $\theta_0$ . Is this a good idea?
- Still terrible!  $\hat{\theta}_0$  will still be biased!

# Exploring Options

Recall our treatment effects model:

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- **Approach 1:** run LASSO of  $y_i$  on  $d_i$  and  $\mathbf{x}_i$ . Is this a good idea?
- $\hat{\theta}_0$  will be severely biased and might even be set to zero!
- **Approach 2:** run LASSO but with no penalty on  $\theta_0$ . Is this a good idea?
- Still terrible!  $\hat{\theta}_0$  will still be biased!
- **Approach 3:** do Approach 2, but then run OLS on the selected  $\mathbf{x}_i$  and  $d_i$ . Is this good?



Recall our treatment effects model:

$$y_i = d_i\theta_0 + \mathbf{x}_i'\boldsymbol{\beta} + \varepsilon_i$$

- **Approach 1:** run LASSO of  $y_i$  on  $d_i$  and  $\mathbf{x}_i$ . Is this a good idea?
- $\hat{\theta}_0$  will be severely biased and might even be set to zero!
- **Approach 2:** run LASSO but with no penalty on  $\theta_0$ . Is this a good idea?
- Still terrible!  $\hat{\theta}_0$  will still be biased!
- **Approach 3:** do Approach 2, but then run OLS on the selected  $\mathbf{x}_i$  and  $d_i$ . Is this good?
- [Leeb and Pötscher \(2008a\)](#) and [Leeb and Pötscher \(2008b\)](#): terrible idea again!
- Main problem: *omitted variable bias* if some relevant controls are not selected!
- If some  $\mathbf{x}_j$  is correlated with  $d_i$  and affects  $y_i$ , omitting it biases  $\hat{\theta}_0$ !

## Something That Finally Works!

Belloni et al. (2014a) thought about how  $d_i$  and  $\mathbf{x}_i$  interact:

$$d_i = \mathbf{x}_i' \gamma + u_i, \quad \mathbb{E}[u_i \mid \mathbf{x}_i] = 0$$

- What if  $\gamma$  is also sparse, i.e., only a few  $x_j$ 's affect  $d_i$ ?
- Can we find a small subset of  $\mathbf{x}_i$  that *predicts* treatment well?

## Something That Finally Works!

Belloni et al. (2014a) thought about how  $d_i$  and  $\mathbf{x}_i$  interact:

$$d_i = \mathbf{x}_i' \boldsymbol{\gamma} + u_i, \quad \mathbb{E}[u_i \mid \mathbf{x}_i] = 0$$

- What if  $\boldsymbol{\gamma}$  is also sparse, i.e., only a few  $x_j$ 's affect  $d_i$ ?
- Can we find a small subset of  $\mathbf{x}_i$  that *predicts* treatment well?

They proposed the **Double LASSO** procedure:

1. Run LASSO of  $y_i$  on  $\mathbf{x}_i$  to select controls  $\hat{S}_y$ ;
2. Run LASSO of  $d_i$  on  $\mathbf{x}_i$  to select controls  $\hat{S}_d$ ;
3. Run OLS of  $y_i$  on  $d_i$  and  $\mathbf{x}_i$  with  $x_j \in \hat{S}_y \cup \hat{S}_d$ ;

## A Really Cool Result

Belloni et al. (2014b) provide conditions under which:

$$\sqrt{n}(\hat{\theta}_0 - \theta_0) \xrightarrow{d} \mathcal{N}(0, \sigma^2)$$

where  $\sigma^2$  is complicated by consistently estimated.

# A Really Cool Result

Belloni et al. (2014b) provide conditions under which:

$$\sqrt{n}(\hat{\theta}_0 - \theta_0) \xrightarrow{d} \mathcal{N}(0, \sigma^2)$$

where  $\sigma^2$  is complicated but consistently estimated.


## The impressive stuff:

- This convergence is uniform over a large class of DGPs;
- Convergence still happens at the rate  $\sqrt{n}$ , even if  $p \gg n$ !
- Under homoskedasticity, it attains semi-parametric efficiency!
- Construct confidence intervals in the usual ways;


**Key assumption:** sparse representation;

# Some Monte-Carlo Reassurance

post-single-selection estimator



post-double-selection estimator



**Questions?**

## **Limitations and Generalizations**

---



- What if we want to allow for non-linearities?
- What if we want to use other ML methods?
- What if sparsity is not a good assumption?
- What if treatment has heterogenous effects?
- What if outcomes are function-valued?

Belloni et al. (2017) and Chernozhukov et al. (2018) generalize all of this:

$$\begin{aligned}y_i &= g_0(d_i, \mathbf{x}_i) + \varepsilon_i, & \mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] &= 0 \\d_i &= m_0(\mathbf{x}_i) + u_i, & \mathbb{E}[u_i \mid \mathbf{x}_i] &= 0\end{aligned}$$

- $g_0(\cdot)$  and  $m_0(\cdot)$  are unknown (possibly non-linear) functions;
- You can use several different ML method to estimate  $g_0(\cdot)$  and  $m_0(\cdot)$ ;
- Sparsity is not necessary anymore;

# Generalization

Belloni et al. (2017) and Chernozhukov et al. (2018) generalize all of this:

$$\begin{aligned}y_i &= g_0(d_i, \mathbf{x}_i) + \varepsilon_i, & \mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] &= 0 \\d_i &= m_0(\mathbf{x}_i) + u_i, & \mathbb{E}[u_i \mid \mathbf{x}_i] &= 0\end{aligned}$$

- $g_0(\cdot)$  and  $m_0(\cdot)$  are unknown (possibly non-linear) functions;
- You can use several different ML method to estimate  $g_0(\cdot)$  and  $m_0(\cdot)$ ;
- Sparsity is not necessary anymore;
- Secrete sauce I: Neyman Orthogonal Scores  $\psi$

$$\mathbb{E} \left[ \psi(\text{data}, \underbrace{\text{param of interest}}_{\equiv \theta_0}, \underbrace{\text{nuisance params}}_{\equiv \eta_0}) \right] = 0, \quad \frac{\partial}{\partial \eta} \mathbb{E} [\psi(\text{data}, \theta_0, \eta)] \Big|_{\eta=\eta_0} = 0$$

# Generalization

Belloni et al. (2017) and Chernozhukov et al. (2018) generalize all of this:

$$\begin{aligned}y_i &= g_0(d_i, \mathbf{x}_i) + \varepsilon_i, & \mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] &= 0 \\d_i &= m_0(\mathbf{x}_i) + u_i, & \mathbb{E}[u_i \mid \mathbf{x}_i] &= 0\end{aligned}$$

- $g_0(\cdot)$  and  $m_0(\cdot)$  are unknown (possibly non-linear) functions;
- You can use several different ML method to estimate  $g_0(\cdot)$  and  $m_0(\cdot)$ ;
- Sparsity is not necessary anymore;
- Secrete sauce I: Neyman Orthogonal Scores  $\psi$

$$\mathbb{E} \left[ \psi(\text{data}, \underbrace{\text{param of interest}}_{\equiv \theta_0}, \underbrace{\text{nuisance params}}_{\equiv \eta_0}) \right] = 0, \quad \frac{\partial}{\partial \eta} \mathbb{E} [\psi(\text{data}, \theta_0, \eta)] \Big|_{\eta=\eta_0} = 0$$

- Secrete sauce II: cross-fitting  $\implies$  efficiency vs strict assumptions;

# Generalization

Belloni et al. (2017) and Chernozhukov et al. (2018) generalize all of this:

$$\begin{aligned}y_i &= g_0(d_i, \mathbf{x}_i) + \varepsilon_i, & \mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] &= 0 \\d_i &= m_0(\mathbf{x}_i) + u_i, & \mathbb{E}[u_i \mid \mathbf{x}_i] &= 0\end{aligned}$$

- $g_0(\cdot)$  and  $m_0(\cdot)$  are unknown (possibly non-linear) functions;
- You can use several different ML method to estimate  $g_0(\cdot)$  and  $m_0(\cdot)$ ;
- Sparsity is not necessary anymore;
- Secrete sauce I: Neyman Orthogonal Scores  $\psi$

$$\mathbb{E} \left[ \psi(\text{data}, \underbrace{\text{param of interest}}_{\equiv \theta_0}, \underbrace{\text{nuisance params}}_{\equiv \eta_0}) \right] = 0, \quad \frac{\partial}{\partial \eta} \mathbb{E} [\psi(\text{data}, \theta_0, \eta)] \Big|_{\eta=\eta_0} = 0$$

- Secrete sauce II: cross-fitting  $\implies$  efficiency vs strict assumptions;
- Independence across  $i$  is essential;

## A Concrete Example (A Partially Linear Model)

$$\begin{aligned}y_i &= d_i\theta_0 + g_0(\mathbf{x}_i) + \varepsilon_i, & \mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] &= 0 \\d_i &= m_0(\mathbf{x}_i) + u_i, & \mathbb{E}[u_i \mid \mathbf{x}_i] &= 0\end{aligned}$$

## A Concrete Example (A Partially Linear Model)

$$y_i = d_i\theta_0 + g_0(\mathbf{x}_i) + \varepsilon_i, \quad \mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] = 0$$
$$d_i = m_0(\mathbf{x}_i) + u_i, \quad \mathbb{E}[u_i \mid \mathbf{x}_i] = 0$$

Steps:

- Divide the data into two folds;
- On fold 1, estimate  $\hat{g}_0(\mathbf{x}_i)$  and  $\hat{m}_0(\mathbf{x}_i)$  using ML methods;
- On fold 2, compute residuals:

$$\hat{\varepsilon}_i = y_i - \hat{g}_0(\mathbf{x}_i)$$

$$\hat{u}_i = d_i - \hat{m}_0(\mathbf{x}_i)$$

- Regress  $\hat{\varepsilon}_i$  on  $\hat{u}_i$  to get  $\hat{\theta}_0$ ;

## A Concrete Example (A Partially Linear Model)

$$y_i = d_i\theta_0 + g_0(\mathbf{x}_i) + \varepsilon_i, \quad \mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] = 0$$
$$d_i = m_0(\mathbf{x}_i) + u_i, \quad \mathbb{E}[u_i \mid \mathbf{x}_i] = 0$$

Steps:

- Divide the data into two folds;
- On fold 1, estimate  $\hat{g}_0(\mathbf{x}_i)$  and  $\hat{m}_0(\mathbf{x}_i)$  using ML methods;
- On fold 2, compute residuals:

$$\hat{\varepsilon}_i = y_i - \hat{g}_0(\mathbf{x}_i)$$

$$\hat{u}_i = d_i - \hat{m}_0(\mathbf{x}_i)$$

- Regress  $\hat{\varepsilon}_i$  on  $\hat{u}_i$  to get  $\hat{\theta}_0$ ;
- Repeat switching folds and average  $\hat{\theta}_0$ 's;



## A Concrete Example (A Partially Linear Model)

$$y_i = d_i\theta_0 + g_0(\mathbf{x}_i) + \varepsilon_i, \quad \mathbb{E}[\varepsilon_i \mid d_i, \mathbf{x}_i] = 0$$
$$d_i = m_0(\mathbf{x}_i) + u_i, \quad \mathbb{E}[u_i \mid \mathbf{x}_i] = 0$$

Steps:

- Divide the data into two folds;
- On fold 1, estimate  $\hat{g}_0(\mathbf{x}_i)$  and  $\hat{m}_0(\mathbf{x}_i)$  using ML methods;
- On fold 2, compute residuals:

$$\hat{\varepsilon}_i = y_i - \hat{g}_0(\mathbf{x}_i)$$

$$\hat{u}_i = d_i - \hat{m}_0(\mathbf{x}_i)$$

- Regress  $\hat{\varepsilon}_i$  on  $\hat{u}_i$  to get  $\hat{\theta}_0$ ;
- Repeat switching folds and average  $\hat{\theta}_0$ 's;
- In practice you can use  $K$  folds!
- See [Chernozhukov et al. \(2017\)](#) for a practical guide!

# Where do we go now?

Some open problems:

- Weak identification, in special in the IV context (see [Scheidegger et al. \(2025\)](#));
- Time series  $\implies$  it's impossible to do cross-fitting (see [Lewis and Syrgkanis \(2021\)](#));
- Panel data  $\implies$  usual estimators leverage linearity (see [Chernozhukov et al. \(2021\)](#) and [Clarke and Polselli \(2025\)](#));

# Where do we go now?

Some open problems:

- Weak identification, in special in the IV context (see [Scheidegger et al. \(2025\)](#));
- Time series  $\implies$  it's impossible to do cross-fitting (see [Lewis and Syrgkanis \(2021\)](#));
- Panel data  $\implies$  usual estimators leverage linearity (see [Chernozhukov et al. \(2021\)](#) and [Clarke and Polselli \(2025\)](#));

Good news! Plenty of dissertation topics!

**Questions?**

Thank you!  
See you tomorrow, stay tuned!



# Appendix and References

# References



Athey, Susan and Guido W. Imbens (Aug. 2019). "Machine Learning Methods That Economists Should Know About". en. In: *Annual Review of Economics* 11.1, pp. 685–725. ISSN: 1941-1383, 1941-1391. DOI: [10.1146/annurev-economics-080217-053433](https://doi.org/10.1146/annurev-economics-080217-053433). URL: <https://www.annualreviews.org/doi/10.1146/annurev-economics-080217-053433> (visited on 12/03/2025).



Belloni, A., V. Chernozhukov, and C. Hansen (Apr. 2014a). "Inference on Treatment Effects after Selection among High-Dimensional Controls". en. In: *The Review of Economic Studies* 81.2, pp. 608–650. ISSN: 0034-6527, 1467-937X. DOI: [10.1093/restud/rdt044](https://doi.org/10.1093/restud/rdt044). URL: <https://academic.oup.com/restud/article-lookup/doi/10.1093/restud/rdt044> (visited on 12/03/2025).



Belloni, Alexandre, Victor Chernozhukov, Iván Fernandez-Val, and Christian Hansen (2017). "Program Evaluation and Causal Inference With High-Dimensional Data". en. In: *Econometrica* 85.1, pp. 233–298. ISSN: 0012-9682. DOI: [10.3982/ECTA12723](https://doi.org/10.3982/ECTA12723). URL: <https://www.econometricsociety.org/doi/10.3982/ECTA12723> (visited on 12/04/2025).



Belloni, Alexandre, Victor Chernozhukov, and Christian Hansen (May 2014b). "High-Dimensional Methods and Inference on Structural and Treatment Effects". en. In: *Journal of Economic Perspectives* 28.2, pp. 29–50. ISSN: 0895-3309. DOI: [10.1257/jep.28.2.29](https://doi.org/10.1257/jep.28.2.29). URL: <https://pubs.aeaweb.org/doi/10.1257/jep.28.2.29> (visited on 12/03/2025).



Chernozhukov, Victor, Denis Chetverikov, Mert Demirer, Esther Duflo, Christian Hansen, and Whitney Newey (May 2017). "Double/Debiased/Neyman Machine Learning of Treatment Effects". en. In: *American Economic Review* 107.5, pp. 261–265. ISSN: 0002-8282. DOI: [10.1257/aer.p20171038](https://doi.org/10.1257/aer.p20171038). URL: <https://pubs.aeaweb.org/doi/10.1257/aer.p20171038> (visited on 12/02/2025).



Chernozhukov, Victor, Denis Chetverikov, Mert Demirer, Esther Duflo, Christian Hansen, Whitney Newey, and James Robins (Feb. 2018). "Double/debiased machine learning for treatment and structural parameters". en. In: *The Econometrics Journal* 21.1, pp. C1–C68. ISSN: 1368-4221, 1368-423X. DOI: [10.1111/ectj.12097](https://doi.org/10.1111/ectj.12097). URL: <https://academic.oup.com/ectj/article/21/1/C1/5056401> (visited on 12/03/2025).



Chernozhukov, Victor, Wolfgang Karl Härdle, Chen Huang, and Weining Wang (June 2021). "LASSO-driven inference in time and space". In: *The Annals of Statistics* 49.3. ISSN: 0090-5364. DOI: [10.1214/20-aos2019](https://doi.org/10.1214/20-aos2019). URL: <http://dx.doi.org/10.1214/20-AOS2019>.



Clarke, Paul S and Annalivia Polselli (Apr. 2025). "Double machine learning for static panel models with fixed effects". In: *Econometrics Journal*. ISSN: 1368-423X. DOI: [10.1093/ectj/utaf011](https://doi.org/10.1093/ectj/utaf011). URL: <http://dx.doi.org/10.1093/ectj/utaf011>.

