

# Intensional infect proportion of susceptible

April 20, 2018

Here is the system we are investigating:

$$\frac{dS}{d\tau} = \epsilon - \mathcal{R}_0 S(I_N + I_T) - \frac{p}{\gamma + \mu} S - \epsilon S \quad (1)$$

$$\frac{dI_T}{d\tau} = \frac{p}{\gamma + \mu} S + \mathcal{R}_0 S I_T - I_T \quad (2)$$

$$\frac{dI_N}{d\tau} = \mathcal{R}_0 S I_N - I_N \quad (3)$$

$$\frac{dR}{d\tau} = (1 - \epsilon)(I_N + I_T) - \epsilon R \quad (4)$$

We are trying to find the endemic equilibrium. Specifically the  $I_N$  value at E.E. By letting (3) equal to 0, we should get  $S = \frac{1}{\mathcal{R}_0}$ , but when substituting into (2), we get  $\frac{p}{\gamma + \mu} S = 0$ , which is inconsistent with our simulation result. Result shows that all variables stabilizes when  $t$  is large enough.

Thus, with the help of our plot. Thus, it is possible to conclude that  $S = \frac{1}{\mathcal{R}_0}$  is actually not the value of value of  $S$  at E.E.. Instead,  $I_N = 0$  is actually the E.E., for all  $p \neq 0$ .