

Intentional infection as a method of population level disease control

Newborn infection

Roger Zhang

McMaster University Department of Mathematics and Statistics

June 18th 2018

Motivation

- What are the historical examples?

Motivation

- What are the historical examples?
 - ▶ Variolation with smallpox

Motivation

- What are the historical examples?
 - ▶ Variolation with smallpox
 - ★ use scabs from infected person to infect someone else via skin

Motivation

- What are the historical examples?
 - ▶ Variolation with smallpox
 - ★ use scabs from infected person to infect someone else via skin
 - ★ variolation led to death less often than natural infection.

Motivation

- What are the historical examples?
 - ▶ Variolation with smallpox
 - ★ use scabs from infected person to infect someone else via skin
 - ★ variolation led to death less often than natural infection.
 - ▶ Pox parties

Motivation

- What are the historical examples?
 - ▶ Variolation with smallpox
 - ★ use scabs from infected person to infect someone else via skin
 - ★ variolation led to death less often than natural infection.
 - ▶ Pox parties
 - ★ increase contact rate to cause more natural infections

Motivation

- What are the historical examples?
 - ▶ Variolation with smallpox
 - ★ use scabs from infected person to infect someone else via skin
 - ★ variolation led to death less often than natural infection.
 - ▶ Pox parties
 - ★ increase contact rate to cause more natural infections
- Now illegal or uncommon – why do we care?

Motivation

- What are the historical examples?
 - ▶ Variolation with smallpox
 - ★ use scabs from infected person to infect someone else via skin
 - ★ variolation led to death less often than natural infection.
 - ▶ Pox parties
 - ★ increase contact rate to cause more natural infections
- Now illegal or uncommon – why do we care?
 - ▶ In history, the mechanisms and benefits of intentional infection on a population level was not quite understood.

Motivation

- What are the historical examples?
 - ▶ Variolation with smallpox
 - ★ use scabs from infected person to infect someone else via skin
 - ★ variolation led to death less often than natural infection.
 - ▶ Pox parties
 - ★ increase contact rate to cause more natural infections
- Now illegal or uncommon – why do we care?
 - ▶ In history, the mechanisms and benefits of intentional infection on a population level was not quite understood.
 - ▶ New application to disease control: transmissible vaccines

Method

Standard SIR model:

Method

Standard SIR model:

$$\begin{aligned}\frac{dS}{dt} &= \mu - \beta SI - \mu S, \\ \frac{dI}{dt} &= \beta SI - \gamma I - \mu I, \\ \frac{dR}{dt} &= \gamma I - \mu R,\end{aligned}\tag{1}$$

where S , I and R represent susceptible, infected and recovered.

- μ is birth/death rate (total population size N is constant)
- β is transmission rate
- γ is recovery rate

Assumptions:

- No difference between intentionally infected and naturally infected individuals.
- No disease induced mortality (in first version of model)
- Latent period (time from infection to becoming infectious) is short enough to be ignored.
- All susceptible individuals are equally likely to be infected, and all infected individuals are equally infectious.

Method

With intentional infection of newborns,

$$\begin{aligned}\frac{dS}{dt} &= \mu(1 - p) - \beta SI - \mu S, \\ \frac{dI}{dt} &= \beta SI + \mu p - \gamma I - \mu I, \\ \frac{dR}{dt} &= \gamma I - \mu R,\end{aligned}\tag{2}$$

where

p = proportion of newborns intentionally infected

Non-dimensionalize the above system by scaling time, by

$$\tau = (\gamma + \mu)t, \quad (3)$$

so time unit is “mean time infected”.

Non-dimensionalize the above system by scaling time, by

$$\tau = (\gamma + \mu)t, \quad (3)$$

so time unit is “mean time infected”.

Dimensionless system is:

$$\frac{dS}{d\tau} = \epsilon(1 - p) - \mathcal{R}_0 SI - \epsilon S, \quad (4a)$$

$$\frac{dI}{d\tau} = \mathcal{R}_0 SI + \epsilon p - I, \quad (4b)$$

where

$$\epsilon = \frac{\mu}{\gamma + \mu} \quad (\text{infectious period as proportion of lifetime}), \quad (5a)$$

$$\mathcal{R}_0 = \frac{\beta}{\gamma + \mu} \quad (\text{basic reproduction number}). \quad (5b)$$

Equilibria

Solving equations above to find equilibria, we obtain only one solution,

$$\hat{S} = \frac{1}{\mathcal{R}_0} - \frac{2p}{(\mathcal{R}_0 - 1) + \sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}, \quad (6a)$$

$$\hat{I} = \frac{\epsilon(\mathcal{R}_0 - 1) + \epsilon\sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}{2\mathcal{R}_0}. \quad (6b)$$

Equilibria

Solving equations above to find equilibria, we obtain only one solution,

$$\hat{S} = \frac{1}{\mathcal{R}_0} - \frac{2p}{(\mathcal{R}_0 - 1) + \sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}, \quad (6a)$$

$$\hat{I} = \frac{\epsilon(\mathcal{R}_0 - 1) + \epsilon\sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}{2\mathcal{R}_0}. \quad (6b)$$

- $\hat{I} \neq 0$ for all p between 0 and 1.

Equilibria

Solving equations above to find equilibria, we obtain only one solution,

$$\hat{S} = \frac{1}{\mathcal{R}_0} - \frac{2p}{(\mathcal{R}_0 - 1) + \sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}, \quad (6a)$$

$$\hat{I} = \frac{\epsilon(\mathcal{R}_0 - 1) + \epsilon\sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}{2\mathcal{R}_0}. \quad (6b)$$

- $\hat{I} \neq 0$ for all p between 0 and 1.
- This equilibrium is an **Endemic Equilibrium (EE)**.

Equilibria

Solving equations above to find equilibria, we obtain only one solution,

$$\hat{S} = \frac{1}{\mathcal{R}_0} - \frac{2p}{(\mathcal{R}_0 - 1) + \sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}, \quad (6a)$$

$$\hat{I} = \frac{\epsilon(\mathcal{R}_0 - 1) + \epsilon\sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}{2\mathcal{R}_0}. \quad (6b)$$

- $\hat{I} \neq 0$ for all p between 0 and 1.
- This equilibrium is an **Endemic Equilibrium (EE)**.
- Without intentional infection ($p = 0$) EE reduces to

$$\hat{S} = \frac{1}{\mathcal{R}_0}, \quad \hat{I} = \epsilon\left(1 - \frac{1}{\mathcal{R}_0}\right). \quad (7)$$

Stability of Equilibria

Jacobian Matrix,

$$\mathcal{J} = \begin{bmatrix} -\mathcal{R}_0 I - \epsilon & -\mathcal{R}_0 S \\ \mathcal{R}_0 I & \mathcal{R}_0 S - 1 \end{bmatrix} . \quad (8)$$

What else do we need?

What else do we need?

- How can we compare the results of this model to others and determine advantages or disadvantages of intentional infection?

What else do we need?

- How can we compare the results of this model to others and determine advantages or disadvantages of intentional infection?
- Define “advantage” as “fewer deaths”.

What else do we need?

- How can we compare the results of this model to others and determine advantages or disadvantages of intentional infection?
- Define “advantage” as “fewer deaths”.
- If intentionally infected cases have the same case fatality proportion (CFP) as naturally infected cases, then it will result in *more* deaths.
 - ▶ But variolation had lower CFP

What else do we need?

- How can we compare the results of this model to others and determine advantages or disadvantages of intentional infection?
- Define “advantage” as “fewer deaths”.
- If intentionally infected cases have the same case fatality proportion (CFP) as naturally infected cases, then it will result in *more* deaths.
 - ▶ But variolation had lower CFP \implies revise model to include different CFP for intentional infections.

What else do we need?

- How can we compare the results of this model to others and determine advantages or disadvantages of intentional infection?
- Define “advantage” as “fewer deaths”.
- If intentionally infected cases have the same case fatality proportion (CFP) as naturally infected cases, then it will result in *more* deaths.
 - ▶ But variolation had lower CFP \implies revise model to include different CFP for intentional infections.
- We need to divide I into two separate infective classes. V for intentionally infected class, I for naturally infected class.

Model with disease induced mortality rate

Therefore, our model becomes,

$$\frac{dS}{d\tau} = \epsilon(1 - p) - \mathcal{R}_0 S(V + I) - \epsilon S, \quad (9a)$$

$$\frac{dV}{d\tau} = \mathcal{R}_0 S V + \epsilon p - V, \quad (9b)$$

$$\frac{dI}{d\tau} = \mathcal{R}_0 S I - I, \quad (9c)$$

$$\frac{dM}{d\tau} = p_V(1 - \epsilon)V + p_I(1 - \epsilon)I, \quad (9d)$$

$$\frac{dR}{d\tau} = (1 - p_V)(1 - \epsilon)V + (1 - p_I)(1 - \epsilon)I - \epsilon R, \quad (9e)$$

where p_V and p_I represent the case fatality proportion (CFP) for intentionally infected and naturally infected cases, respectively.

Equilibria

If $p \neq 0$, the only equilibrium is,

$$\hat{S} = \frac{1}{\mathcal{R}_0} - \frac{2p}{(\mathcal{R}_0 - 1) + \sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}, \quad (10a)$$

$$\hat{V} = \frac{\epsilon(\mathcal{R}_0 - 1) + \epsilon\sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}{2\mathcal{R}_0}, \quad (10b)$$

$$\hat{I} = 0. \quad (10c)$$

- No naturally infected cases at endemic equilibrium (EE).
 - ▶ Helpful for disease eradication.

Effect of intentional infection on total mortality

Smallpox

Table: Model parameters and smallpox values.

Symbol	Meaning	Value
$\frac{1}{\mu}$	Average lifespan	50 years
$\frac{1}{\gamma}$	Mean infectious period	22 days
\mathcal{R}_0	Basic reproduction number	4.5
p_V	Intentionally infected CFP	0.01
p_I	Naturally infected CFP	0.3

Effect of intentional infection on total mortality

Mortality rate at EE

$$\left. \frac{dM}{d\tau} \right|_{EE} = \frac{p_V(1 - \epsilon)\epsilon(\mathcal{R}_0 - 1) + p_V(1 - \epsilon)\epsilon\sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0\rho}}{2\mathcal{R}_0}, \quad (11)$$

Effect of intentional infection on total mortality

Mortality rate at EE

$$\left. \frac{dM}{d\tau} \right|_{EE} = \frac{p_V(1 - \epsilon)\epsilon(\mathcal{R}_0 - 1) + p_V(1 - \epsilon)\epsilon\sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}{2\mathcal{R}_0}, \quad (11)$$

- Mortality rate at EE increases as proportion intentionally infected (p) increases.

Effect of intentional infection on total mortality

Mortality rate at EE

$$\left. \frac{dM}{d\tau} \right|_{EE} = \frac{p_V(1 - \epsilon)\epsilon(\mathcal{R}_0 - 1) + p_V(1 - \epsilon)\epsilon\sqrt{(\mathcal{R}_0 - 1)^2 + 4\mathcal{R}_0 p}}{2\mathcal{R}_0}, \quad (11)$$

- Mortality rate at EE increases as proportion intentionally infected (p) increases.
- In the long run, a larger p will lead to more deaths.

Initial state near equilibrium

In history, intentional infection was introduced when the population was endemic, so near the equilibrium for $p = 0$.

Initial state near equilibrium

In history, intentional infection was introduced when the population was endemic, so near the equilibrium for $p = 0$.

$$\hat{S} = \frac{1}{\mathcal{R}_0}, \quad (12a)$$

$$\hat{V} = 0, \quad (12b)$$

$$\hat{I} = \epsilon(1 - \frac{1}{\mathcal{R}_0}) \quad (12c)$$

Initial state being at equilibrium

- What is the time it takes to reach the new EE?
 - ▶ Need to define a threshold.

Initial state being at equilibrium

- What is the time it takes to reach the new EE?
 - ▶ Need to define a threshold.

Since the new equilibrium has $\hat{I} = 0$, define reaching equilibrium by $I \leq 10^{-6}$.




Figure: Determination of time taken to reach equilibrium




Figure: Time to advantage, as a function of p

With a lower proportion of intentional infection, we can gain advantages relatively faster.

Possibility of disease eradication

$\hat{I} = 0$ at EE. We can consider the Naturally infected cases already been eradicated at EE.

Possibility of disease eradication

$\hat{I} = 0$ at EE. We can consider the Naturally infected cases already been eradicated at EE.

For intentionally infected cases, if we stop intentional infection after we reach EE, can they burn out?

Possibility of disease eradication

Vaccination threshold (Herd immunity): $p_c = 1 - \frac{1}{\mathcal{R}_0}$

Possibility of disease eradication

Vaccination threshold (Herd immunity): $p_c = 1 - \frac{1}{\mathcal{R}_0}$

Threshold for susceptible: $S = \frac{1}{\mathcal{R}_0}$

Possibility of disease eradication

Vaccination threshold (Herd immunity): $p_c = 1 - \frac{1}{\mathcal{R}_0}$

Threshold for susceptible: $S = \frac{1}{\mathcal{R}_0}$

If S remains below this threshold until V goes extinct, then we can achieve complete eradication of this disease.

Possibility of disease eradication

Example: $p = 1$ initially, stop intentional infection after reaching EE,

Increase of S after we stop intentional infection




Figure: For more than 10 years after we stop intentional infection, $S < \frac{1}{R_0}$

Possibility of disease eradication

V as a function of time, after we stop intentional infection




Figure: It takes less than 1 year for V to fall below 1×10^{-6}

Future directions

- Model extension

Future directions

- Model extension
 - ▶ Different transmission rate for intentionally or naturally infected cases

Future directions

- Model extension
 - ▶ Different transmission rate for intentionally or naturally infected cases
 - ▶ Different recovery rate

Future directions

- Model extension
 - ▶ Different transmission rate for intentionally or naturally infected cases
 - ▶ Different recovery rate
- Combination of strategies

Future directions

- Model extension
 - ▶ Different transmission rate for intentionally or naturally infected cases
 - ▶ Different recovery rate
- Combination of strategies
- Comparison to traditional vaccination

Future directions

- Model extension
 - ▶ Different transmission rate for intentionally or naturally infected cases
 - ▶ Different recovery rate
- Combination of strategies
- Comparison to traditional vaccination
- Other approaches for intentional infection

Future directions

- Model extension
 - ▶ Different transmission rate for intentionally or naturally infected cases
 - ▶ Different recovery rate
- Combination of strategies
- Comparison to traditional vaccination
- Other approaches for intentional infection
- Use intentional infection models to improve understanding of historical smallpox dynamics

Other strategies for intentional infection

Another approach would be to intentionally infect susceptible individuals, with a certain rate r .

Other strategies for intentional infection

Another approach would be to intentionally infect susceptible individuals, with a certain rate r .

$$\begin{aligned}\frac{dS}{dt} &= \mu - \beta SI - rS - \mu S, \\ \frac{dI}{dt} &= rS + \beta SI - \gamma I - \mu I, \\ \frac{dR}{dt} &= \gamma I - \mu R,\end{aligned}\tag{13}$$



Challenges to this method

- Identify susceptible individuals

Conclusion

- Intentional infection has positive effects on population level disease control.
- Further study is required for determining intentional infection strategies to optimize this method.

References

-  Pauline van den Driessche *Reproduction numbers of infectious disease models*, Infectious Disease Modelling, Volume 2, Issue 3, August 2017, Pages 288-303.
-  R.M. Anderson, R.M. May *Infectious diseases of humans: Dynamics and control*, Oxford University Press, Oxford, UK (1991)