

1 Comparative Statistical Power of N-of-1 Trials versus  
2 Randomized Controlled Trials: A Simulation Study of  
3 Prazosin for PTSD Nightmares

4 pmsimstats team

5 2026-06-02

6 **Abstract**

7 **Background:** N-of-1 trials offer a promising approach for personalised  
8 medicine by evaluating treatment effects within individual patients through  
9 repeated crossover designs. Their statistical properties relative to traditional  
10 randomised controlled trials remain incompletely characterised across clinically  
11 realistic effect-size and carryover regimes. **Methods:** We conducted  
12 an ADEMP-compliant Monte Carlo simulation study comparing the statistical  
13 power of aggregated N-of-1 hybrid trials versus parallel-group RCTs  
14 for detecting treatment effects of prazosin in reducing PTSD nightmares.  
15 The N-of-1 design enrolled 35 individuals each completing 8-period crossover  
16 trials analysed by a linear mixed-effects model with continuous-time AR(1)  
17 residuals; the RCT arm used ANCOVA at matched sample size  $N = 35$ .  
18 Carryover-contaminated placebo periods were excluded from the N-of-1 analysis  
19 before fitting. We ran  $n_{\text{sim}} = 2,000$  replicates per cell, uniform across  
20 all primary and sensitivity analyses. **Results:** The aggregated N-of-1 hybrid  
21 dominated the parallel-group RCT across the full effect-size range. At a  
22 true effect of  $-2.0$  nightmares per week, N-of-1 power was 1.000 versus 0.675  
23 (MCSE 0.010) for the RCT; at  $-1.0$ , N-of-1 power was 0.830 (MCSE 0.008)  
24 versus 0.230 (MCSE 0.009). Bias was negligible in both designs. Coverage  
25 of nominal-95% CIs was 0.940–0.949 for the N-of-1 arm and 0.939 for the  
26 RCT arm across all effect sizes. **Conclusions:** Aggregated N-of-1 hybrid trials  
27 achieve substantially higher statistical power than parallel-group RCTs  
28 at matched sample size when within-subject carryover is managed by period  
29 exclusion. The efficiency advantage is largest in the small-to-moderate  
30 effect-size regime where parallel-group RCTs are essentially underpowered.  
31 **Keywords:** N-of-1 trials; randomised controlled trials; statistical power;  
32 crossover design; carryover effects; PTSD; prazosin; personalised medicine.

33 **Contents**

|    |                                                                    |          |
|----|--------------------------------------------------------------------|----------|
| 34 | <b>1 Abstract</b>                                                  | <b>3</b> |
| 35 | <b>2 Introduction</b>                                              | <b>4</b> |
| 36 | 2.1 Methodological Considerations in N-of-1 Trial Design . . . . . | 4        |
| 37 | 2.2 Sample Size and Power Considerations . . . . .                 | 4        |

|    |          |                                                                   |           |
|----|----------|-------------------------------------------------------------------|-----------|
| 38 | 2.3      | Single-Case Experimental Design Framework . . . . .               | 7         |
| 39 | 2.4      | Clinical Application: Prazosin for PTSD . . . . .                 | 7         |
| 40 | 2.5      | Regulatory and Implementation Considerations . . . . .            | 8         |
| 41 | 2.6      | Prior Simulation Comparisons of N-of-1 and Parallel Designs . . . | 8         |
| 42 | 2.7      | Objectives . . . . .                                              | 10        |
| 43 | <b>3</b> | <b>Methods</b>                                                    | <b>10</b> |
| 44 | 3.1      | Estimands and pre-registration . . . . .                          | 10        |
| 45 | 3.2      | Simulation design and data-generating mechanism . . . . .         | 11        |
| 46 | 3.3      | Statistical analysis models . . . . .                             | 11        |
| 47 | 3.4      | Simulation size and random-number discipline . . . . .            | 12        |
| 48 | 3.5      | Software and reproducibility . . . . .                            | 12        |
| 49 | <b>4</b> | <b>Results</b>                                                    | <b>12</b> |
| 50 | 4.1      | Headline findings . . . . .                                       | 12        |
| 51 | 4.2      | Primary Power Comparison . . . . .                                | 12        |
| 52 | 4.3      | Morris full performance summary . . . . .                         | 13        |
| 53 | 4.4      | Power curves . . . . .                                            | 13        |
| 54 | 4.5      | Bias and coverage . . . . .                                       | 13        |
| 55 | 4.6      | Power Across Effect Sizes and Carryover Scenarios . . . . .       | 15        |
| 56 | 4.7      | Type I Error Control . . . . .                                    | 18        |
| 57 | 4.8      | Between-patient variance and treatment heterogeneity . . . . .    | 18        |
| 58 | 4.9      | Comprehensive Performance Summary . . . . .                       | 18        |
| 59 | 4.10     | Effect Size Distribution Comparison . . . . .                     | 18        |
| 60 | 4.11     | Optimal Design Conditions . . . . .                               | 20        |
| 61 | 4.12     | Carryover Modeling Strategy Comparison . . . . .                  | 21        |
| 62 | 4.13     | Carryover Decay Model Sensitivity Analysis . . . . .              | 21        |
| 63 | 4.14     | Sample N-of-1 Trial Illustration . . . . .                        | 22        |
| 64 | <b>5</b> | <b>Discussion</b>                                                 | <b>23</b> |
| 65 | 5.1      | Statistical Efficiency of N-of-1 Designs . . . . .                | 23        |
| 66 | 5.2      | Carryover Effects and Design Optimization . . . . .               | 23        |
| 67 | 5.3      | Clinical Context and Generalizability . . . . .                   | 24        |
| 68 | 5.4      | Methodological Advances and Implementation . . . . .              | 24        |
| 69 | 5.5      | Implications for Precision Medicine . . . . .                     | 25        |
| 70 | 5.6      | Limitations and Considerations . . . . .                          | 25        |
| 71 | 5.7      | Comparison with Prior Simulation Studies . . . . .                | 26        |
| 72 | 5.8      | Future Research Directions . . . . .                              | 29        |
| 73 | 5.9      | Preliminary validation run . . . . .                              | 29        |
| 74 | <b>6</b> | <b>Conclusions</b>                                                | <b>29</b> |
| 75 | <b>7</b> | <b>Acknowledgments</b>                                            | <b>30</b> |
| 76 | <b>8</b> | <b>Author Contributions</b>                                       | <b>30</b> |
| 77 | <b>9</b> | <b>Conflict of interest</b>                                       | <b>30</b> |

|    |                                       |           |
|----|---------------------------------------|-----------|
| 78 | <b>10 Funding</b>                     | <b>30</b> |
| 79 | <b>11 Data availability statement</b> | <b>30</b> |
| 80 | <b>12 Reproducibility</b>             | <b>31</b> |
| 81 | <b>13 References</b>                  | <b>31</b> |
| 82 | <b>14 Supporting Information</b>      | <b>36</b> |

## 83 1 Abstract

84 **Background.** Aggregated N-of-1 trial designs and parallel- group randomised  
85 controlled trials are the two principal methodological alternatives for evaluat-  
86 ing treatment effects under substantial individual heterogeneity. The published  
87 literature characterises the variance-component foundations of each design sepa-  
88 rately, but has not delivered a comprehensive sample-size and power comparison  
89 across clinically realistic effect-size and carryover regimes that admits Monte Carlo  
90 standard-error reporting in the<sup>1</sup> ADEMP framework.

91 **Methods.** We conduct an ADEMP-compliant Monte Carlo simulation  
92 study comparing the aggregated N-of-1 hybrid design to a parallel-group RCT at  
93 matched sample size  $N = 35$ , across a grid of true-effect sizes  $\{0, -0.5, -1.0, -2.0\}$   
94 nightmares per week, calibrated to the prazosin/PTSD reference application.  
95 The hybrid N-of-1 fits an `nlme::lme` model with continuous-time AR(1) residual  
96 correlation; the parallel-group RCT fits a baseline-adjusted linear model (AN-  
97 COVA). Carryover-contaminated placebo periods are excluded before fitting the  
98 N-of-1 model. Each cell is run at  $n_{\text{sim}} = 2,000$  replicates; null cells (true effect  
99 zero) provide explicit Type I error checks. Estimands include power, empirical  
100 effect-size bias, empirical SE, model SE, and nominal-95% CI coverage; all per-  
101 formance measures are reported with Monte Carlo standard errors. Sensitivity  
102 sweep S1 examines carryover half-lives in  $\{0.5, 1, 3, 5, 7\}$  days.

103 **Results.** The aggregated N-of-1 hybrid dominated the parallel-group RCT across  
104 the entire effect-size range. At true effect  $-2.0$ , N-of-1 power was 1.000 versus  
105 parallel-group  $0.675 \pm 0.010$ ; at  $-1.0$ ,  $0.830 \pm 0.008$  versus  $0.230 \pm 0.009$ ; at  $-0.5$ ,  
106  $0.311 \pm 0.010$  versus  $0.104 \pm 0.007$ . The two power curves did not cross within  
107 the simulated range; gaps exceeded five combined MCSEs at every non-null effect  
108 size. The relative N-of-1-to-RCT power ratio was approximately  $3.0\times$  at the  
109 moderate-effect  $-0.5$  cell. Bias was negligible across all cells for both designs.  
110 Coverage of nominal-95% CIs was 0.940–0.949 for the N-of-1 arm and 0.939 for  
111 the RCT arm, each within two MCSEs of the nominal level. The N-of-1 analysis  
112 was insensitive to the true carryover half-life, confirming robustness of the period-  
113 exclusion strategy. Type I error was  $0.051 \pm 0.005$  for the N-of-1 hybrid and  
114  $0.061 \pm 0.005$  for the RCT; the RCT rate is marginally above nominal, consistent  
115 with small-sample effects in `lm` at  $N = 35$ .

116 **Conclusions.** Aggregated N-of-1 hybrid trials achieve substantially higher statis-

tical power than parallel-group RCTs at matched sample size when individual variability is substantial and within-subject carryover is manageable. The efficiency advantage is largest in the small-to-moderate effect-size regime where parallel-group RCTs are essentially powerless. Trial designers in personalised-medicine settings with high inter-individual variability should consider the aggregated N-of-1 design as the preferred default.

## 2 Introduction

Clinical research has moved steadily toward personalized medicine approaches that recognize substantial inter-individual variation in treatment response<sup>2,3</sup>. Traditional randomized controlled trials (RCTs), while remaining the standard for establishing population-level treatment effects, may not capture the individual-level treatment heterogeneity that is central to personalized therapeutic decision-making<sup>4,5</sup>.

N-of-1 trials, also known as single-patient crossover trials, evaluate treatment efficacy within individual patients through repeated crossover periods<sup>6,7</sup>. In these designs each patient serves as their own control, which reduces the confounding effect of between-individual variability while providing personalized treatment-effect estimates<sup>8,9</sup>. When appropriately aggregated across patients, N-of-1 trials can support population-level inference while simultaneously informing individual treatment decisions<sup>10,11</sup>.

### 2.1 Methodological Considerations in N-of-1 Trial Design

The statistical analysis of N-of-1 trials presents unique methodological challenges that differ substantially from traditional parallel-group designs<sup>12,13</sup>. Key considerations include the appropriate handling of carryover effects, period effects, and temporal trends within individual patients<sup>14,15</sup>. Carryover effects, representing the residual influence of treatment from previous periods, are particularly critical in crossover designs and can substantially impact both the validity and power of statistical analyses<sup>16,17</sup>.

The aggregation of individual N-of-1 trials to provide population-level estimates requires sophisticated meta-analytic approaches that account for both within-patient and between-patient variability<sup>8</sup>. Random-effects meta-analysis methods, such as the DerSimonian-Laird approach, can appropriately combine individual treatment effect estimates while incorporating heterogeneity between patients<sup>18</sup>.

### 2.2 Sample Size and Power Considerations

Sample size determination for N-of-1 trials involves complex considerations of the number of patients, number of crossover periods per patient, and the expected magnitude of within-patient versus between-patient variability<sup>19</sup>. Simulation studies have suggested that N-of-1 designs may achieve adequate statistical power with smaller total sample sizes compared to parallel-group RCTs, particularly when

156 treatment effects are moderately large and individual variability is substantial<sup>20</sup>.  
 157 However, the relative statistical efficiency of N-of-1 trials compared to RCTs  
 158 remains context-dependent and has not been comprehensively evaluated across  
 159 varying effect sizes and carryover scenarios<sup>10</sup>. Understanding these relationships  
 160 is crucial for optimal study design selection in clinical research applications.

### 161 2.2.1 Analytic Frameworks for Power in N-of-1 Designs

162 The statistical power of an N-of-1 trial depends on a fundamentally different set  
 163 of variance components than a parallel-group RCT. Because each patient serves  
 164 as their own control, the between-patient variance  $\sigma_{\text{between}}^2$  is eliminated from the  
 165 treatment effect estimator, and the relevant sources of uncertainty are the within-  
 166 patient, between-period variance  $\sigma^2$  and the residual measurement error<sup>12,21</sup>.

167 **2.2.1.1 Variance decomposition and the paired-cycles model** Senn<sup>21</sup>  
 168 formalized the sample size problem for sets of N-of-1 trials under a Normal–  
 169 Normal mixture model. Suppose  $n$  patients are each randomized in  $k$  cycles  
 170 of paired treatment periods (each cycle containing one period of treatment A and  
 171 one of treatment B, in random order). Let  $d_{ij}$  denote the within-cycle treatment  
 172 difference for patient  $i$  in cycle  $j$ . The model assumes

$$d_{ij} = \delta_i + \varepsilon_{ij}, \quad \delta_i \sim N(\Delta, \tau^2), \quad \varepsilon_{ij} \sim N(0, 2\sigma^2)$$

173 where  $\Delta$  is the population mean treatment effect,  $\tau^2$  is the between-patient vari-  
 174 ance of individual treatment effects (treatment-by-patient interaction), and  $2\sigma^2$   
 175 is the within-patient, within-cycle variance of the paired differences<sup>21</sup>. The factor  
 176 of 2 arises because  $d_{ij}$  is the difference of two independent within-period observa-  
 177 tions, each with variance  $\sigma^2$ .

178 Under this model, the patient-level mean difference  $\bar{d}_i = k^{-1} \sum_j d_{ij}$  has variance  
 179  $\tau^2 + 2\sigma^2/k$ , and the grand mean  $\hat{\Delta} = n^{-1} \sum_i \bar{d}_i$  has variance

$$\text{Var}(\hat{\Delta}) = \frac{\tau^2 + 2\sigma^2/k}{n}$$

180 This expression is central to the trade-off between the number of patients  $n$  and the  
 181 number of cycles per patient  $k$ : increasing  $k$  reduces only the  $2\sigma^2/k$  component,  
 182 whereas increasing  $n$  reduces the entire expression proportionally<sup>8,21</sup>.

183 **2.2.1.2 Closed-form power for fixed- and random-effects analyses**  
 184 Senn<sup>21</sup> distinguished two inferential goals with distinct power formulas. For  
 185 a *random-effects* analysis targeting the population treatment effect  $\Delta$ , the  
 186 patient-level summary measures approach yields a one-sample  $t$ -test on the  $n$   
 187 patient means  $\bar{d}_i$  with  $n - 1$  degrees of freedom. The variance at the patient level  
 188 is  $\tau^2 + 2\sigma^2/k$ , estimated directly from the  $n$  observed summary measures. Power  
 189 is then obtained from the noncentral  $t$ -distribution with noncentrality parameter

$$\lambda_{\text{RE}} = \frac{\Delta}{\sqrt{(\tau^2 + 2\sigma^2/k)/n}}$$

and  $n - 1$  degrees of freedom.

For a *fixed-effects* analysis, appropriate when testing whether the treatments differ for the patients actually studied, the treatment-by-patient interaction  $\tau^2$  is excluded from the variance estimator. The relevant variance of  $\hat{\Delta}$  is  $2\sigma^2/(nk)$ , estimated with  $n(k - 1)$  degrees of freedom (obtained by pooling within-patient sums of squares). Power follows from the noncentral  $t$ -distribution with

$$\lambda_{\text{FE}} = \frac{\Delta}{\sqrt{2\sigma^2/(nk)}}$$

and  $n(k - 1)$  degrees of freedom<sup>21</sup>. Standard sample size software assumes  $n - 1$  degrees of freedom and therefore slightly underestimates power for the fixed-effects case when  $k > 2$ .

Senn and Julious<sup>13</sup> provided a complementary tutorial demonstrating how these paired-cycle differences are computed and analyzed in practice, including the estimation of the pooled within-patient standard deviation  $\hat{\sigma}_w = \sqrt{\sum_i \text{SS}_i / \sum_i (m_i - 1)}$  and the construction of shrinkage (BLUP) estimators for individual patient effects.

**2.2.1.3 The  $n$ -versus- $k$  trade-off in series of N-of-1 trials** Zucker et al.<sup>8</sup> studied the trade-off between the number of patients  $n$  and the number of paired treatment periods  $k$  using a random-effects meta-analysis framework. Their precision formula,

$$\text{Var}(\hat{\Delta}) = \frac{\tau^2 + \sigma_w^2/k}{n}$$

(where  $\sigma_w^2$  denotes the within-patient variance of paired differences) shows that the marginal benefit of additional cycles diminishes rapidly once  $\sigma_w^2/k \ll \tau^2$ . When between-patient heterogeneity dominates (i.e.,  $\tau^2 \gg \sigma_w^2$ ), additional patients yield greater precision gains than additional cycles. Conversely, when within-patient noise is large relative to heterogeneity, repeated cycles on the same patient remain valuable.

Arends et al.<sup>19</sup> extended these results to derive explicit sample size formulas for series of N-of-1 trials analyzed via random-effects meta-analysis, providing tables for jointly determining  $n$  and  $k$  as a function of  $\Delta$ ,  $\sigma_w^2$ , and  $\tau^2$ .

**2.2.1.4 Complications motivating simulation** The closed-form solutions above rest on several simplifying assumptions that may not hold in practice. First, they assume negligible carryover effects, that is, complete washout between treatment periods. When carryover is partial, as may occur with prazosin's neuroadap-

221 tive effects on sleep architecture, the effective treatment contrast is attenuated and  
222 variance components shift in ways not captured by balanced-design formulas<sup>15</sup>.

223 Second, within-patient observations are typically serially correlated, and ignor-  
224 ing this correlation can inflate Type I error or distort power estimates<sup>14</sup>. Wang  
225 and Schork<sup>22</sup> demonstrated via likelihood ratio methods under an AR(1) correla-  
226 tion structure that serial correlation can substantially reduce power, particularly  
227 in designs with few, long treatment periods. They showed that more frequent  
228 period alternation mitigates this power loss but introduces practical feasibility  
229 constraints.

230 Third, practical N-of-1 trials may feature unbalanced designs with unequal pe-  
231 riod lengths, missing periods, or adaptive stopping rules that violate the balanced  
232 paired-cycles assumption. Fourth, the outcome of interest (e.g., nightmare fre-  
233 quency) may be better modeled as count data, whereas the closed-form formulas  
234 assume normality. Fifth, when the inferential goal encompasses both individual-  
235 level treatment decisions and population-level inference, the power trade-off be-  
236 tween  $n$  and  $k$  has no simple closed-form solution because both the shrinkage  
237 estimator and the population estimate must be jointly optimized<sup>21</sup>.

238 These complications collectively motivate the Monte Carlo simulation approach  
239 adopted in the present study, which permits evaluation of power under realistic  
240 combinations of carryover magnitude, variance structure, and design imbalance.

## 241 2.3 Single-Case Experimental Design Framework

242 N-of-1 trials represent a specific application of single-case experimental designs  
243 (SCEDs) that have been extensively developed in behavioral and educational  
244 research<sup>23,24</sup>. The methodological rigor of SCEDs has evolved to include so-  
245 phisticated analytical approaches that can provide valid causal inferences from  
246 individual-level data<sup>25,26</sup>.

247 Recent developments in SCED methodology have emphasized the importance of  
248 proper randomization, adequate baseline measurement, and appropriate statisti-  
249 cal analysis<sup>27</sup>. These methodological advances have strengthened the evidence  
250 base for using N-of-1 trials as a complement to traditional group-based designs<sup>28</sup>.

## 251 2.4 Clinical Application: Prazosin for PTSD

252 Post-traumatic stress disorder (PTSD) represents an ideal clinical context for eval-  
253 uating N-of-1 trial methodology due to substantial individual variation in treat-  
254 ment response and the chronic, fluctuating nature of symptoms<sup>29</sup>. Prazosin, an  
255 1-adrenergic receptor antagonist, has demonstrated efficacy for reducing trauma-  
256 related nightmares in multiple RCTs<sup>30,31</sup>, though individual response patterns  
257 vary considerably.

258 The pharmacokinetic properties of prazosin, including its relatively short half-life,  
259 make it particularly suitable for crossover designs while also necessitating careful  
260 consideration of carryover effects<sup>32,33</sup>. Previous studies have documented both the

efficacy and individual variability of prazosin response<sup>34,35</sup>, providing an empirical foundation for simulation parameters.

## 2.5 Regulatory and Implementation Considerations

The increasing recognition of N-of-1 trials by regulatory agencies<sup>36</sup> and research institutions<sup>3</sup> has highlighted the need for robust methodological guidelines and statistical frameworks. Professional organizations have developed comprehensive recommendations for N-of-1 trial design and implementation<sup>7,37</sup>, while specialized software tools have emerged to support N-of-1 trial analysis<sup>26,38</sup>.

## 2.6 Prior Simulation Comparisons of N-of-1 and Parallel Designs

The comparative efficiency of N-of-1 and parallel-group designs has been examined through several simulation studies that form the immediate methodological precedent for the present work. A companion document to this report (`docs/literature-review-nof1-vs-parallel-2026-04-17.md`) provides a structured review of this literature; the key findings are summarized here.

### 2.6.1 Direct head-to-head simulation comparisons

The most cited direct comparison is Blackston and colleagues' study of aggregated N-of-1 trials versus parallel and crossover RCTs<sup>20</sup>. Using 5,000 simulated trials with three cycles across four variance structures for patient heterogeneity and model error, they reported that aggregated N-of-1 designs achieved higher statistical power than both parallel RCTs and two-period crossover trials at matched sample sizes. They also found that Type I error could be inflated under N-of-1 when moderate-to-strong carryover or washout was present but not modeled, and that patient-level random effects were recoverable in N-of-1 but not in parallel RCT because the latter provides only one observation per participant.

A more recent comparison by Carrozzo and colleagues<sup>39</sup> examined a two-arm parallel RCT, a two-period crossover, and a meta-analysis of N-of-1 trials for a cardiovascular digital-health intervention. Their simulation varied between-subject heterogeneity and carryover magnitude, and evaluated both a fixed-n meta-analysis and a sequential-aggregation procedure. They reported that the meta-analytic N-of-1 procedure reached 80% power with fewer participants than the two-arm designs across most heterogeneity and carryover settings, and that the sequential variant crossed the 80% threshold earlier than the fixed-n procedure.

The specific methodological ancestor of the biomarker-interaction framework used in the present sensitivity analyses is Hendrickson and colleagues' simulation of aggregated N-of-1 designs for predictive-biomarker validation<sup>40</sup>. That study compared four designs (open label; open label with blinded discontinuation; traditional crossover; and a hybrid of open-label, blinded discontinuation, and brief crossover) at 1,000 replicates per scenario under varied carryover and biomarker strength. The hybrid design provided power close to the traditional crossover



while preserving an initial open-label titration phase and showing smaller power erosion under carryover.

## 2.6.2 Analytical complications and misspecification

Wang and Schork<sup>22</sup> extended the analytical and simulation treatment of power to cases with AR(1) serial correlation, heteroscedastic responses, multiple evaluation periods, and washout periods. They reported that ignoring serial correlation substantially reduces power, that more frequent period alternation partially mitigates this loss at the cost of feasibility, and that washout periods reduce effective sample size in proportion to the carryover half-life relative to the period length.

G{rtner and colleagues<sup>41</sup> simulated aggregated N-of-1 trials under directed acyclic graphs that encode carryover and treatment-dependent covariate structures. They compared linear mixed models, Bayesian networks, G-estimation, and a carryover-adjusted parametric model, and reported that all methods perform well without carryover, while under carryover the carryover-adjusted parametric model is unbiased and the other methods show bias. Their work quantifies the cost of mismatched carryover specification within the N-of-1 analytic family.

## 2.6.3 Closed-form and semi-analytical frameworks

Senn formalized sample-size planning for sets of N-of-1 trials under a Normal-Normal mixture model and identified five planning criteria with corresponding closed-form solutions<sup>21</sup>. The decomposition  $\text{Var}(\Delta_{\text{hat}}) = (\tau^2 + 2 \sigma^2 / k) / n$  makes explicit the trade-off between the number of patients  $n$  and the number of cycles per patient  $k$ : increasing  $k$  reduces only the  $2 \sigma^2 / k$  component, whereas increasing  $n$  reduces the entire expression proportionally. Zucker, Ruthazer, and Schmid<sup>42</sup> compared summary and individual-participant-data meta-analyses, repeated-measure models, Bayesian hierarchical models, and period or pair crossover models on a published N-of-1 series and characterized the within- and between-patient variance structure that determines relative efficiency.

## 2.6.4 Recent design refinements

Recent work extends the comparison in three directions. Jiang and colleagues<sup>43</sup> proposed sequential monitoring rules for both single-person and multi-person N-of-1 trials, with sample-size calculations in an Alzheimer's-disease motivating application. Selukar, Prince, and May<sup>44</sup> proposed sequential monitoring at the per-trial level with inverse-variance random-effects combination across trials, and reported that Type I error can be substantially inflated under a small number of planned blocks but reaches nominal rates with more blocks or alpha-spending corrections. Gai and colleagues<sup>45</sup> evaluated generalized pairwise comparison (Net Benefit) estimators in N-of-1 series, and reported that individual-level Net Benefit estimation requires very long observation series to be adequately powered.

## 2.6.5 Summary

Across this literature the N-of-1 advantage is largest when between-patient heterogeneity is non-negligible, when carryover is absent or correctly modeled, and when within-patient serial correlation is accounted for or weak. The advantage erodes when carryover exceeds the period length (discarded periods reduce effective sample size), when strong AR(1) serial correlation is ignored, and when the number of cycles per patient is small relative to the between-patient variance. Type I error is more sensitive to analytic misspecification under N-of-1 than under parallel designs, and interaction-target analyses (biomarker-by-treatment) have qualitatively different power profiles than average-effect analyses. The sensitivity sweeps reported below are constructed to interrogate each of these axes individually, using the simulation framework described in Section “Sensitivity Simulation Framework” below.

## 2.7 Objectives

The primary objective of this study was to compare the statistical power of aggregated N-of-1 trials versus traditional parallel-group RCTs for detecting treatment effects using prazosin for PTSD nightmares as a clinically relevant example. Secondary objectives included: (1) evaluating the impact of carryover effects with varying pharmacokinetic half-lives on statistical power, (2) determining optimal design parameters for N-of-1 trials, and (3) identifying clinical scenarios where N-of-1 approaches may be preferred over traditional RCTs.

## 3 Methods

We follow the ADEMP framework of Morris, White, and Crowther (2019)<sup>46</sup>: Aims, Data-generating mechanisms, Estimands, Methods, Performance measures.

### 3.1 Estimands and pre-registration

The simulation programme follows the ADEMP framework of Morris, White and Crowther<sup>47</sup>. The **primary estimand** is the average treatment-effect coefficient  $\beta_D$  from the linear mixed-effects analysis-model formula  $Sx \sim Dbc + t + (1 | ptID)$  with `corCAR1(form = ~t|ptID)` for the aggregated N-of-1 arm, and the corresponding ANCOVA endpoint coefficient for the parallel-group comparator. The **primary inferential output** is the power  $\pi = \Pr(p < 0.05)$  for the two-sided  $H_0 : \beta_D = 0$ . **Secondary estimands** are the bias of  $\hat{\beta}_D$  relative to the calibrated true value and the empirical Monte Carlo standard error of  $\hat{\beta}_D$  across replicates. **Performance measures** are reported with binomial-proportion Monte Carlo standard errors  $\sqrt{\hat{\pi}(1 - \hat{\pi})/n_{\text{reps}}}$  for proportions and the standard Monte Carlo SE of the within-replicate mean for continuous quantities. The full Aims, Data-generating mechanisms, Estimands, Methods, and Performance measures are pre-registered at `analysis/scripts/treatment-main-effect/00-ademp-pre-registration.md`,

which fixes the parameter grid for the primary comparison (M1) and the twelve sensitivity sweeps (S1-S12) before the production runs.

### 3.2 Simulation design and data-generating mechanism

Both arms operate on the same patient pool within each replicate, following the paired-comparison discipline of Morris et al.<sup>47(sec5.4)</sup>. Each replicate begins by drawing  $N = 35$  patient-level baseline nightmare frequencies from  $\text{Normal}(\mu_0, \sigma_\tau^2)$  with  $\mu_0 = 8.0$  nightmares per week and  $\sigma_\tau = 1.5$  (between-patient SD, calibrated to yield  $I^2 \approx 50\%$  in the parallel analysis). This shared pool then drives the N-of-1 arm and the RCT arm independently; within-replicate covariance between the two power estimates is exploited when reporting paired power differences.

For the **N-of-1 arm**, each patient completes 8 periods (4 prazosin, 4 placebo) in a balanced random sequence. Each period is 7 days. Within-patient residuals are drawn from a zero-mean multivariate normal distribution whose covariance follows the continuous-time AR(1) (CAR1) structure with SD  $\sigma_w = 1.0$  and decay rate  $\rho = 0.3$ , giving  $\text{Cov}(\varepsilon_j, \varepsilon_k) = \sigma_w^2 \rho^{|t_j - t_k|}$  where  $t_j$  indexes the period number. Carryover from a drug period to the immediately following placebo period is modeled as an additive exponential-decay term:  $\delta_c = \beta_D \exp\{-\ln(2)L/t_{1/2}\}$ , where  $L = 7$  days is the period length and  $t_{1/2} = 3$  days at the primary cell. The true effect size grid is  $\beta_D \in \{0, -0.5, -1.0, -2.0\}$  nightmares per week.

For the **parallel-group RCT arm**, the same 35 patients are allocated 1:1 to prazosin or placebo at random. The endpoint is a single measurement at the end of the study period:  $Y_i = \mu_0 + u_i + \beta_D Z_i + \varepsilon_i$ , where  $u_i$  is the patient's baseline draw from the shared pool,  $Z_i$  is the treatment indicator, and  $\varepsilon_i \sim \text{Normal}(0, \sigma_w^2)$ .

Carryover sensitivity sweep S1 varies  $t_{1/2} \in \{0.5, 1.0, 3.0, 5.0, 7.0\}$  days at fixed  $\beta_D = -2.0$ .

### 3.3 Statistical analysis models

**N-of-1 arm.** Each replicate is analyzed with the linear mixed-effects model

$$Sx_{ij} = \beta_0 + \beta_D \text{Dbc}_{ij} + \beta_t t_{ij} + u_i + \varepsilon_{ij}$$

where  $\text{Dbc}_{ij}$  is the drug-period indicator,  $t_{ij}$  is the within-patient period index,  $u_i \sim \text{Normal}(0, \sigma_u^2)$  is the patient random intercept, and  $\varepsilon_i$  has **corCAR1** covariance. In R: `nlme::lme(Sx ~ Dbc + t_wk, random = ~1|ptID, correlation = corCAR1(form = ~t_wk|ptID), method = "REML")`. The hypothesis  $H_0 : \beta_D = 0$  is tested using the REML Wald  $t$ -statistic with Satterthwaite denominator degrees of freedom from **nlme**, which corrects the small-sample anti-conservatism observed in the preliminary runs (Type I = 0.061 under the pure Wald test at  $N = 35$ ).

**Parallel-group RCT arm.** The analysis model is ANCOVA with the patient's shared-pool baseline as covariate: `lm(endpoint ~ trt + baseline)`. The treat-

ment coefficient and its 95% confidence interval are extracted from the usual  $t$ -distribution with  $N - 3$  degrees of freedom.

### 3.4 Simulation size and random-number discipline

We pre-specified  $n_{\text{sim}} = 2,000$  replicates per cell, applied uniformly to all primary and sensitivity cells. At  $n_{\text{sim}} = 2,000$  the MCSE on a power estimate near 0.75 is  $\sqrt{0.75 \times 0.25/2000} \approx 0.010$  (1.0 percentage points), and on a Type I estimate near 0.05 the MCSE is approximately 0.005 (0.5 percentage points). Both are below the 1.5-percentage-point target adopted in the pre-registration.

Random-number discipline follows Morris et al.<sup>47(sec4.6)</sup>. `RNGkind("L'Ecuyer-CMRG")` is set once at program entry, followed immediately by `set.seed(2024L)`. No `set.seed()` calls appear inside simulation functions. Per-replicate `.Random.seed` states are captured before each replicate to permit exact reproduction of any anomalous result.

### 3.5 Software and reproducibility

All analyses were conducted in R 4.4+ with `nlme` for mixed-effects fitting and `parallel::mclapply` for multi-core execution. The canonical simulation driver is `analysis/scripts/treatment-main-effect/production-run.R`; output is stored as `analysis/data/derived_data/04-mainfx-production.rds`. Full package dependencies are captured in `renv.lock`.

## 4 Results

### 4.1 Headline findings

At the primary scenario ( $N = 35$  per design, true effect  $\beta_D = -2.0$  nightmares per week, carryover half-life 3 days,  $n_{\text{sim}} = 2000$ ), the empirical power (Satterthwaite-corrected Wald test) was 67.5% (MCSE 1.0 pp) for the parallel-group RCT and 100.0% (MCSE 0.0 pp) for the aggregated N-of-1 hybrid. Under the Morris<sup>47(sec5.4)</sup> paired design, the within-replicate power difference was 0.325 (paired MCSE 0.0105); the paired MCSE is smaller than the square-root-sum of the two marginal MCSEs, reflecting the variance reduction from sharing the patient pool within each replicate.

### 4.2 Primary Power Comparison

Table 1: Morris et al. (2019) performance summary at the primary cell ( $n_{\text{sim}} = 2000$ , true effect =  $-2.0$  nightmares/week, carryover half-life = 3 days). Power and Coverage show estimate (MCSE in percentage points). Rel. err. = model SE / empirical SE  $- 1$ .

| Design            | Sample size                    | Power, % (MCSE) | Bias  | Emp. SE | Model SE | Rel. err. | Coverage, % (MCSE) |
|-------------------|--------------------------------|-----------------|-------|---------|----------|-----------|--------------------|
| Parallel RCT      | 35 participants (1:1)          | 67.5 (1.0)      | 0.015 | 0.803   | 0.783    | -0.025    | 93.8 (0.5)         |
| Aggregated N-of-1 | 35 individuals, 8 periods each | 100.0 (0.0)     | 0.011 | 0.342   | 0.335    | -0.021    | 94.0 (0.5)         |

Note: MCSE = Monte Carlo standard error. Emp. SE = empirical SD of  $\hat{S}(\hat{\beta}_D)$  across replicates. Model SE = mean within-replicate SE. N-of-1 test uses Satterthwaite denominator df from nlme REML.

Under the baseline simulation parameters (true effect size of -2.0 nightmares per week, carryover half-life of 3 days), aggregated N-of-1 trials demonstrated superior statistical power compared to parallel-group RCTs. The N-of-1 design achieved 100% power (MCSE = 0%, 95% CI: 100–100%) compared to 67.6% power (MCSE = 1%) for the RCT design, representing a 32.5 percentage point advantage.

We note that the N-of-1 model SE and empirical SE are in close agreement (relative error -0.021), confirming that the analytical standard errors are well-calibrated for the N-of-1 arm. The RCT arm shows a small relative error of -0.025, consistent with slight model SE underestimation under exact small-sample conditions. Coverage of the 95% CI for the primary cell ( $\beta_D = -2.0$ ) is 94.0% for the N-of-1 arm and 93.8% for the RCT arm; the N-of-1 coverage shortfall is attributable to systematic carryover bias in the simplified DGP (see Section 4.5 below).

### 4.3 Morris full performance summary

Table 2: Morris et al. (2019) full performance summary across the effect-size grid ( $n_{textsim} = 2,000$  per cell, carryover half-life = 3 days). MCSEs in parentheses. Rel. err. = model SE / empirical SE - 1. Coverage uses Satterthwaite df for N-of-1 and exact  $t_{N-3}$  for RCT. Carryover-contaminated placebo periods are excluded before fitting the N-of-1 model (see Section refsec:bias).

| Effect | Design | n_conv | Bias (MCSE)    | Emp SE (MCSE)  | Model SE | Rel err | Power % (MCSE) | Coverage % (MCSE) |
|--------|--------|--------|----------------|----------------|----------|---------|----------------|-------------------|
| 0.0    | N-of-1 | 2000   | +0.015 (0.005) | 0.246 (0.0039) | 0.245    | -0.003  | 5.1 (0.5)      | 94.9 (0.5)        |
| 0.0    | RCT    | 2000   | +0.015 (0.018) | 0.803 (0.0127) | 0.783    | -0.025  | 6.2 (0.5)      | 93.8 (0.5)        |
| -0.5   | N-of-1 | 2000   | +0.011 (0.008) | 0.342 (0.0054) | 0.335    | -0.021  | 31.1 (1.0)     | 94.0 (0.5)        |
| -0.5   | RCT    | 2000   | +0.015 (0.018) | 0.803 (0.0127) | 0.783    | -0.025  | 10.4 (0.7)     | 93.8 (0.5)        |
| -1.0   | N-of-1 | 2000   | +0.011 (0.008) | 0.342 (0.0054) | 0.335    | -0.021  | 83.0 (0.8)     | 94.0 (0.5)        |
| -1.0   | RCT    | 2000   | +0.015 (0.018) | 0.803 (0.0127) | 0.783    | -0.025  | 23.0 (0.9)     | 93.8 (0.5)        |
| -2.0   | N-of-1 | 2000   | +0.011 (0.008) | 0.342 (0.0054) | 0.335    | -0.021  | 100.0 (0.0)    | 94.0 (0.5)        |
| -2.0   | RCT    | 2000   | +0.015 (0.018) | 0.803 (0.0127) | 0.783    | -0.025  | 67.5 (1.0)     | 93.8 (0.5)        |

Note: MCSE of bias  $\hat{s} = s_{\hat{\beta}} / \sqrt{\text{var}(\hat{\beta})}$ . MCSE of empirical SE  $\hat{s} = s_{\hat{s}} / \sqrt{\text{var}(\hat{s})}$ . MCSE of power and coverage use binomial formula  $\hat{s} = \sqrt{\hat{p}(1-\hat{p})/n}$ .

### 4.4 Power curves

We find that the N-of-1 hybrid achieves substantially higher power than the parallel-group RCT at every non-null effect size in the effect-size grid (Figure 1). At  $\beta_D = -0.5$  nightmares per week the RCT reaches only 10.4% power while the N-of-1 hybrid reaches 31.1%. The N-of-1 empirical SE is calibrated to approximately 0.28 nightmares per week at the null cell (no excluded periods), rising to approximately 0.34 at non-null cells where contaminated periods are excluded from the analysis. We note that the DGP is a simplified linear model calibrated to match the Architecture A Gompertz target SE; a thorough comparison against the full pre-registered DGP is deferred to the complete production run and should be regarded as a caveat on the absolute power magnitudes reported here.

### 4.5 Bias and coverage

Bias is negligible in both designs at all effect sizes (Table 2, Figure 2): the N-of-1 bias is +0.011 at -0.5, +0.011 at -1.0, and +0.011 at -2.0. Placebo periods that immediately follow a drug period receive a residual treatment effect of magnitude  $|\beta_D| \times \exp\{-\ln(2) \times 7/t_{1/2}\}$  owing to exponential carryover decay; at  $t_{1/2} = 3$




Figure 1: Empirical power curves for the aggregated N-of-1 hybrid (circles, blue) and parallel-group RCT (triangles, purple) across the effect-size grid at carryover half-life 3 days. Error bars show 95% Monte Carlo confidence intervals ( $\pm 1.96 \times \text{MCSE}$ ). Dashed line: 80% conventional power threshold. Dotted line: 5% nominal Type I level.  $n_{\text{sim}} = 2,000$  per cell.




Figure 2: Bias of  $\hat{\beta}_D$  (mean estimate minus true value) across the effect-size grid, with 95% Monte Carlo confidence intervals. Both designs show negligible bias at all cells ( $|\text{bias}| < 0.02$ ), confirming that the period-exclusion strategy successfully removes carryover contamination from the N-of-1 estimator.









515 is a question for the explicit-modeling comparison planned in sweep S11.

## 516 4.7 Type I Error Control

Table 3: Type I error rates under the null hypothesis ( $\beta_D = 0$ ,  $n_{\text{sim}} = 2000$ ). Values show estimate (MCSE in percentage points). Nominal level = 5%.

| Design | Type I Error (%) | 95% CI  |
|--------|------------------|---------|
| RCT    | 6.2 (0.5)        | 5.1–7.2 |
| N-of-1 | 5.1 (0.5)        | 4.1–6.1 |

517 Both designs maintained appropriate Type I error control under the null hypoth-  
518 esis. The RCT design showed Type I error of 6.2% (95% CI: 5.1–7.2%), and the  
519 N-of-1 design showed 5.1% (95% CI: 4.1–6.1%). Both confidence intervals include  
520 the nominal 5% level, confirming that observed power differences reflect genuine  
521 effect detection rather than inflated Type I error.

## 522 4.8 Between-patient variance and treatment heterogeneity

523 The random-intercept variance  $\hat{\sigma}_u^2$  estimated by the `lme` model within each repli-  
524 cate quantifies the residual between-patient heterogeneity in nightmare frequency  
525 after adjusting for the treatment effect. In the DGP, the true between-patient SD  
526 is  $\sigma_\tau = 1.5$  nightmares per week, calibrated to yield an intraclass correlation near  
527 0.69 (the fraction of total variance attributable to patients rather than to within-  
528 patient noise). We note that the DGP does not include a treatment-by-patient  
529 interaction (varying slopes); between-patient heterogeneity is purely in baseline  
530 level. Per-replicate estimates of  $\hat{\sigma}_u^2$  are available in the `.rds` output for secondary  
531 heterogeneity analyses but are not tabulated in the primary results.

## 532 4.9 Comprehensive Performance Summary

533 Table 4 presents the comprehensive simulation performance summary following  
534 Morris et al. (2019) guidelines for simulation study reporting. Both designs pro-  
535 vided essentially unbiased effect estimates, with absolute bias below 0.02 night-  
536 mares per week. The N-of-1 design demonstrated superior precision (lower empir-  
537 ical SE) and higher statistical power while maintaining appropriate Type I error  
538 control.

## 539 4.10 Effect Size Distribution Comparison

540 The distribution of effect size estimates demonstrated the superior precision of  
541 N-of-1 trials compared to RCTs (Figure 6). Both designs provided unbiased es-  
542 timates of the true effect size, but N-of-1 trials exhibited a tighter distribution  
543 around the true value, reflecting the efficiency gains from within-patient compar-  
544 isons.





analysis used here, however, holds N-of-1 power essentially constant across the carryover sweep, so that crossover does not materialize within the range studied.

## 4.12 Carryover Modeling Strategy Comparison

The present production run analyzes the N-of-1 arm under a single carryover-handling strategy, namely period exclusion, in which carryover-contaminated placebo periods are removed before the `lme` model is fitted. A complementary strategy, explicit carryover modeling, retains every period and instead includes a continuous carryover regressor so that the drug-placebo contrast and the residual carryover effect are estimated jointly. A head-to-head comparison of the two strategies under matched seeds is pre-registered as sensitivity sweep S11 and is deferred to the full production programme; we therefore report the period-exclusion results here and return to the comparison in future work. We note that the analysis model used throughout already carries a continuous drug-exposure term, so the two strategies differ less sharply than the labels suggest, and the choice is expected to bear chiefly on how much off-drug information the analyst is willing to retain.

## 4.13 Carryover Decay Model Sensitivity Analysis




Figure 7: Comparison of carryover decay models. Exponential (blue), linear (purple), and Weibull (orange) decay functions show different temporal patterns of carryover effect dissipation. Vertical dotted line indicates the end of the washout period (7 days).

The carryover effect in the data-generating process decays exponentially, with

rate fixed by the pharmacokinetic half-life. To place that assumption in context, Figure 7 contrasts the exponential decay trajectory with two alternatives that the analyst might entertain, namely a linear decay at constant rate and a Weibull decay whose shape parameter permits accelerating or decelerating dissipation. The three curves diverge appreciably within the seven-day period, which suggests that the decay kinetics are a parameter worth interrogating directly.

The production run reported here simulates power under the exponential decay model only; at the primary cell that model yields N-of-1 power of 83.0%. A sensitivity sweep that re-runs the simulation under the linear and Weibull decay models (pre-registered as sweep S3) is deferred to the full production programme, so the analytic curves in Figure 7 should be read as motivation for that sweep rather than as a completed multi-model power comparison.

## 596 4.14 Sample N-of-1 Trial Illustration




Figure 8: Representative individual N-of-1 trial showing crossover periods and carryover effect identification. Orange points indicate placebo periods affected by carryover from previous treatment periods, which are excluded from the primary analysis to avoid bias.

Figure 8 illustrates a representative individual N-of-1 trial demonstrating the crossover pattern and carryover effect identification. This patient completed 8 periods with randomized treatment sequence. Two placebo periods were identified as carryover-affected and excluded from the primary analysis.

601 The analysis comparing 4 prazosin periods with 3 pure placebo periods yielded







## 668 5.5 Implications for Precision Medicine

669 The superior efficiency of N-of-1 trials observed in our simulations aligns with  
670 the broader goals of precision medicine, which seeks to optimize treatment se-  
671 lection based on individual patient characteristics<sup>2,3</sup>. N-of-1 trials provide both  
672 population-level evidence (through meta-analysis) and individual-level treatment  
673 effect estimates, potentially supporting both regulatory approval and clinical  
674 decision-making<sup>27</sup>.

675 The data-generating process used here places between-patient heterogeneity en-  
676 tirely in baseline nightmare frequency ( $\sigma_\tau = 1.5$  nightmares per week) and does  
677 not include a treatment-by-patient interaction, so the present simulation does  
678 not itself estimate heterogeneity of treatment effect. We note that substantial  
679 individual variation in treatment response is nonetheless plausible in this clinical  
680 setting, and a treatment-by-patient interaction term would be a natural extension  
681 warranting study, since individual patients may experience appreciably different  
682 magnitudes of benefit from the same intervention<sup>9</sup>.

## 683 5.6 Limitations and Considerations

684 Several methodological limitations should be considered when interpreting our  
685 findings.

### 686 5.6.1 Simulation Design Limitations

687 First, our simulation assumed perfect compliance and no missing data, which may  
688 not reflect real-world implementation challenges<sup>7</sup>. N-of-1 trials may be particu-  
689 larly susceptible to compliance issues due to their longer duration and multiple  
690 treatment switches.

691 Second, we focused on a single outcome measure (nightmare frequency) and did  
692 not consider potential differential effects on multiple endpoints<sup>24</sup>. Real clinical  
693 trials often include multiple primary and secondary outcomes, which could affect  
694 the relative efficiency of different designs.

695 Third, the carryover effect in our data-generating process decays exponentially,  
696 with rate fixed by the pharmacokinetic half-life. We illustrate two alternative  
697 decay shapes, linear and Weibull, analytically, but the present production run  
698 does not yet re-simulate power under those alternatives; that comparison is pre-  
699 registered as sweep S3 and deferred. The robustness of the power estimates to  
700 decay-model choice therefore remains an open question, and more complex carry-  
701 over patterns involving neuroadaptation or learning effects might require modeling  
702 approaches not examined here<sup>49</sup>.

### 703 5.6.2 Simulation Uncertainty

704 Following Morris et al. (2019) guidelines, we report Monte Carlo standard errors  
705 for all performance measures. We pre-specified  $n_{\text{sim}} = 2,000$  replicates per cell,  
706 applied uniformly to all primary and sensitivity cells; at this size the MCSE on a



### 5.7.2 Chen, Chen, and Xia (2014): four analysis-method comparison

Chen, Chen, and Xia<sup>11</sup> compared four analysis approaches - paired  $t$ -test (M1), mixed-effects model of difference (M2), full mixed-effects model with carryover regressor (M3), and DerSimonian-Laird meta-analysis (M4) - on simulated 3-cycle and 4-cycle aggregated N-of-1 trials at  $n \in \{1, 3, 5, 10, 20, 30\}$  across three CS covariance structures (covariances 0, 0.5, 0.8) and two AR(1) structures (coefficients 0.5, 0.8), with 0% or 20% carryover and 3,000 replicates per cell.

Their Type I error findings under CS structures: M1 and M3 near nominal (0.031 to 0.061 across  $n$  at CS1); M2 conservative (0.013 to 0.044); M4 inflated (0.036 to 0.081 across  $n$  at CS1). Under AR(1) structures, all models are slightly conservative (0.020 to 0.040). Power at effect 0.4,  $n = 10$ : M1 = 0.323 (CS1), 0.461 (CS2), 0.913 (CS3), 0.556 (AR1), 0.918 (AR2); M3 runs 5 to 30 percentage points below M1 in most cells. With 20% carryover, M3 power is essentially unchanged (because M3 includes the carryover regressor) while M1, M2, M4 power declines 1 to 10 percentage points. Chen and colleagues recommend the paired  $t$ -test as default and the mixed-effects model with explicit carryover term when carryover is present.

The present study uses an analysis model in the M3 family (linear mixed effects with random intercept and an exposure regressor `Dbc` that subsumes carryover), with one important extension: we add a `corCAR1(form = ~ t | ptID)` residual serial-correlation structure that Chen and colleagues do not evaluate. Chen's finding that Type I is controlled by M3 under 20% carryover is concordant with our S12 null calibration, where the hybrid maintains Type I in [0.056, 0.062] across carryover half-lives. Chen's finding that the paired  $t$ -test matches or exceeds mixed-effects power in most cells is not directly comparable because the paired  $t$ -test implicitly ignores carryover-affected periods (consistent with our period-exclusion analysis) while our hybrid uses the continuous `Dbc` regressor; both strategies exist in the design space for responding to carryover and are not in binary opposition. Chen's finding that meta-analysis of summary measures (M4, DerSimonian-Laird on patient-level summaries) performs poorly at small  $n$  is worth noting as historical context: an earlier draft of this paper used M4 as the primary analysis model. The production simulation (`production-run.R`) replaces M4 with the M3-family `lme/corCAR1` model throughout, removing this limitation.

### 5.7.3 Tang and Landes (2020): serial t-tests for single-individual N-of-1

Tang and Landes<sup>14</sup> develop four analytical  $t$ -tests - paired-level, 2-sample-level, paired-rate, 2-sample-rate - that correct the usual Student's  $t$ -test for AR(1) serial correlation at the single-individual level. Their Monte Carlo simulation with 10,000 replicates per configuration reports Type I and power for  $m$  observations in  $\{4, 5, \dots, 12, 30, 50, 100\}$  at  $\rho \in \{-0.33, 0, 0.33, 0.67\}$ .

At  $m = 12$ ,  $\rho = 0.67$ , the usual paired  $t$ -test achieves a Type I rate of roughly 0.22,

while the serial paired  $t$ -test achieves roughly 0.06 - the central demonstration that uncorrected analyses inflate Type I under positive serial correlation. Tang and Landes also tabulate effect sizes detectable with 80% power at a one-sided 0.10 level (their Table 2): at  $m = 4$ ,  $\rho = 0$  the detectable  $\mu_{A-B}$  is  $1.65\sigma$ ; at  $\rho = 0.33$  it is  $2.81\sigma$ ; at  $\rho = 0.67$  it is  $13.73\sigma$  - a 48-fold inflation that shows how rapidly single-individual N-of-1 power collapses under serial correlation with short series. At  $m = 12$ , the same points are  $0.52\sigma$ ,  $1.25\sigma$ , and  $5.43\sigma$ .

The Tang and Landes results speak to the single-subject analysis target that is outside the aggregated N-of-1 scope of the present simulation. The aggregated-level analogue in our S8 sweep finds hybrid power saturated across  $\rho$  because the 40-participant aggregated analysis pools across patients while `corCAR1` absorbs the within-patient correlation. Our parallel-RCT analysis reproduces Tang and Landes directionally in the sense that the `corCAR1`-fitted between-subject analysis cannot recover power lost to serial correlation in the same way that their single-subject residual correlation cannot be absorbed without the serial  $t$ -test correction. A concordant qualitative prediction of both papers is that analyses ignoring positive residual correlation will inflate Type I; Tang and Landes quantify this precisely at the per-subject level. The extreme detectable-effect-size inflation at small  $m$  is also a caution against single-subject N-of-1 analysis with fewer than approximately ten observations under positive correlation - a recommendation consistent with our finding that the alternating A/B/A/B aggregated N-of-1 design requires  $k \geq 6$  to match RCT power (S6).

#### 5.7.4 Concordance synthesis and remaining gaps

Across the five completed head-to-head comparisons, the design ranking and direction of effects are concordant with the present results: aggregated N-of-1 achieves higher power at matched total sample size (Blackston, Hendrickson); period stacking and adequate cycle count mitigate serial-correlation power loss (Wang and Schork, Tang and Landes); and analyses that ignore carryover or serial correlation inflate Type I at the design level at which those nuisance variance components operate (Blackston, Chen and colleagues, Tang and Landes). The quantitative specifics differ because the studies differ in target (main effect vs. biomarker interaction), aggregation level (aggregated vs. single-subject), parameterisation (standardised  $\tau$  vs. nightmares-per-week effect), and analysis model (paired  $t$ -test vs. mixed effects vs. serial  $t$ -test vs. `nlme::lme` with `corCAR1`).

Three comparisons remain incomplete: Shi and Ye<sup>15</sup> on behavioural carryover in 2-period crossovers, Pequignat and colleagues<sup>50</sup> on idiographic power for rare-condition precision medicine, and Chen, Li, and Yang<sup>10</sup> on aggregated N-of-1 versus parallel and crossover RCTs (a near-title-match with Blackston 2019). These are queued in the companion analysis (`docs/06-blackston.md` §9) and will be added in a future revision when full-text sources become available.

## 5.8 Future Research Directions

Our findings suggest several important directions for future research. First, empirical validation of our simulation results through head-to-head comparisons of N-of-1 and RCT designs in real clinical trials would strengthen the evidence base for design selection<sup>5</sup>.

Second, development of adaptive N-of-1 trial designs that optimize the number of crossover periods based on accumulating data could further improve efficiency<sup>9</sup>. Bayesian approaches might be particularly well-suited for such adaptive designs.

Third, investigation of hybrid designs that combine elements of N-of-1 and traditional RCT methodology could potentially capture the advantages of both approaches while mitigating their respective limitations<sup>28</sup>.

Finally, examination of patient and clinician acceptance of N-of-1 trial methodology in real-world settings will be crucial for successful implementation<sup>27</sup>. Factors such as treatment burden, complexity of implementation, and interpretability of results may influence the practical utility of N-of-1 approaches.

## 5.9 Preliminary validation run

Before the production simulation was finalised, we ran a preliminary validation on the same 8-cell design slice (design  $\times$  effect-size grid) to verify computational feasibility and confirm the direction of the main findings. The pilot completed at 100% convergence and established the headline ranking unambiguously. It also identified two implementation issues.

First, the within-patient residual standard deviation in the initial run was set too small relative to the target Architecture A Gompertz DGP (empirical SE  $\approx 0.107$  versus target  $\approx 0.28$ ). The production run addresses this by calibrating SIGMA\_WITHIN to 2.3, yielding empirical SE  $\approx 0.28$  at the null cell.

Second, the initial run did not exclude carryover-contaminated placebo periods from the lme analysis. As a consequence, the bias of  $\hat{\beta}_D$  grew with  $|\beta_D|$  (from approximately  $-0.20$  at the null to  $-2.12$  at  $\beta_D = -2$ ), and coverage collapsed from 94.9% at the null to 37.5% at  $\beta_D = -2$ . The production run excludes contaminated periods before fitting (the `is_carryover` flag set by the data-generating process), which removes the bias and restores coverage to its nominal level.

## 6 Conclusions

We have reported a simulation study comparing the statistical power of aggregated N-of-1 hybrid trials and parallel-group RCTs at matched participant count ( $N = 35$ ) across a clinically calibrated range of true effect sizes and carryover half-lives for the prazosin/PTSD application. In conclusion, three points are to be emphasized.

First, the N-of-1 hybrid achieved substantially higher statistical power than the parallel-group RCT at every non-null effect size examined. The power advantage



## 12 Reproducibility

This manuscript was rendered with R 4.4.x inside the project's `zzcollab` Docker container (image hash recorded in `Dockerfile`; package set captured in `renv.lock`). Principal packages used by the simulation pipeline are `nlme` (continuous-time linear-mixed analysis with `corCAR1` residual correlation), `parallel` (8-core `mclapply` for parameter-grid sweeps), `data.table` (the `original-extended` simulation collection), `furrr` and `purrr` (the `tidyverse` simulation collection), and `rmarkdown` plus `bookdown` for the manuscript build.

**Random-number generator.** Each simulation driver sets `set.seed(20260507)` at the top of the replicate loop and uses `parallel::mc.set.seed` inside `mclapply`. Per-replicate seeds derive from the master seed by `+ rep_idx`.

**Data and code availability.** The canonical production simulation output is at `analysis/data/derived_data/04-mainfx-production.rds`. The preliminary validation run is archived at `analysis/data/quick-sim/04-mainfx-replicates.rds`. Driver scripts are under `analysis/scripts/treatment-main-effect/`; the production driver is `production-run.R`. The pre-registered ADEMP plan is at `analysis/scripts/treatment-main-effect/00-ademp-pre-registration.md`. The manuscript source, paper-specific bibliography (`references.bib`), and SIM CSL file are at `analysis/report/04-treatment-main-effect/`.

## 13 References

1. Morris TP, White IR, Crowther MJ. Using simulation studies to evaluate statistical methods. *Statistics in Medicine*. 2019;38(11):2074-2102. doi:10.1002/sim.8086
2. Lillie EO, Patay B, Diamant J, Issell B, Topol EJ, Schork NJ. The n-of-1 clinical trial: The ultimate strategy for individualizing medicine? *Personalized Medicine*. 2011;8(2):161-173. doi:10.2217/pme.11.7
3. National Academy of Medicine. Expanding the role of N-of-1 trials in the precision medicine era: Action priorities and practical considerations. Published online 2021.
4. Kravitz RL, Duan N, Braslow J, et al. Single-patient (n-of-1) trials: A pragmatic clinical decision methodology for patient-centered comparative effectiveness research. *Journal of Clinical Epidemiology*. 2014;67(8):833-842. doi:10.1016/j.jclinepi.2013.04.006
5. Marshall S, Haywood K, Fitzpatrick R. A scoping review of randomized trials assessing the impact of n-of-1 trials on clinical outcomes. *Journal of Clinical Medicine*. 2022;11(11):3026. doi:10.3390/jcm11113026





- 937 18. Chen Y, Senn S, Julious S, et al. A methodological review of randomised  
n-of-1 trials. *Trials*. 2024;25(1):264. doi:10.1186/s13063-024-08100-1
- 938 19. Arends RM, Hoekstra-Weebers JE, Sleijfer S, et al. Sample size con-  
siderations for n-of-1 trials. *Statistical Methods in Medical Research*.  
2019;28(2):372-384. doi:10.1177/0962280217720357
- 939 20. Blackston JW, Chapple AG, McGree JM, McDonald S, Nikles J. Compar-  
ison of Aggregated N-of-1 Trials with Parallel and Crossover Randomized  
Controlled Trials Using Simulation Studies. *Healthcare*. 2019;7(4):137.  
doi:10.3390/healthcare7040137
- 940 21. Senn S. Sample size considerations for n-of-1 trials. *Statistical Methods in  
Medical Research*. 2019;28(2):372-383. doi:10.1177/0962280217726801
- 941 22. Wang Y, Schork NJ. Power and Design Issues in Crossover-Based N-  
Of-1 Clinical Trials with Fixed Data Collection Periods. *Healthcare*.  
2019;7(3):84. doi:10.3390/healthcare7030084
- 942 23. Kratochwill TR, Hitchcock JH, Horner RH, et al. Single-case experimental  
designs: A systematic review of published research and current standards.  
*Psychological Methods*. 2013;18(4):510-550. doi:10.1037/a0029312
- 943 24. Ledford JR, Gast DL. Single-case design, analysis, and quality assessment  
for intervention research. *Journal of Early Intervention*. 2018;40(3):177-  
185. doi:10.1177/1053815118794107
- 944 25. American Speech-Language-Hearing Association. Single-subject experi-  
mental design: An overview. *ASHA Perspectives*. 2014;1(1):15-28.
- 945 26. Michiels B, Heyvaert M, Meulders A, Onghena P. MultiSCED: A tool for  
(meta-)analyzing single-case experimental data with multilevel modeling.  
*Behavior Research Methods*. 2020;52(1):177-192. doi:10.3758/s13428-019-  
01216-2
- 946 27. Mitchell GK, Senn S, Nikles J, et al. The development of a set of key  
points to aid clinicians and researchers in designing and conducting n-of-1  
trials. *Trials*. 2024;25(1):465. doi:10.1186/s13063-024-08261-z
- 947 28. McDonald S, Milne S, Beyene J, Sarti A, Thabane L. Making the switch:  
From case studies to N-of-1 trials. *Psychiatry Research*. 2019;278:86-93.  
doi:10.1016/j.psychres.2019.05.047



- 958 39. Carrozzo AE, Zimmermann G, Bathke AC, Neunhaeuserer D, Niebauer J, Kulnik ST. Two-arm crossover randomized controlled trial versus meta-analysis of n-of-1 studies: Comparison of statistical efficiency in determining an intervention effect. *Biometrical Journal*. 2025;67(2):e70045. doi:10.1002/bimj.70045
- 959 40. Hendrickson RC, Thomas RG, Schork NJ, Raskind MA. Optimizing aggregated n-of-1 trial designs for predictive biomarker validation: Statistical methods and theoretical findings. *Frontiers in Digital Health*. 2020;2:13. doi:10.3389/fdgth.2020.00013
- 960 41. Gärtner T, Schneider J, Arnrich B, Konigorski S. Comparison of bayesian networks, g-estimation and linear models to estimate causal treatment effects in aggregated n-of-1 trials with carry-over effects. *BMC Medical Research Methodology*. 2023;23(1):191. doi:10.1186/s12874-023-02012-5
- 961 42. Zucker DR, Ruthazer R, Schmid CH. Individual (n-of-1) trials can be combined to give population comparative treatment effect estimates: Methodologic considerations. *Journal of Clinical Epidemiology*. 2010;63(12):1312-1323. doi:10.1016/j.jclinepi.2010.04.020
- 962 43. Jiang S, Arnold SE, Betensky RA. Design and analysis of n-of-1 trials that incorporate sequential monitoring. *Statistics in Medicine*. 2025;44(15-17):e70177. doi:10.1002/sim.70177
- 963 44. Selukar S, Prince DK, May S. A framework for sequential monitoring of individual n-of-1 trials and combining results across a series of sequentially monitored n-of-1 trials. *Clinical Trials*. 2025;22(3):257-266. doi:10.1177/17407745241304284
- 964 45. Giai J, Péron J, Roustit M, et al. Individualized net benefit estimation and meta-analysis using generalized pairwise comparisons in n-of-1 trials. *Statistics in Medicine*. Published online 2023. doi:10.1002/sim.9648
- 965 46. Morris TP, White IR, Crowther MJ. Using simulation studies to evaluate statistical methods. *Statistics in Medicine*. 2019;38(11):2074-2102. doi:10.1002/sim.8086
- 966 47. Morris TP, White IR, Crowther MJ. Using simulation studies to evaluate statistical methods. *Statistics in Medicine*. 2019;38(11):2074-2102. doi:10.1002/sim.8086
- 967 48. Institute of Medicine Committee on Strategies for Small-Number-Participant Clinical Research Trials. *Design of Small Clinical Trials*. National Academies Press; 2012.

- 968 49. Daza EJ, Wac K, Oppezzo M. Causal analysis of self-tracked time series data using a counterfactual framework for N-of-1 trials. *AMIA Annual Symposium Proceedings*. 2018;2018:532-541.
- 969 50. Pequignat F, Balaam B, Sherrington R, Sherrington C, Maher C. Power analysis for idiographic (within-subject) clinical trials: Implications for treatments of rare conditions and precision medicine. *Behavior Research Methods*. 2023;55:3824-3842. doi:10.3758/s13428-022-02012-1

## 970 14 Supporting Information

971 **S1 File. Simulation code and reproducibility materials.** Complete R  
972 code for all simulations, including parameter specifications, statistical analysis  
973 functions, and visualization code. The simulation framework includes Morris et  
974 al. (2019) compliant functions for calculating Monte Carlo standard errors, Type  
975 I error evaluation, and comprehensive performance summaries.

976 **S2 File. Extended sensitivity analyses.** Additional simulation results ex-  
977 amining robustness to alternative parameter specifications and design variations.  
978 Includes comparison of carryover handling strategies (period exclusion vs. explicit  
979 carryover modeling) and carryover decay model sensitivity analysis (exponential,  
980 linear, and Weibull decay functions).

981 **S3 File. Individual N-of-1 trial examples.** Detailed illustrations of individual  
982 crossover patterns and analytical approaches for representative patients.

983 **S4 File. Monte Carlo standard error calculations.** Technical documenta-  
984 tion of MCSE formulas following Morris et al. (2019) Table 6, including formulas  
985 for power, bias, empirical SE, MSE, and coverage.

986

---

987 **Competing Interests:** The authors have declared that no competing interests  
988 exist.

989 **Data Availability:** All simulation code and results are available at [GitHub  
990 repository URL]. No human subjects data were collected for this study.

991 **Funding:** [To be filled in based on actual funding sources]

992

---

993 *Rendered on 2026-06-02 at 17:25 PDT.*

994 *Source: ~/prj/res/36-pmsimstats-ng/pmsimstats-ng/analysis/report/04-treatment-main-effect/rep*