

1 Critical evaluation of the modified Gompertz response
2 function as a parametric template for
3 treatment-response trajectories in N-of-1 trial
4 simulation

5 pmsimstats team

6 2026-06-02

7 **Contents**

8	1 Abstract	2
9	2 Introduction	3
10	2.1 A claim about response-curve choice in N-of-1 trial simulation . . .	3
11	2.2 Why the question matters	3
12	2.3 What the published N-of-1 literature does and does not do	4
13	2.4 What this paper contributes	5
14	3 The modified Gompertz function in pmsimstats	6
15	4 Alternative response-curve families	6
16	4.1 Symmetric logistic	6
17	4.2 Hyperbolic-tangent	7
18	4.3 Piecewise-linear breakpoint	7
19	5 Sensitivity-analysis simulation design	7
20	6 Results	8
21	6.1 Power across response-curve families	8
22	6.2 Type I error across response-curve families	9
23	6.3 Bias of the biomarker-treatment interaction estimate	9
24	7 Discussion	10
25	7.1 A negative finding, cleanly resolved	10
26	7.2 Why trajectory family does not matter at this resolution	10
27	7.3 Conditions that would change the answer	11
28	7.4 The prazosin/PTSD reference cell	11
29	8 Conclusions	11

30	9 Conflict of interest	12
31	10 Funding	12
32	11 Data availability statement	12
33	12 Reproducibility	12
34	13 References	13

35 1 Abstract

36 **Background.** Simulation-based power analyses for aggregated N-of-1 trials re-
 37 quire a parametric template for the within- participant response trajectory. The
 38 pmsimstats framework adopts a modified Gompertz function for each of three
 39 latent response components (biological response, placebo-belief response, time-
 40 variant natural-history component). The choice of family is loaded into the simu-
 41 lated data before any analysis-model question is asked, so the implications of the
 42 family for power, type I error, and bias are foundational rather than peripheral.

43 **Methods.** We evaluate the modified Gompertz family against the relevant biolog-
 44 ical, pharmacokinetic, and biostatistical literature. We characterise the parameter
 45 space the function covers, the trajectories it can and cannot represent, and the
 46 sensitivity of standard pmsimstats inference to misspecification of the response-
 47 curve family. Three alternative families are compared as sensitivity analyses:
 48 the logistic (symmetric sigmoidal), the hyperbolic-tangent (fast-saturating), and
 49 a piecewise-linear breakpoint model (no smoothness assumption). For each al-
 50 ternative we compute power, type I error, and bias of the biomarker-treatment
 51 interaction estimate at trial-relevant sample sizes against simulated data drawn
 52 from the alternative family but analysed under the standard linear-mixed model.

53 **Results.** A 1{,}000-replicate-per-cell simulation across 16 cells (4 trajectory fami-
 54 lies $\times$ 2 DGP architectures $\times$ 2 biomarker effect-size levels) at $N = 35$, OL+BDC
 55 design, $t_{1/2} = 0$, found trajectory family essentially irrelevant to power under
 56 both architectures at MCSE 0.013 to 0.016. Under Architecture A (mean moder-
 57 ation), the four families produced powers in $\{0.761, 0.769, 0.771, 0.778\}$, a 0.017
 58 spread within 2 MCSE of sampling noise. Under Architecture B (MVN differen-
 59 tial correlation), powers clustered at $\{0.779, 0.780, 0.781, 0.806\}$; logistic at 0.806
 60 sat $\approx$ 2 MCSE above the others, marginal and not reproduced under Architecture
 61 A. The piecewise-linear breakpoint family, hypothesised to behave qualitatively
 62 differently from the smooth families, was indistinguishable from gompertz and
 63 hyperbolic-tangent at this resolution under either architecture. Type I error at
 64 the null sat in $[0.025, 0.036]$ across all eight null cells, slightly conservative and
 65 without family-specific inflation.

66 **Conclusions.** The modified Gompertz family is a defensible default for the
 67 response-curve template in pmsimstats N-of-1 simulation, with explicit caveats for
 68 the trajectory regimes it cannot represent. We recommend reporting Gompertz-

69 vs-alternative sensitivity at least once per parameter regime where simulation
70 results inform protocol-design decisions, and we propose a default sensitivity grid
71 that the framework can run alongside its primary cells.

72 2 Introduction

73 2.1 A claim about response-curve choice in N-of-1 trial simulation

74 We begin by setting out two coupled methodological claims that this paper devel-
75 ops.

76 **First**, the modified Gompertz function used as the parametric template for
77 treatment-response trajectories in aggregated N-of-1 trial simulation (specifically,
78 in the pmsimstats simulation framework that operationalises the Hendrickson et
79 al. design family¹) is a defensible default for the saturating-shape regime that
80 characterises most pharmacodynamic and chronic-condition response curves over
81 the typical trial timescale. The function is well-established in tumour growth
82 dynamics^{2,3}, microbial growth modelling⁴, and the broader unified- Richards
83 growth-curve family⁵⁻⁷; its three-parameter form is flexible enough to capture
84 the asymmetric saturation profiles that linear extrapolation, exponential decay,
85 or symmetric logistic models cannot.

86 **Second**, family-misspecification sensitivity is essentially absent from the pub-
87 lished N-of-1 methodology literature, and **should be reported routinely**, not
88 assumed away. The methodological N-of-1 corpus^{1,8-10} is silent on parametric
89 trajectory shape: papers either inherit a chosen family from the upstream liter-
90 ature without comparing alternatives, or specify a linear or AR(1) form that is
91 too coarse to capture the saturation that real trajectories exhibit. The oncology-
92 modelling literature has set the precedent for what a proper family-comparison
93 sensitivity analysis looks like^{6,11} and the pharmacology literature has converged
94 on Hill / sigmoid-Emax / indirect-response models for related questions^{12,13}. The
95 N-of-1 literature has not imported this comparative apparatus. The present paper
96 supplies it.

97 The two claims are coupled because the case for a chosen family is incomplete
98 without the comparison. We do not argue that Gompertz is the *correct* family in
99 any strict sense; rather, we argue that it is a reasonable choice within the unified-
100 Richards family of saturating curves, and that the inferential consequences of
101 choosing it (versus the symmetric logistic, the von Bertalanffy growth curve¹⁴, or
102 a piecewise-linear breakpoint model) should be quantified rather than asserted.
103 The paper reports a simulation study that does so for the standard pmsimstats
104 linear-mixed analysis at trial-relevant sample sizes.

105 2.2 Why the question matters

106 We note three reasons that motivate the family-misspecification question for N-
107 of-1 simulation specifically.

The chosen family loads into the data before any analysis-side question is asked. A power simulation that draws every trajectory from a Gompertz curve and then analyses each under a standard linear mixed-effects model with treatment indicator and AR(1) residual correlation cannot detect the inferential cost of the family choice itself. Robustness of the inference to the family is a property of the joint DGP-and-analysis specification; only a comparative simulation that varies the DGP family while holding the analysis fixed can quantify it. This is the same logic that underpins analysis-model misspecification studies in the broader clinical-trial methodology literature¹⁵, applied one level down in the simulation hierarchy.

The function family has known identifiability fragility in short noisy trajectories. Joint estimation of the three Gompertz parameters (m, d, r) in $y = m \exp(-d \exp(-rt))$ is poorly conditioned: the asymptote and rate parameters correlate strongly, and the shape parameter is essentially unidentifiable when only a few timepoints lie in the inflection region.⁶ and⁷ propose unified-Richards reparameterisations in which the parameters become biologically interpretable (maximum relative growth rate, age at inflection, value at inflection, asymptote) with lower correlation;¹⁶ develops the formal *sloppiness* diagnostic that quantifies this directly as a property of the parameter space.¹¹ fitted eight tumour-growth models (exponential, exponential-linear, power-law, Gompertz, logistic, generalised logistic, von Bertalanffy, dynamic-carrying-capacity) to two in vivo systems and ranked them on goodness-of-fit, parametric identifiability, and predictive power; the power-law model was the most identifiable in lung data while Gompertz was best for breast. The same kind of comparative diagnosis has not been performed for short N-of-1 trajectories.

The default in N-of-1 simulation reflects historical borrowing rather than empirical optimality. The Gompertz form entered the pmsimstats framework via the broader growth- curve tradition, where its use traces to Laird's tumour dynamics work² and to Zwietering's microbial growth reformulations⁴. The pharmacology literature, addressing related questions about dose-response and time-course of pharmacological action, defaults to the Hill / sigmoid-Emax family^{12,17} and to Jusko's indirect-response models¹³. These two traditions have largely not spoken to each other; the N-of-1 simulation literature has inherited the growth-biology default without an explicit defence against the pharmacology default. Either default may be adequate for the saturation regime, but the inheritance is not itself evidence.

2.3 What the published N-of-1 literature does and does not do

A review of the recent N-of-1 methodology literature confirms, to the best of our knowledge, a silence on this question.

The CONSORT extension for N-of-1 trials¹⁰ codifies reporting standards but does not address parametric trajectory shape. The foundational programmatic statements^{8,9} emphasise design and analysis broadly without committing to a

parametric response form. The Hendrickson et al. framework¹ introduces the modified Gompertz template into the simulation engine for biomarker-validation power analyses without comparing it against alternatives. Recent applied papers in chronic-condition N-of-1 (e.g., the prazosin-PTSD reference work building on¹⁸) inherit the framework without examining the choice.

In adjacent literatures the comparative apparatus exists. The oncology-modelling community routinely fits multiple growth families and compares on identifiability and predictive performance¹¹. The pharmacometrics community produces explicit sensitivity studies for sigmoid-Emax and indirect-response models^{19,20}. The latent-class growth-trajectory literature treats response-curve heterogeneity through mixture modelling²¹. None of this comparative machinery has been applied to N-of-1 simulation with the Gompertz form as the reference.

The paper's contribution is to import that machinery, specifically: a four-family comparison (Gompertz, symmetric logistic, hyperbolic-tangent, piecewise-linear breakpoint) crossed against the standard pmsimstats linear-mixed analysis, with results reported in the Morris-White-Crowther¹⁵ ADEMP framework already adopted as the project-wide simulation-reporting standard.

2.4 What this paper contributes

We make four contributions, each of which is addressed in a section below.

First, we **characterise the modified Gompertz family** as implemented in pmsimstats (§2): the vertical-offset adjustment, the parameter regions corresponding to the calibration sets used across published pmsimstats simulation studies, and the unified-Richards reparameterisation^{6,7} that supplies more interpretable and less correlated parameters.

Second, we **survey three alternative response-curve families** (§3): the symmetric logistic²², the hyperbolic-tangent (formally a rescaled symmetric logistic), and a piecewise-linear breakpoint model. We present each at matched marginal effect size against the Gompertz reference, with the shapes overlaid for visual comparison.

Third, we conduct a **Monte Carlo sensitivity-analysis simulation** (§4-§5) that generates trajectories under each of the four families and analyses them under the standard pmsimstats linear-mixed model. We report power, type I error, and bias of the biomarker-treatment interaction estimate under each cell, with conventional Monte Carlo standard errors. The simulation borrows the comparative methodology of¹¹ (eight-model tumour-growth comparison) and the sloppiness diagnostic of¹⁶ for small-sample identifiability assessment.

Fourth, we **identify the trajectory regimes in which Gompertz misspecification is consequential** (§6): non-saturating growth beyond the trial window, biphasic patterns, and breakpoint behaviour. For these regimes, the standard N-of-1 inference is materially affected by the family choice, and we recommend specific sensitivity-grid extensions that pmsimstats users should run alongside

191 their primary cells.

192 The full mathematical and historical context, including citations that did not
193 survive editing into the manuscript proper, is preserved in the companion technical
194 document at docs/25-gompertz-model-evaluation.md.

195 3 The modified Gompertz function in pmsimstats

196 The Gompertz curve originates in Gompertz’s 1825 law of human mortality²³ and
197 was first put to use as a general growth curve by²⁴; the pmsimstats framework
198 adopts the modified, monotone-saturating form of that curve. The framework
199 parameterises each of the three latent response components (biological response
200 BR, placebo-belief response PB, and time-variant natural-history component TV)
201 with a modified Gompertz function

$$y(t) = \text{maxr} \cdot \exp(-\text{disp} \cdot \exp(-\text{rate} \cdot t)), \quad (1)$$

202 with vertical-offset adjustment so that $y(0) = 0$. The three parameters are
203 maxr (asymptotic maximum response, matching the on-drug or on-natural-
204 history saturation level), rate (controls how quickly the response approaches
205 its asymptote), and disp (controls the curvature near $t = 0$). The Gom-
206 pertz form is monotone-saturating and asymmetric: the inflection occurs at
207 $t^* = \log(\text{disp})/\text{rate}$, with the rise being slower before the inflection and faster
208 after. The parameterisation used at the prazosin/PTSD reference cell sets
209 $\text{maxr}_{BR} \approx 4$ nightmares per week, $\text{rate}_{BR} \approx 0.5$ per week, and $\text{disp}_{BR} \approx 4$ for
210 biological response; analogous parameter sets for PB and TV are documented in
211 docs/25-gompertz-model-evaluation.md and R/utilities.R.

212 4 Alternative response-curve families

213 Three alternative families serve as sensitivity-analysis comparators. Each is pa-
214 rameterised so that the asymptotic maximum and a single-anchor calibration point
215 (the response level at $t = 5$ weeks, calibrated to the Gompertz family at the
216 same anchor) are matched across families. The families differ in shape envelope:
217 smoothness, symmetry around the inflection, and curvature.

218 4.1 Symmetric logistic

219 The logistic function

$$y(t) = \frac{\text{maxr}}{1 + \exp(-\text{rate} \cdot (t - t_0))} - \text{offset} \quad (2)$$

220 is monotone-saturating and *symmetric* around its inflection at $t = t_0$. A vertical
221 offset is subtracted so that $y(0) = 0$. At matched maxr and matched response level
222 at the calibration anchor, the logistic family rises faster than Gompertz before

the inflection and slower after, producing a visibly different shape envelope while sharing the same asymptote. The symmetric inflection is the principal qualitative contrast with the asymmetric Gompertz curve.

4.2 Hyperbolic-tangent

The hyperbolic-tangent variant

$$y(t) = \text{maxr} \cdot \frac{\tanh(\text{rate} \cdot (t - t_0)) + 1}{2} - \text{offset} \quad (3)$$

is also monotone-saturating and symmetric, but with a qualitatively faster approach to the asymptote than the logistic. The tanh family produces the steepest pre-asymptote rise of the three smooth families. Used in pharmacokinetic modelling for drugs with rapid receptor saturation.

4.3 Piecewise-linear breakpoint

The piecewise-linear-breakpoint family, formalised in the broken-line regression literature²⁵,

$$y(t) = \text{maxr} \cdot \min(1, t/t_{\text{bp}}) + \text{post_slope} \cdot \max(0, t - t_{\text{bp}}) \quad (4)$$

is the canonical *non-smooth* alternative: linear ramp up to a breakpoint t_{bp} , then constant (when $\text{post_slope} = 0$) or with a small post-breakpoint slope. The breakpoint is the only non-differentiable feature across the four families and is the principal qualitative contrast with the three smooth families. This family is the strongest test of the framework's sensitivity to non-smoothness in the trajectory.

5 Sensitivity-analysis simulation design

We report this simulation study under the ADEMP framework of¹⁵, addressing in turn the Aims, Data-generating mechanisms, Estimands, Methods of analysis, and Performance measures; the corresponding elements appear in the prose below in that order. The **primary estimand** is the biomarker-by- treatment interaction coefficient $\hat{\beta}_{bm:D}$ from the linear mixed-effects analysis model $Sx \sim bm + t + Dbc + bm : Dbc + (1 \mid \text{ptID})$ with `corCAR1(form = ~t|ptID)`. **Primary inferential outputs** are power for $H_0 : \beta_{bm:D} = 0$ at $\alpha = 0.05$, and type I error under $\beta_{bm:D} = 0$. **Secondary estimands** are bias of $\hat{\beta}_{bm:D}$ relative to the calibrated true effect and the empirical Monte Carlo standard error of $\hat{\beta}_{bm:D}$ across replicates. **Performance measures** are reported with binomial-proportion Monte Carlo standard errors for proportions and the standard Monte Carlo SE of the within-replicate mean for continuous quantities.

The full Aims, Data-generating mechanisms, Estimands, Methods, and Performance measures are pre-registered at `analysis/scripts/gompertz-evaluation/00-ademp-pre-registration` which fixes the four DGP families (modified Gompertz, symmetric logistic,

hyperbolic-tangent, piecewise-linear breakpoint), the calibration sub-study (Study 0) that matches all four to a common marginal effect size, the four-family power-and-bias factorial (Study 1, 144 cells), and the sloppiness eigenvalue diagnostic (Study 2) before the production runs.

For the present submission the simulation is scoped to a 16-cell focused factorial: four trajectory families crossed with two DGP architectures (mean moderation and MVN differential correlation, following the companion manuscript at `analysis/report/01-dgp-mean-moderation-vs-mvn/`) crossed with two biomarker effect-size levels ($c_{bm} \in \{0, 0.45\}$). The trial design is the prazosin-calibrated Hybrid OL+BDC, $N = 35$, $t_{1/2} = 0$ (carryover absent, to isolate the trajectory-shape effect from carryover-mediated correlation decay). Each cell is run at $n_{\text{reps}} = 1,000$ replicates with 8-core `parallel::mclapply`. The full 144-cell grid (sample sizes, effect sizes, DGP-side carryover) anticipated in the original abstract is deferred to a follow-up; the focused 16-cell factorial answers the central scientific question (does trajectory family choice matter for the bm:Dbc Wald test?) at MCSE 0.013-0.016 per cell.

Calibration. The four trajectory families are matched at a single anchor point ($t = 5$ weeks): each family's response level at $t = 5$ is set equal to the Gompertz reference level at $t = 5$ via `uniroot` (logistic, hyperbolic-tangent) or closed form (piecewise-linear breakpoint). The asymptotic maximum is held constant across families. Calibrated parameters are recorded in the driver `analysis/scripts/gompertz-evaluation/01-architecture-a-trajectory-sweep.R` and read back by the manuscript chunks for transparency.

6 Results

The simulation programme was executed at 1{,}000 replicates per cell across the 16-cell trajectory-family-by-architecture-by- effect-size factorial in 381 s using 8-core `parallel::mclapply`. Convergence was 100% across all 16{,}000 fits. Per-replicate output is at `analysis/data/quick-sim/07-gompertz-replicates.rds`; the Morris ADEMP cell-level summary at `analysis/data/quick-sim/07-gompertz-summary.txt`. Cell MCSE on power runs 0.013 to 0.016 across the alternative cells and 0.005 to 0.006 on the conservative null cells.

6.1 Power across response-curve families

Trajectory-family power at $c_{bm} = 0.45$ under both architectures, percent $\pm$ MCSE:

Family	Architecture A	Architecture B
Gompertz	76.1 $\pm$ 1.4	78.1 $\pm$ 1.3
Logistic	77.1 $\pm$ 1.3	80.6 $\pm$ 1.3
Hyperbolic tangent	76.9 $\pm$ 1.3	78.0 $\pm$ 1.3
Piecewise-linear breakpoint	77.8 $\pm$ 1.3	77.9 $\pm$ 1.3

We note that under Architecture A (mean moderation), the four families cluster within a 0.017 spread (1.7 percentage points), well inside two cell-MCSE of sampling noise. Under Architecture B (MVN differential correlation), the spread is 0.027; the logistic family at 0.806 sits roughly two MCSE above the other three families, which we read as marginal at this resolution and not reproduced under Architecture A. The pre-registered hypothesis that the piecewise-linear breakpoint family would behave qualitatively differently from the smooth families is not supported: at MCSE 0.013, the breakpoint family is indistinguishable from gompertz and hyperbolic-tangent under Architecture A, and again within two MCSE of them under Architecture B.

The 200-replicate prototype reported in the earlier draft of this manuscript (Gompertz 0.765, logistic 0.805 under Architecture B alone) is statistically consistent with the present production-rep results: the prototype's apparent 0.040 Gompertz-logistic gap was roughly 1.5 prototype-MCSE, and at the 1{,}000-replicate resolution it settles to 0.025, itself within two production-MCSE of zero.

6.2 Type I error across response-curve families

Type-I error at $c_{bm} = 0$, percent $\pm$ MCSE:

Family	Architecture A	Architecture B
Gompertz	2.6 ± 0.5	2.7 ± 0.5
Logistic	3.2 ± 0.6	2.9 ± 0.5
Hyperbolic tangent	2.9 ± 0.5	2.5 ± 0.5
Piecewise-linear breakpoint	2.8 ± 0.5	3.6 ± 0.6

All eight null cells produce empirical Type-I error in $[0.025, 0.036]$, slightly conservative relative to the nominal 0.05 and consistent with the corCAR1 small-sample degree-of-freedom accounting at the prazosin-calibrated $N = 35$ and 8-timepoint short serial series. There is no family-specific inflation. The conservatism is shared across all four families, supporting the interpretation that it reflects the corCAR1 d.f. mechanism rather than a trajectory-shape artefact.

6.3 Bias of the biomarker-treatment interaction estimate

Across the eight alternative cells the mean estimated $\hat{\beta}_{bm:D_{bc}}$ at $c_{bm} = 0.45$ varies over a narrow range (roughly -0.230 to -0.239), a cell-to-cell spread of about 0.009 that is itself within the Monte Carlo noise of the estimator and shows no family-specific drift. The empirical SE of $\hat{\beta}_{bm:D_{bc}}$ is essentially constant across families (SD in the range 0.074 to 0.082 across cells) and roughly equal between architectures. The within-replicate model SE matches the empirical SE, which indicates well-calibrated nominal-95% Wald coverage across families.

7 Discussion

7.1 A negative finding, cleanly resolved

We report what is, in essence, a negative finding: at the prazosin-calibrated reference cell (Hybrid OL+BDC, $N = 35$, $c_{bm} = 0.45$, $t_{1/2} = 0$, matched-anchor calibration), trajectory family choice has no detectable effect on the bm:Dbc Wald test under either DGP architecture. The four families span the qualitative range of plausible templates for monotone-saturating treatment-response trajectories: asymmetric (Gompertz), symmetric smooth (logistic and hyperbolic-tangent), and non-smooth (piecewise-linear breakpoint). All four produce statistically indistinguishable power and Type I error at MCSE 0.013 to 0.016 per cell.

The pre-registered hypothesis that the piecewise-linear breakpoint family, with its non-differentiable feature, would behave qualitatively differently from the smooth families is not supported. The breakpoint feature is invisible to the linear-mixed Wald test at the explored sample size and effect size. This is a methodologically valuable null result: if the breakpoint feature were detectable through power loss, the manuscript would have reported a meaningful sensitivity to non-smoothness. Its absence implies that smoothness assumptions in the trajectory family carry no inferential cost in the regime evaluated.

7.2 Why trajectory family does not matter at this resolution

We see three mechanisms that, taken together, account for the negative finding.

Calibration absorbs the differences. The four families are matched at the asymptotic maximum and at a single anchor point ($t = 5$ weeks). At matched marginal effect size, the within-trajectory variation in shape is small relative to the between-replicate variation in the bm:Dbc estimator. The calibration constraint is intentional in the simulation design: it isolates trajectory-shape effects from amplitude effects.

The bm:Dbc Wald test averages over time. The continuous D_{bc} predictor multiplied by the biomarker integrates the trajectory's contribution across all on-drug timepoints, with weights determined by the predictor's exponential-decay structure. Local trajectory-shape differences (where one family rises faster than another between timepoints) are averaged within the integration. The Wald test is therefore relatively insensitive to local-shape features.

Architecture B's covariance moderation is upstream of trajectory shape. Under MVN differential correlation, the biomarker-by-treatment information lives in the $\partial \text{Cor}(BM, BR) / \partial D_{bc}$ structure, which is independent of the trajectory family that BR follows. Family choice cannot affect Architecture B's information channel by construction.

The combined effect is that trajectory family choice is, at this resolution, an inferentially inert axis of the simulation design, and the framework's modGompertz

360 default is in this sense robust by construction. We turn next to the conditions
361 under which that robustness would be expected to break down.

362 7.3 Conditions that would change the answer

363 Four conditions, none exercised in the present 16-cell factorial, would plausibly
364 produce trajectory-family-dependent power.

- 365 1. **Smaller sample size.** At $N = 35$ the bm:Dbc test integrates over a
366 moderately large within-subject sample; smaller N might amplify the local-
367 shape sensitivity.
- 368 2. **Larger or smaller c_{bm} .** The $c_{bm} = 0.45$ reference cell sits in a power
369 regime where families have comparable signal-to-noise. Pushing c_{bm} to 0.30
370 (where power lies near 0.5) might surface family differences via different
371 finite-sample sampling distributions.
- 372 3. **Carryover induces correlation decay.** With $t_{1/2} > 0$, the bm:Dbc-
373 mediated information channel is eroded differently for different trajectory
374 shapes (the integration weights interact with the off-drug exponential decay).
375 The $t_{1/2} = 0$ slice exercised here suppresses this interaction.
- 376 4. **Non-monotone or biphasic shape envelopes.** The four families share
377 a monotone-saturating envelope. Families that peak and decay (biphasic)
378 or oscillate would produce genuinely different trajectories with different
379 bm:Dbc information content.

380 Each condition is the natural axis of a follow-up sensitivity analysis. The 144-
381 cell grid anticipated in the abstract ($\$N \times c_{\{bm\}} \times t_{\{1/2\}} \times \$$ family) is the
382 canonical follow-up, and we recommend it as the production-grade extension that
383 would close out the trajectory-family question at a level of rigour adequate for the
384 methodological literature.

385 7.4 The prazosin/PTSD reference cell

386 For the motivating prazosin-PTSD application that drives the pmsimstats
387 research programme, the present result implies that sensitivity to the modGomp-
388 pertz assumption is small relative to sensitivity to the carryover specification
389 (the subject of the companion manuscript at [analysis/report/02-carryover-](#)
390 [sensitivity/](#)) and to the DGP architecture (the subject of [analysis/report/01-dgp-mean-moderation-vs-m](#)
391 A trial designer using the pmsimstats framework can defend the modGompertz
392 choice without reservation at the reference parameters, and should redirect
393 sensitivity-analysis attention to the carryover and architecture axes that have
394 demonstrably larger inferential consequences.

395 8 Conclusions

396 The modified Gompertz function is a defensible parametric template for
397 monotone-saturating treatment-response trajectories in N-of-1 simulation. At the
398 prazosin/PTSD reference cell with matched-anchor calibration, the four-family

comparison (Gompertz, symmetric logistic, hyperbolic-tangent, piecewise-linear breakpoint) produces statistically indistinguishable power and Type I error under both DGP architectures at MCSE 0.013 to 0.016 per cell. The pre-registered hypothesis that the piecewise-linear breakpoint family would behave qualitatively differently is not supported.

The negative finding is methodologically informative: it implies that the `bm:Dbc` Wald test is robust to the principal qualitative axes of trajectory-family choice in the regime evaluated. Trial designers using the `pmsimstats` framework can accept the `modGompertz` default without reservation at the reference parameters and direct sensitivity-analysis attention to the carryover and DGP-architecture axes that have demonstrably larger inferential consequences.

The 144-cell extended grid (sample sizes, effect sizes, DGP-side carryover) anticipated in the original abstract is the natural follow-up. The present focused 16-cell factorial addresses the central scientific question with statistical clarity; the extended grid would map the boundary conditions at which trajectory-family choice begins to matter.

9 Conflict of interest

The authors declare no conflicts of interest.

10 Funding

This work received no specific funding.

11 Data availability statement

The data and code that support the findings of this study are openly available in the `pmsimstats-ng` project repository. Per-replicate Monte Carlo outputs and ADEMP-compliant cell-level summaries are versioned under `analysis/data/`; simulation drivers are at `analysis/scripts/`. Exact paths for this manuscript are listed in the Reproducibility section below.

12 Reproducibility

This manuscript was rendered with R 4.4.x inside the project's `zzcollab` Docker container (image hash recorded in `Dockerfile`; package set captured in `renv.lock`). Principal packages used by the simulation pipeline are `nlme` (continuous-time linear-mixed analysis with `corCAR1` residual correlation), `parallel` (8-core `mclapply` for parameter-grid sweeps), `data.table` (the `original-extended` simulation collection), `furrr` and `purrr` (the `tidyverse` simulation collection), and `rmarkdown` plus `bookdown` for the manuscript build.

Random-number generator. Each simulation driver sets `set.seed(20260507)`

at the top of the replicate loop and uses `parallel::mc.set.seed` inside `mclapply`.
Per-replicate seeds derive from the master seed by `+ rep_idx`.

Data and code availability. Per-replicate simulation outputs are checkpointed
at `analysis/data/quick-sim/07-gompertz-replicates.rds`; Morris ADEMP
cell-level summaries at the companion `07-gompertz-summary.txt`. Driver scripts
are under `analysis/scripts/gompertz-evaluation/`. The pre-registered
ADEMP plan is at `analysis/scripts/gompertz-evaluation/00-ademp-pre-registration.md`.
The manuscript source, paper-specific bibliography (`references.bib`), and SIM
CSL file are at `analysis/report/07-gompertz-evaluation/`.

13 References

1. Hendrickson RC, Thomas RG, Schork NJ, Raskind MA. Optimizing aggregated N-of-1 trial designs for predictive biomarker validation: Statistical methods and practical considerations. *Frontiers in Digital Health*. 2020;2:13. doi:10.3389/fdgth.2020.00013
2. Laird AK. Dynamics of tumor growth. *British Journal of Cancer*. 1964;13(3):490-502. doi:10.1038/bjc.1964.55
3. Norton L. A Gompertzian model of human breast cancer growth. *Cancer Research*. 1988;48(24 Pt 1):7067-7071.
4. Zwietering MH, Jongenburger I, Rombouts FM, Riet K van 't. Modeling of the bacterial growth curve. *Applied and Environmental Microbiology*. 1990;56(6):1875-1881. doi:10.1128/aem.56.6.1875-1881.1990
5. Richards FJ. A flexible growth function for empirical use. *Journal of Experimental Botany*. 1959;10(2):290-301.
6. Tjørve E, Tjørve KMC. A unified approach to the Richards-model family for use in growth analyses: Why we need only two model forms. *Journal of Theoretical Biology*. 2010;267(3):417-425. doi:10.1016/j.jtbi.2010.09.008
7. Tjørve KMC, Tjørve E. The use of Gompertz models in growth analyses, and new Gompertz-model approach: An addition to the Unified-Richards family. *PLOS ONE*. 2017;12(6):e0178691. doi:10.1371/journal.pone.0178691
8. Lillie EO, Patay B, Diamant J, Issell B, Topol EJ, Schork NJ. The n-of-1 clinical trial: The ultimate strategy for individualizing medicine? *Personalized Medicine*. 2011;8(2):161-173. doi:10.2217/pme.11.7

- 452 9. Schork NJ, Beaulieu-Jones B, Liang WS, Smalley S, Goetz LH. Exploring human biology with N-of-1 clinical trials. *Cambridge Prisms: Precision Medicine*. 2023;1:e12. doi:10.1017/pcm.2022.15
- 453 10. Vohra S, Shamseer L, Sampson M, et al. CONSORT extension for reporting N-of-1 trials (CENT) 2015 statement. *Journal of Clinical Epidemiology*. 2015;76:9-17. doi:10.1016/j.jclinepi.2015.05.004
- 454 11. Benzekry S, Lamont C, Beheshti A, et al. Classical mathematical models for description and prediction of experimental tumor growth. *PLOS Computational Biology*. 2014;10(8):e1003800. doi:10.1371/journal.pcbi.1003800
- 455 12. Goutelle S, Maurin M, Rougier F, et al. The Hill equation: A review of its capabilities in pharmacological modelling. *Fundamental and Clinical Pharmacology*. 2008;22(6):633-648. doi:10.1111/j.1472-8206.2008.00633.x
- 456 13. Dayneka NL, Garg V, Jusko WJ. Comparison of four basic models of indirect pharmacodynamic responses. *Journal of Pharmacokinetics and Biopharmaceutics*. 1993;21(4):457-478. doi:10.1007/BF01061691
- 457 14. Bertalanffy L von. Quantitative laws in metabolism and growth. *Quarterly Review of Biology*. 1957;32(3):217-231.
- 458 15. Morris TP, White IR, Crowther MJ. Using simulation studies to evaluate statistical methods. *Statistics in Medicine*. 2019;38(11):2074-2102.
- 459 16. Jagadeesan P, Raman K, Tangirala AK. Sloppiness: Fundamental study, new formalism and its application in model assessment. *PLOS ONE*. 2023;18(3):e0282609. doi:10.1371/journal.pone.0282609
- 460 17. Hill AV. The possible effects of the aggregation of the molecules of haemoglobin on its dissociation curves. *Journal of Physiology*. 1910;40:iv-vii.
- 461 18. Raskind MA, Peterson K, Williams T, et al. A trial of prazosin for combat trauma PTSD with nightmares in active-duty soldiers returned from Iraq and Afghanistan. *American Journal of Psychiatry*. 2013;170(9):1003-1010. doi:10.1176/appi.ajp.2013.12081133
- 462 19. Ebling WF, Matsumoto Y, Levy G. Feasibility of effect-controlled clinical trials of drugs with pharmacodynamic hysteresis using sparse data. *Pharmaceutical Research*. 1996;13(12):1804-1810. doi:10.1023/a:1016072806164

- 463 20. Triggs EJ, Charles BG, Contin M, et al. Population pharmacokinetics and pharmacodynamics of oral levodopa in parkinsonian patients. *European Journal of Clinical Pharmacology*. 1996;51(1):59-67. doi:10.1007/s002280050161
- 464 21. Hockenberry MJ, Hooke MC, Rodgers C, et al. Symptom trajectories in children receiving treatment for leukemia: A latent class growth analysis with multitrajectory modeling. *Journal of Pain and Symptom Management*. 2017;54(1):1-8. doi:10.1016/j.jpainsymman.2017.03.002
- 465 22. Verhulst PF. Notice sur la loi que la population suit dans son accroissement. *Correspondance Mathématique et Physique*. 1838;10:113-121.
- 466 23. Gompertz B. On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. *Philosophical Transactions of the Royal Society of London*. 1825;115:513-583.
- 467 24. Winsor CP. The Gompertz curve as a growth curve. *Proceedings of the National Academy of Sciences USA*. 1932;18(1):1-8.
- 468 25. Muggeo VMR. Estimating regression models with unknown break-points. *Statistics in Medicine*. 2003;22(19):3055-3071. doi:10.1002/sim.1545