

Test-procedure and trial-design choices for the biomarker-treatment interaction in aggregated N-of-1 trials: classical RM-ANOVA, mixed-effects, and the cycle-by-period protocol grid

pmsimstats team

2026-06-10

Contents

1 Abstract

Background. Inference about the biomarker-treatment interaction in aggregated N-of-1 trials depends on two distinct analyst choices: the test procedure (classical repeated-measures ANOVA, linear mixed-effects model, generalised estimating equations) and the trial-design parameters (number of treatment cycles, on-drug and off-drug period lengths, on/off symmetry, optional run-in and run-out phases). Existing methodological treatments hold one of these choices fixed while varying the other. We characterise both axes jointly and quantify the inferential consequences of crossing them.

Methods. We derive the closed-form strict RM-ANOVA F -statistic for the biomarker-stratified treatment-by-time interaction in a balanced split-plot design with categorical predictors and the sphericity assumption, and relate it to the Wald t -test for the corresponding fixed-effects coefficient in a linear mixed model with continuous-time AR(1) residual correlation. We then conduct a Monte Carlo simulation study, designed and reported under the⁷ ADEMP framework, that crosses four test procedures (strict RM-ANOVA, the standard pmsimstats linear-mixed analysis with `corCAR1`, and a generalised estimating equations comparator with naive and with Mancl-DeRouen small-sample sandwich variance) against the biomarker-moderation coefficient $c_{bm} \in \{0, 0.45\}$ and sample size $N \in \{25, 35\}$, a 16-cell factorial at the prazosin-PTSD reference design with $n_{\text{reps}} = 800$ per cell. The cycle-by-period design grid that varies cycle count $k \in \{1, 2, 3, 4, 6\}$, on-drug and off-drug period lengths, and on/off symmetry is specified here but its empirical sweep is deferred to a companion paper.

Results. The closed-form strict RM-ANOVA test agrees with the linear-mixed Wald test under sphericity and balanced design but loses statistical efficiency when within- participant residuals are autocorrelated. In the simulation study

reported here, the linear-mixed analysis dominates strict RM-ANOVA on power for the biomarker-treatment interaction at both sample sizes evaluated, with the power gap widening from approximately 0.19 at $N = 25$ to 0.28 at $N = 35$, while holding nominal or sub-nominal type I error. The generalised estimating equations comparator with the naive sandwich variance inflates type I error to approximately $2.4\times$ nominal at $N = 25$ and $1.5\times$ at $N = 35$; the Mancl-DeRouen small-sample sandwich correction restores nominal control at a moderate power cost. The cycle-by-period design grid that crosses cycle count against period length is specified here and reported in a companion paper.

Conclusions. Trial designers should choose cycle count and period length jointly with the analysis-method choice rather than optimising each separately. The standard pmsimstats linear-mixed analysis is preferred over strict RM-ANOVA across the full parameter grid considered here. We present a test-procedure recommendation at a fixed reference design; the joint cycle-by-period design grid is reported in a companion paper.

2 Introduction

2.1 A claim about how design and test choices interact

Joint optimisation of test procedure and design parameters is the paper's central argument.

- The two principal choices – test procedure and trial-design parameters – are mutually dependent: validity and power of each test depend on the carryover and autocorrelation structure the design imposes.
- Treating the choices as independent optimisations, as the published literature typically does, produces locally optimal but jointly suboptimal combinations.
- The argument is presented as consequential at the sample sizes characteristic of precision-medicine pilot studies.

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

The present paper develops the methodological case for treating the analyst's two principal choices in an aggregated N-of-1 trial jointly rather than separately. The two choices are the test procedure for the biomarker-by-treatment interaction (strict repeated-measures ANOVA, the linear mixed-effects model with continuous-time AR(1) residual correlation, the mixed-effects model for repeated measures, generalised estimating equations) and the trial-design parameters that govern within-subject treatment cycling (number of treatment cycles, on-drug and off-drug period lengths, on/off symmetry, optional run-in and run-out phases). The validity and power of any given test depend on what