Summary of paper 01: tables and figures of the extended data-generating process

pmsimstats team

November 07, 2026

Problem and approach

Validating a predictive biomarker requires an interaction test, which needs about four times the sample of a main-effect test. Hendrickson et al. (2020) proposed a data-generating process (DGP) for simulating aggregated N-of-1 trials. In that original DGP the predictive biomarker is a correlation with the drug response that is present on drug and absent off drug. Paper 01 extends the original DGP for the Hybrid design with carryover. It judges each extension by five criteria: existence, identifiability under carryover, fidelity to the pharmacology, analyzability and calibration status. All simulations use the original response and baseline parameters of Hendrickson et al. This summary reproduces every table and figure of the paper in the paper's order. Each is generated by the same code as in the paper.

Methods

Table 1 states which analysis targets which quantity. The paired-difference statistic E9 is the one with a closed form.

Table 1: The analyses used in this paper and the quantity each estimates.

Analysis	Drug indicator	Estimates	Used in
Paired-difference statistic E9, E9b (closed form)	binary	on-minus-off contrast, $\beta_{bm:D}$ times the on/off correlation difference	Stages 1, 2, 4, 5; planning

Analysis	Drug indicator	Estimates	Used in
The original analysis (mixed model)	binary	the same contrast	Stage 2 (original code)
Stress test: the original analysis	binary	the on-minus-off contrast	Stage 6
Stress test: fixed-half-life analyses	$D_{bc,it}$ at an analysis half-life of 0.5 week, the true half-life except in the cells that vary it	on-drug moderation $\beta_{bm:D}$ where the half-lives match	Stage 6
Stress test: analyses with the half-life chosen by AIC	$D_{bc,it}$ at the AIC-chosen half-life (0, 0.25, 0.5, 1 or 2 weeks)	on-drug moderation $\beta_{bm:D}$ where the chosen half-life is the true one	Stage 6

Table 2 lays out the three simulation studies. They verify the closed forms, rerun the original code and stress-test ten analyses under a prognostic biomarker. A fourth run simulates the analysis G10 for the planning section.

Table 2: Design of the three simulation studies in the ADEMP structure. MCSE: Monte Carlo standard error of a rejection rate, $\sqrt{p(1-p)/R}$ for R replicates.

	1. Closed-form verification	2. The original code	3. Stress test under a prognostic biomarker
Aim	Check the closed-form moments and power of Appendix A, and the loss of power under the original rule	Show on the original code that the correlation rule alone determines the loss of power	Find which analyses keep their nominal size under features the extended DGP adds

	1. Closed-form verification	2. The original code	3. Stress test under a prognostic biomarker
Data-generating mechanisms	The correlation encoding with the original rule and with the decaying correlation; compound symmetry and the separable AR(1) form; carryover on the drug-response mean decaying from the last on-drug value; Hybrid, crossover and OL+BDC designs; $N = 70$; grids of c_{bm} and $t_{1/2}$	The original `generateData()` at commit 58b32a9, as published and with the decaying correlation only (Appendix B.1); $c_{bm} = 0.25$. (a) Hybrid, $N = 70$, no censoring, $t_{1/2} \in \{0, 0.1, 0.2, 1\}$; (b) four designs, $N \in \{35, 70\}$, $t_{1/2} \in \{0, 0.1, 0.2\}$, the original censoring patterns	The extended DGP (Table 12), run three times: the reported run as defined, the first with the original carryover rule and the second with randomized allocation; base cell separable AR(1), $\rho = 0.7$, $N = 70$, $t_{1/2} = 0.5$ week; 15 null cells (11 without and 4 with a prognostic pathway of strength 0.3) and 26 alternative cells: power curves under the separable form (c_{bm} from -0.30 to 0.45, 7 cells) and under compound symmetry ($\rho_{cs} = 0.8$, c_{bm} from -0.20 to 0.25, 6 cells), and 13 cells at $c_{bm} = 0.30$ that each vary one feature (N, half-life, decay form, dropout, measurement error or a prognostic pathway)
Estimand	$\beta_{bm:D}$, and the mean of the paired-difference statistic	$\beta_{bm:D}$	$\beta_{bm:D}$

	1. Closed-form verification	2. The original code	3. Stress test under a prognostic biomarker
Methods	Paired-difference statistics E9 and E9b, closed form and simulated; the original mixed model fitted with `lmer`	The original analysis pipeline (`lmer`, model-based test) on both versions	Fourteen analyses fitted to the same simulated trials, of which the ten in Table 9 are reported; two-sided tests at $\alpha = 0.05$
Performance measures	Mean, SD and power of the statistic; agreement as (simulated − closed form)/MCSE	Power; mean interaction estimate; number of repaired correlation matrices	Null rejection rate; power; failed fits
Replicates	5,000 per cell (MCSE of power at most 0.007); 500 to 1,000 for `lmer`	(a) 500 per cell (MCSE at most 0.022); (b) 250 per cell (at most 0.032)	1,000 per null cell (MCSE 0.0069 at a rate of 0.05), 500 per alternative cell (at most 0.022)

Results

Stage 1. Validity of the joint distribution

Table 3 counts correlation matrices that are not positive definite. The original code repaired 20 matrices (25%) on its own grid without a warning. The extended DGP stops instead, and it finds 75% of the grid infeasible at $c_{bm} = 0.6$.

Table 3: Correlation matrices that are not positive definite. Figure 4 grid: open label, open label plus blinded discontinuation, crossover and Hybrid designs; $t_{1/2} \in \{0, 0.1, 0.2\}$ week. Extended grid: the last three designs; $t_{1/2} \in \{0, 0.1, 0.2, 0.5, 1\}$ week. The original-DGP counts reproduce those of the original code exactly. The extended-DGP counts are the same at $\rho = 0.7$ (the recommended value) and $\rho = 0.8$ because both feasibility bounds lie between 0.3 and 0.6.

Grid (number of matrices)	Original DGP (constant correlation, compound symmetry, $\rho_{cs} = 0.8$)	Extended DGP (decaying correlation, separable AR(1)), $\rho = 0.8$	Extended DGP, $\rho = 0.7$
Figure 4 grid of the original study, $c_{bm} \in \{0, 0.3, 0.6\}$ (81)	20 (25%)	21 (26%)	21 (26%)
Extended grid, $c_{bm} = 0.25$ (40)	0 (0%)	0 (0%)	0 (0%)
Extended grid, $c_{bm} = 0.3$ (40)	12 (30%)	0 (0%)	0 (0%)
Extended grid, $c_{bm} = 0.6$ (40)	12 (30%)	30 (75%)	30 (75%)

Stage 2. The biomarker-response correlation under carryover

Table 4 gives closed-form power by half-life. Under the original rule E9 power falls to the test size at every positive half-life, and E9b keeps a little power from the baseline visit. Under the decaying rule it stays near its no-carryover value at 0.1 to 0.2 week and declines smoothly after.

Table 4: Closed-form power by carryover half-life (weeks). Hybrid design, $N = 70$, $c_{bm} = 0.25$, compound symmetry. Constant: the original rule (constant correlation after first exposure). Decaying: the extended rule (correlation decaying at the carryover half-life). E9: paired-difference statistic on post-baseline visits; E9b: the same with baseline counted as an off-drug visit, as in the original analysis. Against Monte Carlo estimates (5,000 replicates per cell), 2 of the 24 cell-variants in this table differ by more than 2 Monte Carlo SE, and 16 of all 566 cell-variants of study 1 (about 26 expected by chance).

Correlation rule	Statistic	0	0.1	0.2	0.5	1	2
Constant (original)	E9	0.721	0.050	0.050	0.050	0.050	0.050
	E9b	0.654	0.081	0.081	0.081	0.081	0.081
Decaying	E9	0.721	0.721	0.712	0.631	0.473	0.278
	E9b	0.654	0.653	0.647	0.586	0.472	0.327

Figure 1 shows the same result as lines, with Monte Carlo points on them.




Figure 1: Closed-form power (lines) against carryover half-life, with Monte Carlo estimates (points; 5,000 replicates per cell), under the original rule (constant correlation after first exposure) and the extended rule (correlation decaying at the carryover half-life). Hybrid design, $N = 70$, $c_{bm} = 0.25$, compound symmetry; left panel E9b, right panel E9.

Figure 2 plots power against the on/off correlation difference. Power depends on the correlation rule only through this difference.




Figure 2: Closed-form power against the on/off correlation difference, with the positions of the two correlation rules. Hybrid design, $N = 70$, $c_{bm} = 0.25$, E9.

Table 5 repeats the comparison in the original code, with only the correlation lines replaced. A rerun of the original version at 0.1 week with 4,000 new replicates gave power 0.090.

Table 5: The original code at commit 58b32a9, as published (original) and with the decaying correlation (decaying): Hybrid design, $N = 70$, $c_{bm} = 0.25$, 500 replicates per cell, same seeds. Monte Carlo SE of power at most 0.022. A rerun of the original version at 0.1 week with 4,000 new replicates gave power 0.090 (Stage 2 text).

$t_{1/2}$ (weeks)	Mean estimate, original	Mean estimate, decaying	Power, original	Power, decaying
0	-0.131	-0.131	0.668	0.668
0.1	-0.028	-0.129	0.126	0.638
0.2	-0.029	-0.131	0.092	0.656
1	-0.033	-0.107	0.094	0.462

Figure 3 repeats the comparison for all four designs, both sample sizes and five censoring patterns. The decaying correlation keeps power where the original rule loses it.




Figure 3: Power to detect the biomarker-by-treatment interaction, in the layout of Figure 4B of the original study, from the original code at commit 58b32a9 as published (upper two panel rows) and with the decaying correlation (lower two panel rows). Rows within panels: censoring patterns; columns: carryover half-life; panels: trial design (N-of-1 is the Hybrid design) and total sample size. $c_{bm} = 0.25$, 250 replicates per cell (Monte Carlo SE at most 0.032), the same seeds for both versions. No matrix was repaired in either version.

Figure 4 repeats the grid under the extended DGP with the recommended analysis G10 (a random-intercept CAR(1) mixed model with an analysis half-life chosen by AIC on the model without the interaction, and CR2 standard errors). At $N = 70$ without censoring, Hybrid power is 0.34, 0.38 and 0.34 at half-lives of 0, 0.1 and 0.2 week. The open-label design has no off-drug visit, so the

interaction is not estimable under G10.




Figure 4: Power of the analysis G10 to detect the biomarker-by-treatment interaction under the extended DGP (separable AR(1), $\rho = 0.7$, $c_1 = 0.2$, decaying correlation, carryover decaying from the last on-drug mean, original response parameters), in the layout of Figure 4B of the original study. Rows within panels: censoring patterns; columns: carryover half-life; panels: trial design (N-of-1 is the Hybrid design) and total sample size. $c_{bm} = 0.25$, 250 replicates per cell (Monte Carlo SE at most 0.031). Censoring patterns use the corrected dropout rule, under which dropout rises with worsening; the original-code figure uses the published rule of commit 58b32a9. Gray cells (n/e): the open-label design, in which every visit is on drug and the interaction is not estimable under G10.

Figure 5 extends the closed-form comparison over interaction strengths and longer half-lives. Gray cells mark effect sizes with no valid joint distribution.




Figure 5: Closed-form power of the paired-difference statistic by interaction strength (c_{bm}) and carryover half-life, in the layout of Figure 4B of the original study: the original rule and the decaying correlation. Upper row: E9b, which counts baseline as an off-drug visit as the original analysis does; lower row: E9. Gray cells: c_{bm} at or above the compound-symmetry feasibility bound, where the correlation matrix is not positive definite. Hybrid design, $N = 70$, compound symmetry.

Stage 3. Carryover on the mean

Figure 6 shows the drug-response mean by visit for the four Hybrid paths under both carryover rules.




Figure 6: Drug-response mean by visit in the Hybrid design, by path, under the extended carryover rule (decay from the last on-drug mean, the extended DGP) and the original carryover rule (compounded decay), at carryover half-lives of 0.2 and 1 week. Filled points: on-drug visits; open points: off-drug visits. D: drug; P: placebo.

Stage 4. Serial correlation

Table 6 shows what the serial correlation does to the feasibility bound and to power.

Table 6: Serial correlation: the feasibility bound on c_{bm} (Ceiling; decaying correlation, no carryover, minimum over the Hybrid, crossover and OL+BDC designs) and closed-form E9 power (Hybrid, $N = 70$, no carryover) by ρ, under compound symmetry and under the separable AR(1) form of Stage 5. Under compound symmetry the Hybrid and crossover designs tie for the minimum. Under the separable form the minimum is attained in the Hybrid design at $\rho = 0.8$ and in the OL+BDC design at $\rho \leq 0.7$, and the crossover design is never binding. At $t_{1/2} = 0.2$ week every power value is lower by 0.004 to 0.009.

	Compound symmetry		Separable AR(1)	
ρ_{cs} or ρ	Ceiling	Power at $c_{bm} = 0.25$	Ceiling	Power at $c_{bm} = 0.25$
0.5	0.458	0.492	0.448	0.261
0.6	0.410	0.553	0.462	0.270
0.7	0.346	0.628	0.474	0.290
0.8	0.256	0.721	0.440	0.335

Table 7 extends the separable form to ρ from 0.4 to 0.95. The bound is largest at 0.7 and falls steeply above it.

Table 7: Serial correlation under the separable AR(1) form of the extended DGP ($c_1 = 0.2$, decaying correlation, carryover decaying from the last on-drug mean): the feasibility bound on c_{bm}, minimum over the Hybrid, crossover and OL+BDC designs, by carryover half-life, the design that attains the minimum without carryover, and the closed-form total N for 80 percent power of E9 in the Hybrid design at the 0.2-week half-life. Not feasible: c_{bm} lies above the Hybrid bound.

	Feasibility bound				Hybrid N, E9	
ρ	$t_{1/2} = 0$	0.2 week	1 week	Binding design	$c_{bm} = 0.25$	$c_{bm} = 0.35$
0.40	0.437	0.438	0.445	OL+BDC	318	160
0.50	0.448	0.450	0.465	OL+BDC	314	158
0.60	0.462	0.465	0.494	OL+BDC	302	152
0.65	0.469	0.473	0.512	OL+BDC	292	148
0.70	0.474	0.479	0.533	OL+BDC	278	140
0.75	0.465	0.471	0.550	Hybrid	258	130
0.80	0.440	0.446	0.544	Hybrid	234	118
0.85	0.400	0.407	0.519	Hybrid	200	100
0.90	0.340	0.346	0.457	Hybrid	154	not feasible
0.95	0.247	0.251	0.343	Hybrid	92	not feasible

Stage 5. Cross-component structure

Table 8 compares three correlation structures.

Table 8: Largest feasible biomarker-response correlation c_{bm} under the decaying correlation, without carryover, minimum over the Hybrid, crossover and OL+BDC designs. Under compound symmetry the Hybrid and crossover designs tie for the minimum. Under the two AR(1) forms the minimum is attained in the Hybrid design at $\rho = 0.8$ and in the OL+BDC design at $\rho = 0.7$ (separable form: Hybrid 0.480), and the crossover design is never binding.

ρ_{cs} or ρ	Compound symmetry	AR(1), two-constant cross-component	Separable AR(1)
0.8	0.256	0.291	0.440
0.7	0.346	0.452	0.474

Stage 6. Features the extended DGP adds

Table 9 gives null rejection rates under a prognostic biomarker in the third run of study 3, the extended DGP with fixed allocation. The first run, with the original carryover rule, is in Table 17.

Table 9: Null rejection rates under a prognostic biomarker ($c_{bm} = 0$). Each pathway has strength 0.3: a correlation of 0.3 with the natural-history component (constant), the placebo-response component or baseline severity, and for the growing natural-history pathway a mean shift of the natural-history component of 0.3 of its SD (3 points) per SD of biomarker at the end of its curve, growing along the curve. 1,000 replicates per cell (two-MCSE band 0.036 to 0.064 for a single cell). Third run of study 3: the extended DGP, with carryover decaying from the last on-drug mean and fixed path allocation. Rates above 0.075 in bold. The first run is in Table 17; the changes between runs are given in the text.

Analysis	Natural history, constant	Natural history, growing	Placebo response	Baseline severity
Original analysis (random intercept, model-based)	**0.108**	0.056	**0.157**	0.064
Random intercept + CAR(1), CR2	**0.103**	0.049	**0.105**	0.052
The same, half-life chosen by AIC	**0.130**	0.059	**0.125**	0.057
Unstructured covariance, Kenward-Roger	**0.413**	**0.200**	**0.224**	0.048
The same, half-life chosen by AIC	**0.506**	**0.258**	**0.270**	0.054
Random intercept + CAR(1) with expectancy terms, CR2	0.045	0.040	0.052	0.055
The same, with phase-specific residual variances, CR2	0.056	0.037	0.051	0.059
The same, with phase-specific residual variances, model-based	0.054	0.042	0.049	0.059
Expectancy terms, half-life chosen by AIC, CR2	0.058	0.050	0.060	0.060
Random intercept + CAR(1) with free baseline slope, CR2	0.051	0.066	0.060	0.054

Table 10 gives the feasibility bounds when a prognostic correlation shares the biomarker row with c_{bm}. Each correlation lowers the room left for the other.

Table 10: Largest feasible prognostic correlation given the biomarker-response correlation, and the reverse, under the extended DGP (separable AR(1), $\rho = 0.7$, $c_1 = 0.2$), Hybrid design, minimum over its four paths. Correlation: the quantity whose bound is given. c_{PB}, c_{TV}, c_{BL}: correlation of the biomarker with the placebo-response component, the natural-history component and baseline severity. None: no value is feasible.

Correlation	Given	$t_{1/2} = 0.2$ week	$t_{1/2} = 0.5$ week
c_{PB} or c_{TV}	alone	0.513	0.513
	$c_{bm} = 0.25$	0.466	0.477
	$c_{bm} = 0.30$	0.435	0.451
	$c_{bm} = 0.45$	0.243	0.306
c_{BL}	alone	1.000	1.000
	$c_{bm} = 0.25$	0.857	0.876
	$c_{bm} = 0.30$	0.786	0.815
	$c_{bm} = 0.45$	0.374	0.496
c_{bm}	alone	0.485	0.518
	$c_{PB} = 0.3$	0.422	0.454
	$c_{PB} = 0.6$	none	none

Table 11 shows what each optional feature does to size and power.

Table 11: The optional features of the extended DGP: null rejection rate ($c_{bm} = 0$; 1,000 replicates, MCSE 0.007) and power ($c_{bm} = 0.30$; 500 replicates, MCSE at most 0.022) for the original analysis (Original), the random-intercept CAR(1) model with CR2 and the expectancy-term model with CR2 (Expect.). Hybrid design, third run of study 3 (the extended DGP: carryover decaying from the last on-drug mean, fixed allocation). Each row changes one feature of the base cell.

	Null rejection			Power		
Cell	Original	CAR(1)	Expect.	Original	CAR(1)	Expect.
Base cell ($N = 70$, $t_{1/2} = 0.5$, exponential, no dropout)	0.077	0.051	0.067	0.474	0.486	0.404
Measurement error (SD 3 and 6)	0.069	0.050	0.037	0.422	0.394	0.314
Dropout 20 percent, completely at random	0.078	0.053	0.055	0.466	0.464	0.392
Dropout 20 percent, dependent on last score	0.069	0.060	0.047	0.424	0.426	0.360
Weibull decay, shape 0.5	0.072	0.046	0.044	0.378	0.400	0.300
Weibull decay, shape 2	0.066	0.039	0.046	0.526	0.540	0.496
$t_{1/2} = 0$	0.071	0.040	0.041	0.536	0.566	0.498
$t_{1/2} = 1$ week	0.066	0.047	0.061	0.418	0.416	0.348
$N = 35$	0.077	0.053	0.048	0.270	0.246	0.206
$N = 150$	0.060	0.048	0.045	0.768	0.790	0.688

Stage 7. The extended DGP

Table 12 collects the choices. The Kind column marks each element as an extension, a choice, an option or a retained element.

Table 12: The extended data-generating process. The Default column gives the value of each element in the extended DGP; where the stress test of Stage 6 used another value it is stated. The stress test's reported run uses fixed allocation; its second run assigned participants to paths at random, a driver option outside the extended DGP (Appendix B.5).

Element	Specification	Default	Kind	Justified in
Components	$Y_{it} = \mathrm{BL}_i - (TV_{it} + PB_{it} + BR_{it})$, Gompertz means as in the original DGP	original parameters (Methods): BR $m = 10.99$, $r = 0.42$, SD 8; PB and TV $m = 6.51$, $r = 0.35$, SD $10\eta_t$ and 10; $d = 5$; biomarker 124.3 (SD 15.4); baseline 83.1 (SD 18.5)	retained	Methods
Validity	stop if the correlation matrix is not positive definite	always	extension	Stage 1
Biomarker-response correlation	c_{bm} on drug; $c_{bm}2^{-t_{sd}/t_{1/2}}$ off drug after exposure; 0 before exposure	c_{bm} below the feasibility bound of every design in the study	extension	Stage 2
Carryover on the mean	decay from the last on-drug value, $\mu_{BR,\mathrm{last}}\, 2^{-t_{sd}/t_{1/2}}$	$t_{1/2} = 0.1$ to 0.2 week (pharmacological, assumed); 0.5 in the stress test	choice	Stage 3

Element	Specification	Default	Kind	Justified in
Re-exposure	as in the original DGP: time on drug restarts at zero	on, for titrated drugs	retained	Stage 3
Within-component correlation	AR(1) in weeks, $\rho^{\lvert w_t - w_s \rvert}$, one ρ for all three components	$\rho = 0.7$ (convention; report and vary)	choice	Stage 4
Cross-component correlation	separable, $c_1\rho^{\lvert w_t - w_s \rvert}$ at every lag; the Kronecker form applies to the correlation, since the PB SD scales with the expectancy	$c_1 = 0.2$ (original value)	choice	Stage 5
Path allocation	fixed numbers, as in the original DGP	original values	retained	Methods
Prognostic correlations	correlation of the biomarker with TV or PB, or with baseline severity; or a mean shift of TV growing along its curve	off; 0.3 in the stress test; correlations must lie inside the joint bound with c_{bm} (Table 10)	option	Stage 6
Measurement error	independent, SD by phase	off; 3 open label, 6 blinded in the stress test	option (driver only)	Stage 6
Decay form	exponential or Weibull with the same half-life	exponential	option	Stage 6
Dropout	monotone, completely at random or at random given the last score	off; 20% in the stress test	option	Stage 6

Using the extended DGP to plan a trial

Table 13 gives the total sample size for 80 percent power of the paired-difference statistic E9, and the simulated power of the recommended analysis G10 at $N = 70$. Table 14 gives G10 in every cell. G10 rejected the null at 0.046 to 0.053. At the E9 sample size for 80 percent power, G10 power was 0.844 to 0.936 in the 20 uncapped cells, so the closed-form E9 rule is conservative for G10 in these cells. Table 15 gives the G10 sample size for 80 percent power in the Hybrid design. It is 0.71 to 0.83 times the E9 value.

Table 13: Planning under the extended DGP (separable AR(1), $\rho = 0.7$, $c_1 = 0.2$, decaying correlation, carryover decaying from the last on-drug mean, original response parameters): total N for 80 percent power of the paired-difference statistic E9 at $\alpha = 0.05$, by design, interaction strength and carryover half-life, with the implied slope in symptom points per SD of biomarker ($\sigma_{BR} = 8$) and the power at $N = 70$ and the 0.2-week half-life for E9 (closed form) and for the analysis G10 (simulated, 1,000 replicates per cell, Monte Carlo SE at most 0.016). Participants are allocated to paths in the original fixed proportions.

			N for 80 percent power			Power at $N = 70$, 0.2 week	
Design	c_{bm}	Slope (points per SD)	$t_{1/2} = 0$	0.2 week	1 week	E9	G10
Hybrid	0.25	2.0	272	278	480	0.285	0.357
	0.35	2.8	138	140	244	0.503	0.610
	0.45	3.6	82	84	146	0.728	0.836
OL+BDC	0.25	2.0	394	404	> 800	0.210	0.275
	0.35	2.8	200	206	452	0.368	0.507
	0.45	3.6	120	124	272	0.558	0.722
Crossover	0.25	2.0	424	424	454	0.202	0.243
	0.35	2.8	216	216	230	0.353	0.365
	0.45	3.6	130	130	138	0.537	0.592

Table 14: Rejection rate of the analysis G10 under the extended DGP of the preceding planning table, by design, interaction strength and carryover half-life, at $N = 70$ and at the total N the closed-form E9 rule gives for 80 percent power (the preceding table). Rows with $c_{bm} = 0$ give the size at $N = 70$ (2,000 replicates per cell, Monte Carlo SE at most 0.005); the other rows give power (1,000 replicates per cell, Monte Carlo SE at most 0.016). †: the E9 sample size exceeds 400 and the cell ran at $N = 400$. Rejection rates are over converged fits (at most 0.8 percent of replicates failed in any cell).

		At $N = 70$			At the E9 N for 80 percent power		
Design	c_{bm}	$t_{1/2} = 0$	0.2 week	1 week	$t_{1/2} = 0$	0.2 week	1 week
Hybrid	0.00	0.046	0.051	0.048			
	0.25	0.367	0.357	0.217	0.915	0.900	0.809†
	0.35	0.616	0.610	0.362	0.908	0.899	0.885
	0.45	0.855	0.836	0.531	0.894	0.903	0.874
OL+BDC	0.00	0.052	0.051	0.051			
	0.25	0.295	0.275	0.133	0.936	0.915†	0.601†
	0.35	0.495	0.507	0.235	0.934	0.936	0.866†
	0.45	0.725	0.722	0.362	0.930	0.932	0.891
Crossover	0.00	0.053	0.046	0.050			
	0.25	0.187	0.243	0.220	0.842†	0.840†	0.812†
	0.35	0.356	0.365	0.378	0.861	0.867	0.862
	0.45	0.594	0.592	0.551	0.860	0.869	0.844

Table 15: Total N for 80 percent power in the Hybrid design under the extended DGP with the original parameters: closed-form E9 (Table 13) and the analysis G10. G10 was simulated at 6 values of N per cell (500 replicates each), and the N at which a probit regression of rejection on $\log N$ reaches 0.80 is rounded up.

c_{bm}	$t_{1/2}$ (weeks)	E9 (closed form)	G10 (simulated)	G10 / E9	Simulated N
0.25	0	272	193	0.71	95 to 340
0.25	0.2	278	205	0.73	97 to 348
0.25	1	480	384	0.80	168 to 600
0.35	0	138	104	0.75	48 to 172
0.35	0.2	140	108	0.77	49 to 175
0.35	1	244	193	0.79	85 to 305
0.45	0	82	61	0.74	29 to 102
0.45	0.2	84	64	0.76	29 to 105
0.45	1	146	121	0.83	51 to 182

Table 16 repeats the planning cells at the 0.2-week half-life for $\rho = 0.5$ and 0.9. At 0.9 the Hybrid design needs about half the sample, and 8 to 12 percent of G10 fits failed in each crossover cell at $N = 70$, because the fitted serial correlation reached its limit of 1 and could not be told apart from the random intercept.

Table 16: Planning by serial correlation at the 0.2-week half-life under the extended DGP (separable AR(1), $c_1 = 0.2$, original response parameters): the closed-form total N for 80 percent power of E9, and G10 power at $N = 70$ and at that N, with the percent of G10 fits that failed at $N = 70$. 1,000 replicates per cell. The rows at $\rho = 0.7$ repeat Table 13. At $\rho = 0.9$ the grid of c_{bm} lies inside the lower bound. Not feasible: c_{bm} lies above the bound of that design. †: the E9 sample size exceeds 400 and the cell ran at $N = 400$. Power is over the fits that succeeded.

				G10 power		
ρ	Design	c_{bm}	E9 N	At $N = 70$	At the E9 N	Failed (%)
0.5	Hybrid	0.25	314	0.258	0.849	0
		0.35	158	0.487	0.829	0
		0.45	96	0.693	0.822	0
	OL+BDC	0.25	412	0.225	0.844†	0
		0.35	208	0.391	0.859	0
		0.45	126	0.606	0.856	0
	Crossover	0.25	348	0.267	0.864	0
		0.35	176	0.405	0.825	0
		0.45	106	0.623	0.815	0
0.7	Hybrid	0.25	278	0.357	0.900	0
		0.35	140	0.610	0.899	0
		0.45	84	0.836	0.903	0
	OL+BDC	0.25	404	0.275	0.915†	0
		0.35	206	0.507	0.936	0
		0.45	124	0.722	0.932	0
	Crossover	0.25	424	0.243	0.840†	0
		0.35	216	0.365	0.867	0
		0.45	130	0.592	0.869	0
0.9	Hybrid	0.20	242	0.546	0.979	0
		0.25	154	0.751	0.976	0
		0.30	106	0.897	0.979	0
	OL+BDC	0.20	462	0.416	0.992†	0
		0.25	296	0.627	0.999	0
		0.30	204	0.757	0.997	0
	Crossover	0.20	628	0.212	0.803†	10
		0.25	402	0.315	0.944†	8
		0.30	278	0.451	0.938	9

Appendix A. Closed-form derivations

A.5 The separable structure

Each step between visits contributes the following term to the schedule term of the feasibility bound.

Step	Term
on to on	$(1-\rho_t)/(1+\rho_t)$
on to off	$\rho_t^2/(1-\rho_t^2)$
off to on	$1/(1-\rho_t^2)$
off to off	0

Appendix C. First run of the stress test

Table 17: Null rejection rates under a prognostic biomarker in the first run of study 3 (original carryover rule, fixed allocation), 1,000 replicates per cell (two-MCSE band 0.036 to 0.064 for a single cell). Rates above 0.075 in bold.

Analysis	Natural history, constant	Natural history, growing	Placebo response	Baseline severity
Original analysis (random intercept, model-based)	**0.107**	0.053	**0.159**	0.075
Random intercept + CAR(1), CR2	**0.099**	0.049	**0.100**	0.050
The same, half-life chosen by AIC	**0.115**	0.058	**0.112**	0.055
Unstructured covariance, Kenward-Roger	**0.416**	**0.195**	**0.222**	0.051
The same, half-life chosen by AIC	**0.509**	**0.245**	**0.256**	0.064
Random intercept + CAR(1) with expectancy terms, CR2	0.047	0.046	0.040	0.040
The same, with phase-specific residual variances, CR2	0.055	0.044	0.042	0.042
The same, with phase-specific residual variances, model-based	0.062	0.043	0.045	0.046
Expectancy terms, half-life chosen by AIC, CR2	0.059	0.052	0.047	0.045
Random intercept + CAR(1) with free baseline slope, CR2	0.049	0.063	0.049	0.050

Conclusions

The correlation encoding of the interaction is coherent. The original DGP holds the biomarker-response correlation constant after first exposure, so with any carryover the interaction is absent among the post-baseline visits and power falls nearly to the test size. The extended DGP lets the correlation decay after discontinuation, and power returns to its no-carryover value at pharmacological half-lives. The original effect size of 0.6 relied on repaired matrices, and a separable AR(1) structure at serial correlation 0.7 raises the feasibility bound to 0.474. At the closed-form sample size, G10 power was 0.844 to 0.936 in the uncapped cells. We propose the extended DGP, with a stop on infeasible matrices, as the reference for simulation studies of these designs.