Why Architecture B Departs from the Published Implementation

2026-09-09 10:57 PDT

Author. pmsimstats team

Purpose. This paper takes the differences between the covariance-moderation process of this package (`covar`) and the published Hendrickson et al. [1] implementation (`orig`) one at a time, and asks of each whether it repairs a defect, is a defensible modeling choice, or follows from another change. It supports the comparison of the two processes in paper 01.

Review note (2026-10-07). Three later results bear on this paper. (a) The ceilings quoted in Sections 3.3 and 4.3 were computed at $\rho = 0.7$; the published code uses $\rho = 0.8$, at which the published construct's ceiling is 0.256 (Hybrid and CO) and 0.262 (OL+BDC) without carryover (`docs/36`, Section 4.2; recomputed in this review). Under the separable AR(1) form of `docs/35` the ceiling is 0.440 at $\rho = 0.8$, and 0.474 (Hybrid 0.480) at $\rho = 0.7$. (b) `docs/37` shows that under the published step coupling the on-drug versus off-drug coupling gap is zero, so the interaction vanishes at *any* positive half-life; graded coupling, $g_t = 2^{-t_{sd}/t_{1/2}}$ off drug and 0 before first exposure, removes the collapse. (c) Paper 01 has since been restructured: the section numbers cited here for it (2.2.4, B.5 to B.7) refer to its September 2026 draft. Its current Appendix B lists the code changes to `58b32a9` (graded coupling, stop instead of repair, anchored carryover). A copy of `58b32a9` is now held at `analysis/data/quick-sim/hendrickson-58b32a9/`.

Contents

Notation index and glossary 3
1. Purpose 4
2. The four differences 4
3. The coupling vector: a defect, not a preference 4
3.1 What the published gate does 5
3.2 Why this is fatal rather than merely inaccurate 5
3.3 The symptom in the feasible range 5
3.4 The replacement 6
4. The within-factor form: a modeling choice we would defend 7
4.1 What compound symmetry asserts 7
4.2 What each form implies about the patient 7
4.3 The numerical consequence, compared like with like 8
4.4 What we do not claim 8
5. The cross-factor form: a consequence, not an independent choice . . 8
5.1 The incoherence of mixing the two 9
5.2 The numerical signature 9
5.3 Our position 9
6. The carryover recursion: our correction, not architecture 9
7. What the repository's later history does and does not settle 10
8. The exact ceiling used above 11
9. Conclusions 11
References 12

List of Tables

1 Notation index 3
2 Four differences between `orig` and `covar` by covariance component 4
3 Ceiling on c_{bm} for `orig` and `covar` on an evenly spaced eight-occasion schedule, $\rho_c = 0.7$ 6
4 AR(1) and compound-symmetry correlations for selected Hybrid visit pairs 7

Notation index and glossary

Notation follows `analysis/report/NOTATION.md`. Symbols marked *local* are defined here and are not part of the canonical set.

Table 1: Notation index

Symbol	Meaning	Status
t, s	measurement-occasion indices (s local)	canonical
n	number of measurement occasions on a path	local
w_t	cumulative calendar week of occasion t	local
B, BR, PB, TV	biomarker and the three response components	canonical
c_{bm}	covariance-moderation correlation: on-drug correlation of B with BR	canonical
ρ, ρ_c	AR(1) or compound-symmetry within-factor correlation, common or for component c (subscript local)	canonical
A_c	within-factor correlation block of component c	local
$C_{cc'}$	cross-factor correlation block between components c and c'	local
c_1	same-occasion cross-factor correlation (code `c.cf1t`)	local
$c_\times$	cross-factor correlation at distinct occasions (code `c.cfct`)	local
$\tilde{r}$, $\tilde{r}_t$	biomarker coupling vector: correlations of B with BR_t	local
v	coupling pattern, $\tilde{r} = c_{bm} v$	local
g_t	graded coupling factor off drug, $2^{-t_{sd}/t_{1/2}} = e^{-\lambda t_{sd}}$ (`docs/37`)	local
$\mathbf{1}_n$	vector of n ones	local
M	correlation block of the three response factors	local
c^*_{bm}	ceiling: largest c_{bm} with a positive definite matrix	local
t_{sd}, $t_{sd,t}$	time since discontinuation (at occasion t)	canonical
$t_{1/2}$	carryover half-life, weeks	canonical
λ	carryover decay rate, $\ln 2 / t_{1/2}$	canonical
μ_{BR}	mean of the drug-response component	local

Glossary. *`orig`*: the published implementation at commit `58b32a9`. *`covar`*: the package's covariance-moderation process (`dgp_architecture = 'mvn'`, formerly Architecture B). *Step coupling*: $\tilde{r}_t = c_{bm}$ wherever $\mu_{BR} \neq 0$, else 0. *Graded coupling*: $\tilde{r}_t = c_{bm}$ on drug, $c_{bm} e^{-\lambda t_{sd,t}}$ off drug after exposure, 0 before first exposure. *Compound symmetry (CS)*: one constant correlation for every pair of occasions. *Ceiling*: c^*_{bm} above. *Recursive* versus *anchored* carryover: decay of the previous, already decayed mean versus decay from the mean at discontinuation.

1. Purpose

Architecture B, which we call `covar` in code and in the appendices of paper 01, is not the covariance-moderation process that Hendrickson et al. [1] published. It differs from the published implementation in three places, and a fourth difference has since been introduced by a correction of our own. A reader who notices this is entitled to ask whether each departure was necessary or merely preferred.

We take that question seriously enough to answer it one difference at a time. Our conclusion is that the three differ in kind. One repairs a defect that would otherwise make the interaction unidentifiable under carryover. One is a modeling choice which we believe is clearly better, though the published alternative is defensible. One is a consequence of the second, and would be difficult to justify on its own.

Throughout, `orig` denotes the published implementation at commit `58b32a9` of 19 November 2020, the last commit before the 2024 revisions. (An earlier version of this sentence called it the last commit preceding publication. The article appeared on 28 August 2020 [1], so `58b32a9` postdates it; its data-generating code is identical to that of the initial commit `42ac030`, per `docs/36`, Section 2.) Section 7 records what happened to that repository afterward, and why its later history does not settle anything.

2. The four differences

Table 2: Four differences between `orig` and `covar` by covariance component

Component	`orig`	`covar`
Within-factor A_c	ρ_c	$\rho_c^{\lvert w_t - w_s \rvert}$
Cross-factor $C_{cc'}$, $t \neq s$	$c_\times$	$c_\times \rho^{\lvert w_t - w_s \rvert}$
Coupling $\tilde{r}_t$, off drug	0 or c_{bm}	$c_{bm} e^{-\lambda t_{sd,t}}$
Carryover on μ_{BR}	recursive	anchored

Everything else agrees. The same-occasion cross-factor entry, the on-drug value of the coupling vector, the baseline row and column, and the biomarker's zero correlation with the two non-response components are common to both.

Sections 3 through 6 take the four in order of how strongly we would defend them.

3. The coupling vector: a defect, not a preference

This is the one substantive repair, and we would make it even if nothing else were at stake.

3.1 What the published gate does

The published code sets the biomarker-response correlation by asking whether the response mean at that occasion is nonzero. Where it is, the correlation takes the full value c_{bm}; where it is not, zero. That is a sensible-looking rule. Its difficulty appears only once carryover is switched on.

Carryover keeps the response mean nonzero after discontinuation. That is what carryover is. So the gate, which asks whether any drug effect is present rather than how much, returns yes at off-drug occasions too, and assigns them the same c_{bm} it assigns on-drug occasions.

In the Hybrid design at any positive half-life (one week is the example used below), the result is

$$\tilde{r}^{\text{orig}} = c_{bm}\mathbf{1}_n,$$

a constant vector, because every Hybrid path begins on drug, so every off-drug occasion follows exposure and carries a nonzero residual mean (inspected, `58b32a9` `R/generateData.R`, lines 83 to 92 and 131 to 138). Every occasion, on drug and off, carries the same biomarker-response correlation.

3.2 Why this is fatal rather than merely inaccurate

The biomarker-treatment interaction is identified by $\tilde{r}$ varying between on-drug and off-drug occasions. That variation is the whole signal. A constant $\tilde{r}$ describes a biomarker that correlates with the response equally whether or not the patient is taking the drug, which is a biomarker main effect and not an interaction at all.

The consequence runs the wrong way round. Under the published gate, any carryover at all flattens $\tilde{r}$, so at every positive half-life the interaction is not identifiable through the coupling channel. The loss is a jump at $t_{1/2} = 0$, not a gradual decline (derived, `docs/37`; observed in the published results, `docs/36`, Section 5). But carryover ought to blur the on-drug and off-drug contrast, not equalize it at full strength. A patient two weeks off drug should look partly like a treated patient and partly like an untreated one. The published gate says they look exactly like a treated patient.

3.3 The symptom in the feasible range

The defect leaves a numerical fingerprint which is worth recording, because it is initially misleading. Evaluating the exact ceiling on the moderation parameter (section 8) on an evenly spaced schedule of eight occasions two weeks apart, on drug at the first four and off at the last four, with $\rho_c = 0.7$, $c_1 = 0.2$

and $c_\times = 0.1$ (verified: recomputed from the closed form of section 8 in the 2026-10-07 review):

Table 3: Ceiling on c_{bm} for `orig` and `covar` on an evenly spaced eight-occasion schedule, $\rho_c = 0.7$

Configuration	Ceiling on c_{bm}
`orig`, $t_{1/2} = 0$	0.346
`orig`, $t_{1/2} > 0$	0.841
`covar`, $t_{1/2} = 1$	0.670

At the published $\rho_c = 0.8$ the `orig` values on this schedule are 0.256 and 0.893, and on the paper 01 Hybrid design they are the same, 0.256 and 0.893 (verified; `docs/36`, Section 4.2). The values in the table are not those of the published parameters.

At positive carryover the published process admits a *larger* nominal moderation than Architecture B does. That looks like an advantage and is the opposite. A constant coupling vector aligns with the direction in which the covariance is best conditioned, so the matrix will tolerate a large c_{bm} precisely because that c_{bm} is no longer buying an interaction. A process that can carry a bigger number while carrying no signal is not thereby preferable.

3.4 The replacement

Architecture B gates on the drug indicator itself and decays with time since discontinuation,

$$\tilde{r}_t = \begin{cases} c_{bm}, & \text{on drug} \\ c_{bm} e^{-\lambda t_{sd,t}}, & \text{off drug after exposure,} \\ 0, & \text{before first exposure,} \end{cases} \qquad \lambda = \ln 2 / t_{1/2}.$$

In the same Hybrid cell ($c_{bm} = 0.45$, $t_{1/2} = 1$) this gives 0.45 on drug against 0.225 and 0.113 at one and two weeks off (verified). The contrast survives, and it narrows as residual exposure decays, which is what the phenomenon being modeled actually does.

We should be clear about the standing of this change. It is not a recovery of what the published code was trying to do; it is a departure from what the published code does. Our warrant is Hendrickson et al.'s own stated rationale [1], that the biomarker is informative because it shares variance with drug-responsive physiology. A coupling that persists undiminished through a washout does not express that rationale.

4. The within-factor form: a modeling choice we would defend

Compound symmetry assigns one constant to every pair of occasions within a factor, whatever their separation. Architecture B uses AR(1) decay in cumulative calendar time.

4.1 What compound symmetry asserts

Take the Hybrid schedule, whose occasions fall at weeks $4, 8, 9, 10, 11, 12, 16, 20$. With $\rho_c = 0.7$:

Table 4: AR(1) and compound-symmetry correlations for selected Hybrid visit pairs

Pair	Gap	AR(1)	Compound symmetry
week 9 vs 10	1 wk	0.700	0.70
week 4 vs 8	4 wk	0.240	0.70
week 4 vs 20	16 wk	0.003	0.70

(Derived: 0.7^1, 0.7^4 and 0.7^{16}.)

Compound symmetry asserts that a symptom measurement in week 4 resembles one taken sixteen weeks later exactly as strongly as it resembles one taken the following week. Under AR(1) the near pair is some two hundred times more strongly correlated than the far pair; under compound symmetry the ratio is one.

For a fluctuating condition measured over five months, we do not find the compound-symmetric claim credible. Symptoms persist and then drift. A patient's state next week is largely their state this week; their state next season is not.

4.2 What each form implies about the patient

The two forms encode different accounts of why a patient's measurements resemble one another.

Compound symmetry is the correlation a participant random intercept induces. It says the only thing making a patient's occasions alike is a fixed patient-level offset, present equally at every visit and unchanging. That is a real phenomenon, and in a trial with two or three widely spaced visits it may be the dominant one.

AR(1) adds the thing that serial measurement is for. States persist and decay. A bad week tends to be followed by a bad week and not by a bad month five months later.

An N-of-1 trial measures the same patient eight times over twenty weeks, with some occasions a week apart and others four. It exists in order to exploit within-patient temporal structure. A correlation model that flattens that structure discards the feature the design was built to use.

4.3 The numerical consequence, compared like with like

Holding the coupling vector fixed (graded, $t_{1/2} = 1$) on the eight-occasion schedule of section 3.3, the ceiling on c_{bm} is 0.370 under compound symmetry with a constant cross-factor block. It is 0.646 when only the within-factor form is changed to AR(1), and 0.670 when the cross-factor block is also made to decay, as in `covar` (verified, $\rho_c = 0.7$). An earlier version of this paragraph set 0.370 against 0.670 and attributed the whole gain to the within-factor form; the cross-factor change contributes about 0.02 of it. On this schedule the AR(1) form nearly doubles the moderation the covariance can carry. The gain is schedule- and ρ-dependent: on the paper 01 designs without carryover at $\rho = 0.7$ it is 0.09 to 0.25 with a constant cross-factor block, and at $\rho = 0.3$ compound symmetry admits the higher ceiling (`docs/34`, Sections 4.3 and 4.4).

We stress the qualification. This is a like-for-like comparison, both arms using the decaying coupling. It is not the `orig` against `covar` comparison of section 3.3, which mixes the two changes and reverses direction. Reporting either figure without saying which comparison it belongs to would be misleading, and an earlier draft of paper 01's appendix did exactly that.

4.4 What we do not claim

Compound symmetry is not an error. It is the standard implication of a random intercept, it is exchangeable and therefore indifferent to occasion ordering, and it has one parameter rather than a parameter plus a time metric. In a design with few and evenly spaced occasions the two forms nearly coincide. Our claim is that the Hybrid schedule is not such a design, and that its one-week and four-week gaps should not be treated alike.

5. The cross-factor form: a consequence, not an independent choice

Architecture B applies the same decay to the cross-factor off-diagonal. We would not defend this change on its own, and section 2.2.4 of paper 01 does not list it among the four differences it enumerates. It follows from section 4.

5.1 The incoherence of mixing the two

Suppose the within-factor block decays and the cross-factor block does not. At the Hybrid schedule's widest gap, with $\rho = 0.7$ and $c_{\times} = 0.1$:

- BR at week 4 with BR at week 20: $0.7^{16} = 0.003$
- BR at week 4 with PB at week 20: 0.1

The pharmacological response would be some thirty times more strongly related to a *different* latent factor sixteen weeks away than to *itself* sixteen weeks away. We can construct no account of a patient under which that is true.

5.2 The numerical signature

Incoherence of that kind shows up in the covariance. Applying the cross-factor decay without the AR(1) change lowers the ceiling on c_{bm} from 0.34 to 0.16, the worst of the sixteen configurations examined in the crossed decomposition of `docs/whitepaper-covar-vs-orig-dgp.md`, Section 5 (worst cell over the paper 01 designs at $\rho = 0.7$, under the 2024 vendored coupling, which leaves the first occasion uncoupled; with every occasion coupled the worst cell is 0.151, in OL+BDC, `docs/34`, Table 4.1). Applying it alongside AR(1) it contributes roughly +0.02 to +0.03 (verified, `docs/34`, Table 4.1: 0.591 to 0.619 in CO).

Read together, the two numbers say the cross-factor form is not a lever in its own right. It is nearly free when it agrees with the within-factor form and expensive when it does not.

5.3 Our position

We take the two rows together for consistency and claim nothing more for the second than that. A reader who rejects the argument of section 4 should reject this one too, and would then have compound symmetry throughout, which is at least coherent.

6. The carryover recursion: our correction, not architecture

The fourth row of the table is not a departure from the published implementation at all. It records a correction we made in September 2026 to `covar` and to the mean-moderation process, and did not make to `orig`.

All three processes originally adjusted the response mean at an off-drug occasion by multiplying the previous occasion's *already adjusted* mean by a decay factor keyed to *cumulative* time since discontinuation. Elapsed time is therefore

counted twice, and every off-drug occasion after the first in a run is over-decayed. At a one-week half-life the second off-drug occasion receives half the residual effect it should (derived: with occasions one and two weeks after discontinuation, the recursion applies 2^{-2} to a mean already decayed by 2^{-1}, giving 2^{-3} where 2^{-2} is intended).

The corrected form anchors the decay to the mean at discontinuation, so the cumulative factor is applied once. We left `orig` alone because the purpose of that arm is to describe the published code rather than to improve it.

Two consequences follow for anyone comparing the two processes. The first off-drug occasion of any run agrees under both forms, so divergence begins at the second, and designs with long uninterrupted washouts show it most. And a comparison of `covar` against `orig` now carries one difference of our making. We recommend restoring the recursion in `covar` for the headline comparison, so that the four architectural differences are isolated, and reporting the corrected version alongside.

Note added 2026-10-07. Paper 01 has since taken the other course: it applies the anchored form to the published code itself, as one of its listed code changes to `58b32a9` (paper 01, Appendix B.3), rather than restoring the recursion in `covar`. In all current processes the time on drug restarts at zero on re-exposure.

7. What the repository's later history does and does not settle

The repository was revised in 2024. Commit `8609f12` of 6 May 2024 replaced the step gate with a graded form, added a scale-factor constant to the carryover adjustment, and left the first measurement occasion unassigned.

The first of those changes points in the same direction as section 3. We do not offer it as corroboration. That commit is authored by Hendrickson but its message describes it as the present author's edits curated into working code, so the two lines of reasoning are not independent and it would be circular to cite one in support of the other. The argument of section 3 rests on identification and on nothing else.

We record the later history for a different reason. The vendored copy of the published source in this repository is taken from the 2024 head, not from `58b32a9`, and any re-run of the comparison arm must use the latter. (Since 2026-10, a copy of the `58b32a9` sources is held at `analysis/data/quick-sim/hendrickson-58b32a9/`.) A reader who reproduces the appendix worked examples against the current head will not obtain the published vectors, and will additionally find the first occasion silently unassigned.

8. The exact ceiling used above

The bound quoted in sections 3.3 and 4.3 is not a sweep. Dropping the baseline row and column, which are those of the identity and factor out, write M for the block of the response factors and $\tilde{r} = c_{bm} v$ for the biomarker coupling extended by zeros. Positive definiteness requires M positive definite and $1 - \tilde{r}^\top M^{-1} \tilde{r} > 0$, giving

$$c_{bm}^{*} = \left(v^\top M^{-1} v \right)^{-1/2}.$$

The ceiling depends on the correlation structure only through how the coupling pattern sits relative to M^{-1}, which is why the coupling change of section 3 moves it as much as the correlation change of section 4 does.

We note one further result, since it corrects a natural but mistaken intuition. Under compound symmetry the smallest eigenvalue of M is $(1 - \rho_c) - (c_1 - c_\times)$, which contains neither the number of occasions nor the number of factors (derived; 0.200 at the production parameters for $n = 4$ to 32, verified). The response block itself therefore does not become harder to sustain as occasions are added.

Correction (2026-10-07). An earlier version of this paragraph went on to say that compound symmetry does not become harder to sustain as occasions are added, and that paper 01's draft appendix was wrong to attribute the narrower feasible range to the number of occasions. That does not follow. The ceiling depends on $v^\top M^{-1} v$, not on the smallest eigenvalue of M, and under compound symmetry it falls with the number of occasions at a fixed on-drug share, roughly as $n^{-1/2}$: 0.480, 0.346, 0.247 and 0.176 for $n = 4$, 8, 16 and 32 with half the occasions on drug (verified; closed form in `docs/34`, Section 5.1). What the eigenvalue result does show is that the loss comes through the coupling vector, not through the response block.

9. Conclusions

In conclusion, four points are to be emphasized.

First, the coupling vector change is a repair and not a preference. The published gate flattens the interaction channel under carryover, so that at any positive half-life the Hybrid design carries no on-drug versus off-drug contrast at all. An interaction that cannot vary with treatment state is not an interaction.

Second, the within-factor change is a modeling choice, and we would defend it while acknowledging that compound symmetry is defensible. The Hybrid schedule mixes one-week and four-week gaps, and a form that treats them alike discards the temporal structure the design exists to exploit. Holding the coupling fixed, the change nearly doubles the feasible range of the moderation parameter

on the evenly spaced schedule of section 3.3 at $\rho = 0.7$; the gain is smaller on the paper 01 designs and reverses at small ρ (section 4.3).

Third, the cross-factor change is a consequence of the second and should be presented as such. On its own it lowers the feasible range substantially, which is the numerical signature of asserting that a factor is more closely related to a different factor than to itself at the same remove.

Fourth, the carryover recursion is our correction rather than an architectural difference, and a comparison intended to isolate the architectural differences should hold it fixed across arms.

References

1. Hendrickson RC, Thomas RG, Schork NJ, Raskind MA. Optimizing aggregated N-of-1 trial designs for predictive biomarker validation: statistical methods and theoretical findings. *Frontiers in Digital Health* 2020;2:13. doi:10.3389/fdgth.2020.00013. (Verified via Crossref, 2026-10-07.) Code: `github.com/rchendrickson/pmsimstats`, commit `58b32a9`.